

Table 8.06: Fishergate comparisons by sceatta variety

<u>Sceatta mix</u>	<u>Period</u>	<u>Series/type</u>	<u>Fishergate</u>	<u>Hamwic</u>	<u>Rendlesham</u>	<u>Heckington</u>	<u>Rovston</u>	<u>Sledmere</u>
			Sub- total	Sub- total	Sub- total	Sub- total	Sub- total	Sub- total
Pre-primary	3c	Pa			1			
	3c	Va	0	0	1	0	0	0
Primary	4a	Y		1	1			1
	4b	A			9	2		
	4b	BX			2			
	4b	B	1		18	4	1	2
	4b	C		1	12	6	4	
	4b	F			3	2	2	
	4b	Vernus					1	
	4b	Saraoldo				2		1
	4b	W		3		2		
	4b	Z					2	4
	4b	Æthiliræd		1				3
	4b	?	1	5	1	45	18	10
	4b							10
Primary - Continental	4c	D	2	1	24	11	13	3
	4c	E	1	5	16	6	12	6
Secondary - Continental	4c	Stepped cross	3	6	40	17	5	3
	5c	E	7	11	10	18	9	12
	5c	G	2	1		3		21
Secondary - local	5c	Herstal			2			
	5c	X	9	9	21	13	2	23
	5b	H		48				
Secondary - other	5b	Y	7					13
	5b	R	7	48	6	6	0	0
	5a	J	5	5		1		28
Secondary - exotic species	5b	BIIIA			1		3	1
	5b	K		2		2	28	1
	5b	L		2	2	1	12	
	5b	M		1			1	
	5b	N+		2	1	1	1	
	5b	O		2			3	3
	5b	Q			6	1	1	8
	5b	QR			3	1		
	5b	R	1	1			3	8
	5b	S					9	
	5b	T					3	
	5b	U		5	1		3	1
	5b	V		1			1	
	5b	?		2				
	5b	Fleeing biped			1			
	5b	Animal mask					1	2
	5b	Celtic cross					1	
	5b	Hen			1			
	5b	Monitascorum					1	
	5b	Victory		1			1	1
	5b	Rosette						1
	5b	Saltire					1	2
	5b	Triquetra	1	1				
	5b	Woden	7	1	26	1	8	73
			27	107	122	66	124	125
<u>Sources:</u>			<i>Database 1</i>	EMC	EMC	EMC	EMC	<i>SiEMC 2</i>