

CANINE FAECES: THE MICROBIOLOGY OF AN ENVIRONMENTAL HEALTH PROBLEM

Salah Attia Jaber

BSc., Omar Al-Mukhtar, El-Beida, Libya

MSc., Omar Al-Mukhtar, El-Beida, Libya

MPhil., University of Sheffield, England



Thesis submitted in partial fulfilment for the requirements of the Degree of Doctor of Philosophy in the Department of Molecular Biology and Biotechnology, University of Sheffield, UK.

September 2012

Dedication

I dedicate this thesis to my family, especially to my mother and to my father's soul. To my wonderful wife Maryam Al-Magasbi, who supported me during every step of the way. To my children Azizza, Attia and Mohammed, thank you for patience and understanding. Finally, to all my teachers, my brothers, my sisters and my friends and my neighbours; I respect you all and, I dedicate my modest work to you.

Acknowledgments

First and foremost, I would like to thank Almighty Allah for giving me the patience and determination to carry out and conclude this Thesis. This research project would not have been possible without the support of many people. I wish to express my gratitude to my wonderful supervisor, Prof. Milton Wainwright who was extremely helpful and offered invaluable assistance, support, comments, encouragement and guidance. Deepest gratitude is also due to my advisor, Dr Jim Gilmour and Dr. Lynda Partridge for their valuable advice and suggestion during annual meetings. Not forgetting the members of the supervisory committee, whose suggestion and criticism was highly significant in accomplishing this thesis. I express my appreciation to cultural attaché of Libyan embassy, namely Dr. Abdel- Basit Gadour for tremendous support and encouragement. Dr Sulaiman Al-Naimat, Dr Khalid Al-Abri, Dr Adham Elshawaihde, Krys Bangret and Dr Rada all deserve special credit for their good company and assistance.

The author would also like to convey thanks to my tutor, Dr. James Foley from ELTC for encouragement and help, Words of credits should also go to Mr. Chris Hill for his assistance in electron microscopy also to Mr. Simon J Thorpe for his assistance in GC/MS experiments.

Finally, an honourable mention goes to my family and friends for their understandings and supports on me in completing this project. Special thanks also to all my friends in Prof Wainwright's laboratory, particularly for my best friends Sultan Al-sharari, Khalid Al-Quthami, Sami Alyahya, Waleed Al- Waneen and Mamdouh Bukari.

Abstract

The overall aim of the research work reported in this Thesis was to study a variety of aspects of dog faeces in relation to public health, their fertilizer potential and possibility that such faeces might be remediated using larvae, ultimately to provide a source of biodiesel. The results can be summarized as follows:

- 1) Dog faeces were shown to be source of pathogenic bacteria, notably *Escherichia coli* and *Salmonella*. These bacteria were shown to be transferred to the soil of a local playing field by direct, *in situ*, transfer from dog faeces undergoing weathering. *E. coli* and *Salmonella enterica* were isolated from all four sites while no such isolates were obtained from the fifth location which was uncontaminated with dog faeces
- 2) It was shown here that “common or garden” slugs can transfer potentially pathogenic bacteria from dog faeces to lettuce.
- 3) The feeding of Black Soldier Fly Larvae on faeces led to a statistically significant increase in the number of bacteria inside the BSFL gut and the same trend was seen in relation to dog faeces fed Fruit Beetle Larvae. This trend of increasing bacterial numbers in larvae fed on dog faeces is particularly worrying in relation to the potential feeding of these larvae to animals- post exposure to faeces.
- 4) Dog faeces were shown to have potential inherent fertilizer content; the nutrients present being released over a time period mimicking the natural weathering of dog faeces in the environment.
- 5) As a generalization, the addition of both types of larvae to dog faeces significantly reduced the concentration of indigenous plant nutrients over the entire four week incubation period; exceptions to this were nitrate and phosphate concentrations in BSFL treated faeces, where significant increases were seen at week 4 and 3 respectively and in faeces treated with FBL, where ammonium concentrations were significantly increased at weeks 2-4, and phosphate at week 4. While the addition of both larvae therefore

initially decreased levels of indigenous plant nutrients there was a trend in some of the nutrients to increase the longer the incubation went on. This suggests that perhaps a longer term exposure of dog faeces to the two larvae might have led to increase in ammonium, nitrate, sulphate and phosphate concentrations. The addition of ammonium, elemental sulphur and insoluble phosphate to dog faeces which had been modified by the two larvae led to significant increases in nitrate, sulphate and plant-available phosphate, results which show that dog faeces contains the indigenous microflora required for the transformation of these amendments (which simulate fertilizer addition). The results suggest the possibility that larval modified dog faeces could be used as compost additive fertilizer, or perhaps even be used as an agricultural soil fertilizer.

6) The potential for using fly larvae for the bioremediation of dog faeces was investigated. Black Soldier Fly (BSFL) and Fruit Beetle (FBL) Fly larvae were shown to dramatically improve the physical nature of canine faeces, even after only a short exposure period, giving a bioremediated product which is markedly improved in terms of texture, reduced odour and overall reduced offensiveness. The bioremediated dog faeces product was also found to be suitable as potting compost when “diluted” with proprietary potting compost.

7) The haemolymph and total body extracts of BSFL and FBL were shown to be antibacterial.

8) The potential for using dog faeces and dog faeces which had been treated with BSFL and FB as a source of biodiesel was determined. It was shown that potential biodiesel precursors (mainly fatty acids) were present both in the raw dog faeces and in faeces which were treated with the two different larvae.

9) The number of bacteria present in dog faeces disposed of in plastic bags dramatically increased over exposure to the UK summer, when temperatures were recorded between 10-27⁰C.

Table of Contents

| | |
|---|------------|
| Dedication | I |
| Acknowledgments | II |
| Abstract | III |
| Table of Contents | V |
| List of Figures | X |
| List of Tables | XVI |
| Chapter One: Introduction | 1 |
| 1.1. The environmental problem of canine faeces..... | 2 |
| 1.2. Microbiology of dog faeces..... | 2 |
| 1.3. Parasites and diseases associated with canine faeces | 4 |
| 1.3.1. Toxocariasis | 4 |
| 1.3.2. Toxoplasma contamination of dog faeces..... | 5 |
| 1.3.3. Other protozoan livestock parasite disease related to dog faeces | 6 |
| 1.3.3.1. Neosporosis | 6 |
| 1.4. Chemical composition of dog faeces..... | 8 |
| 1.5. Transmission of pathogenic bacteria from dog faeces to human food..... | 10 |
| 1.6. Slugs and snails as intermediate host-vectors of pathogenic bacteria..... | 11 |
| 1.7. Waste composting | 12 |
| 1.8. Composting and recycling of dogs waste | 13 |
| 1.9. Bio-conversion of putrescent waste using BSFL | 14 |
| 1.9.1. The Black Soldier Fly life cycle | 15 |
| 1.9.2. Bioconversion of dog faeces using Black Soldier Fly larvae (BSFL) | 15 |
| 1.9.3. The Texas Experimental BSF Composting Research Programme | 16 |
| Over a period of one year, approximately 20% by weight of the fresh food waste is converted into fresh larvae. This food waste had an average dry matter content of 37%, and the pre-pupae has an average dry matter content of 44%, i.e., on a dry matter basis, the bioconversion of food waste situates at almost 24%. An input of 100 kg of food waste per day can be handled by three 6-foot bioconversion units. | 16 |
| 1.14. Nitrification | 19 |

| | |
|---|-----------|
| 1.14.1. The Nitrogen Cycle | 19 |
| 1.14.4. Nitrate reduction..... | 22 |
| 1.14.6. Assimilatory nitrate reduction..... | 23 |
| 1.15.1. Forms of S in soil | 26 |
| 1.15.5. S-Mineralization..... | 29 |
| 1.15.7. Sulphur oxidation..... | 29 |
| 1.15.10. The phototrophic sulphur bacteria | 33 |
| 1.17. Aims of the Work Reported in this Thesis | 35 |
| Chapter Two: Transfer of Bacteria by Invertebrate Slugs from Canine Faeces to Lettuce | 36 |
| 2.1. Introduction | 37 |
| 2.2. Materials and Methods | 39 |
| 2.2.1. Slug culturing | 39 |
| 2.2.2. Isolation of bacteria from dog faeces | 39 |
| 2.2.3. Isolation of bacteria from slugs..... | 40 |
| 2.2.4 Isolation of bacteria from lettuce | 42 |
| 2.2.5 Bacterial identification | 42 |
| 2.2.5.1 Identification of bacteria using Chromagar medium | 42 |
| 2.2.5.2. Extraction of Genomic DNA | 43 |
| 2.2.5.3. Agarose preparation | 44 |
| 2.2.5.4. PCR amplification..... | 44 |
| 2.2.5.5. Phylogenetic identification of unknown bacteria..... | 45 |
| 2.2. 6. <i>Mycoplasma</i> identification | 45 |
| 2.2.6.1. Samples description | 45 |
| before BLASTn NCIMB database | 46 |
| 2.2.6.2. Detection of <i>Mycoplasma</i> | 47 |
| 2.2.6.3 Test samples preparation for PCR..... | 47 |
| 2.2.6.4 PCR amplification..... | 47 |
| 2.2.6.5. Agarose gel electrophoresis | 49 |
| 2.2.7. Detection and identification of fungi | 49 |
| 2.2.7.3. PCR amplification of fungal gene..... | 50 |
| 2.2.8. Identification of unknown fungi..... | 51 |
| 2.3. Results and Discussion | 53 |
| 2.3.1. Transfer of bacteria by slugs to lettuce | 53 |

| | |
|--|-----------|
| 2.3.2. Bacteria isolated from dog faeces | 53 |
| 2.3.3. Bacteria isolated from slugs | 54 |
| 2.3.3.1. Bacteria isolated from control slugs..... | 54 |
| 2.3.3.2. Bacteria isolated from slugs fed on faeces | 55 |
| 2.3.3.4. Fungi isolated from dog faeces | 55 |
| 2.3.4. Transmission of bacteria from dog faeces to lettuce..... | 55 |
| 2.3.5. <i>Mycoplasma</i> detection..... | 59 |
| 2.3.5.1. Attempts to isolate <i>Mycoplasma</i> from various samples..... | 59 |
| 2.3.5.2. Comments on the detection of <i>Mycoplasma</i> using PCR procedure | 63 |
| Chapter Three: Reduction in Bacterial Numbers in Dog Faeces Using Black Soldier Fly and Fruit Beetle Larvae | 64 |
| 3.1 Introduction | 65 |
| 3.2. Materials and Methods | 67 |
| 3.2.1 Methods for bacterial isolation and identification | 67 |
| 3.2.2. Dog faeces and insects larvae collection..... | 68 |
| 3.2.3. Bacterial isolation from dog faeces..... | 69 |
| 3.2.4. Bacterial isolation from the larval gut..... | 69 |
| 3.2.5. Determination of pH | 69 |
| 3. 2. 6. Statistical analyses | 70 |
| 3.3. Results and Discussion | 72 |
| 3.3.1. Identification of bacteria associated with dog faeces and larvae | 72 |
| Chapter Four: Modification of Canine Faeces by Black Soldier Fly and Fruit Beetle Larvae in relation to Plant Nutrient Release and Potential Use of Modified Faeces as a Compost Additive | 81 |
| 4.1 Introduction | 82 |
| 4.2. Materials and Methods | 85 |
| 4.2.1. Sample collection | 85 |
| 4.2.1.1. Samples collection from canine waste | 85 |
| 4.2.1.2. Samples collected from treated dog faeces | 85 |
| 4.2.1.3. Effect of larvae on the nutrient content of dog faeces. | 85 |
| 4.2.3. Nutrient transformations in modified dog faeces amended with ammonium nitrate, sulphur and insoluble phosphate..... | 86 |
| 4.2.3.1. Determination of ammonium–N (NH ₄ ⁺ -N) in dog faeces..... | 86 |

| | |
|---|------------|
| 4.2.3.2. Determination of Nitrate – N (NO ₃ ⁻ -N)..... | 87 |
| 4.2.3.3. Determination of Sulphate-SO ₄ ²⁻ - S | 87 |
| 4.2.3.4. Determination of phosphate solubilisation PO ₄ ³⁻ -P | 87 |
| 4.2.4. Measurement of the pH of dog faeces samples..... | 88 |
| 4.2.5. Determination of the effects of modified dog faeces as a compost additive on plant seed germination and growth | 88 |
| 4.3. Results and Discussion | 89 |
| 4.3.1. Effect of larvae on the texture of dog faeces..... | 89 |
| 4.3.5. Determination of the effects of modified dog faeces as a compost additive on plant seed germination and growth | 96 |
| Chapter Five: The Antibacterial Activity of the Haemolymph and Whole Body Extracts from BSFL and FBL against Four Species of Bacteria | 99 |
| 5. 1 Introduction | 100 |
| 5.2. Materials and Methods | 101 |
| 5.2.1. Culture of larvae..... | 101 |
| 5.2.2. Antibacterial properties of BSF and FB larvae | 102 |
| 5.2.2.1. Collection of excretion/secretion (ES) from larval whole-body and haemolymph..... | 102 |
| 5.2.2.2. Antibacterial assay | 102 |
| 5.2.3. Time killing curves for testing the bacterial activity | 103 |
| 5.2.4. Bacterial culture | 104 |
| 5.2.5. Transmission Electron Microscopy of Bacteria treated with haemolymph or whole body secretions of the two larvae | 104 |
| 5.2.6. Statistical analysis | 105 |
| 5.3. Results and Discussion | 105 |
| Chapter Six: The Number of Bacteria in Grassed Soil of Public Parks Contaminated with Dog Faeces and Effect of Faeces-Volatiles on Plant Growth..... | 114 |
| 6. 1. Introduction | 115 |
| 6.2. Materials and methods..... | 116 |
| 6.2.1. Samples collection from public parks | 116 |
| 6.2.2. Effect of incubation of dog faeces in plastic bags left under ambient outside conditions | 116 |
| 6.2.3 The effect of gasses (odour) from dog faeces on plant growth..... | 117 |
| 6.3 Results and Discussion | 119 |

| | |
|---|------------|
| 6.3.1. Occurrence of <i>E. coli</i> and <i>S. enterica</i> in dog faeces contaminated grassed-soil..... | 119 |
| 6.3.2 Effect of leaving dog faeces in plastic bags on bacterial numbers | 120 |
| 6.3.3 The effect of the gasses (odour) released from dog faeces on the plant growth | 122 |
| Chapter Seven: Production of Biodiesel from Dog Waste by BSF and FB Larvae- A Potential Solution To The Dog Faeces Problem..... | 127 |
| 7.1. Introduction | 128 |
| 7.1.1. Fuels from wastes..... | 128 |
| 7.1.2. Anaerobic digestion to methane..... | 128 |
| 7.2. Materials and Methods | 133 |
| 7.2.1. Samples collection | 133 |
| 7. 2.2. Extraction of lipids..... | 133 |
| 7.3. Results and Discussion | 135 |
| Chapter Eight: Final Discussion | 146 |
| 8.1 Final Discussion | 147 |
| 8.2 Future work | 152 |
| References..... | 154 |
| Wright, E. P. (1982). The occurrence of <i>Campylobacter jejuni</i> in dog faeces from a public park. <i>Journal of Hygiene</i> 89 : 191-194..... | 175 |
| Appendices..... | 176 |
| Appendix One: Preparation of chemical solutions and reagents..... | 177 |
| Appendix Two: Identification of bacteria | 181 |
| Appendix Three: ANOVA tables | 197 |
| Appendix Four: Turbo mass spectrometer GC/MS..... | 221 |

List of Figures

| | |
|--|----|
| Fig 1. 1: Anticipated transfer pathways of <i>E. coli</i> O157 by slugs from an environmental source to vegetable crops. | 12 |
| Fig 1. 2: The Nitrogen Cycle | 25 |
| Fig 1. 3: The Sulphur Cycle..... | 27 |
| Fig. 2. 1: Bacterial colonies isolated from dog faeces using Chromagar medium | 40 |
| Fig. 2. 2: Section of digestive track of slug from where the samples were taken | 41 |
| Fig. 2. 3: Isolation of Bacteria from slugs: (a) colonies inside slugs, and (b) colonies from the outside of slugs. | 41 |
| Fig. 2. 4: Standard hyperladder I with 14 lanes indicates higher intensity bands, 1000 and 10,000 and each lane (5µl) provides 720ng of DNA (BIOLINE supplier). | 46 |
| Fig. 2. 5: FinchTV software that manually adjusts errors of consensus sequences..... | 46 |
| Fig. 2. 6: Fungi isolated from dog faeces grown on a) PDA medium b) Czapek Dox... | 52 |
| Fig. 2. 7: Purification of fungi speces; a) green mould; b) black mould on PDA medium | 52 |
| Fig. 2. 8: Extraction of genomic DNA of isolated bacterial species; (lane JS1) <i>Enterobacter amnigenus</i> ; (lane JS 2) <i>Acinetobacter sp</i> ; (lane JS3) <i>Comamonas sp</i> ; (lane JS 4) <i>Acinetobacter sp</i> (lane L); hyper ladder and(lane JS5) <i>Delftia sp</i> | 57 |
| Fig. 2. 9: PCR- 16Sr RNA, amplification products of four isolated bacterial species analyzed by electrophoresis in agarose gel lanes represent; (lane L); hyper ladder;(lane JS1) outside control slugs (Metallic blue); (lane JS 2) inside slugs fed on faeces (B); (lane JS3) outside slugs fed on faeces(B); (lane JS 4) outside slugs fed on faeces (C); (lane 5) contaminated lettuce(Pink-red)..... | 57 |

Fig. 2. 10: Extraction of genomic DNA of isolated bacterial species; (lane L); hyper ladder and (lane MR1) *Delftia sp* ; (lane MR 2) *Staphylococcus lentus*; (lane MR3) *E. coli* 58

Fig. 2. 11: PCR-16Sr RNA, amplification products of isolated bacterial species analyzed by electrophoresis in agarose gel lanes represent; (lane L); hyper ladder;(lane MR1), contaminated lettuce (Blue); (lane MR 2) Inside slugs fed on faeces (Metallic blue); (lane MR3), Inside slugs fed on faeces (Pink-.red) 58

Fig. 2. 12: Extraction of genomic DNA fungi species ; (lane L); hyper ladder and (lane 6 and 7) fungi isolated from dog faeces 60

Fig. 2. 13: PCR-18SrRNA, amplification products of fungal species analyzed by electrophoresis in agarose gel lanes represent; (lane L); hyper ladder; (lane5) *Trichoderma asperellum* (lane 6) no result..... 60

Fig. 2. 14: Polymerase chain reaction (PCR) detection of *Mycoplasma* species in Earth worm samples, EZ-PCR-*Mycoplasma* test analyzed by electrophoresis in 2% agarose gel; the lanes represent (lane M); 1kb hyper ladder, (lane 1); negative control(distilled water). ; (lane 2); earth worm sample; (lane 3); negative result; (lane 4) positive control 61

Fig. 2. 15: Polymerase chain reaction (PCR) detection of *Mycoplasma* species in Dog faeces and lettuce samples, EZ-PCR- *Mycoplasma* test analyzed by electrophoresis in 2% agarose gel; the lanes represent (lane L); 1kb hyper ladder, (lane 1); negative control (distilled water).; (lane 2); negative result; (lane 3); dog faeces samples ; (lane 4) control lettuce; (lane 5); contaminated lettuce 61

Fig. 2. 16: Polymerase chain reaction (PCR) detection of *Mycoplasma* species in Dog faeces and lettuce samples, EZ-PCR- *Mycoplasma* test analyzed by electrophoresis in 2% agarose gel; the lanes represent (lane L); 1kb hyper ladder, (lane 1); Outside control slug (negative result) (lane 2); Outside contaminated slug; (lane 3); Outside

| | |
|---|----|
| control snails (lane 4) Outside contaminated snails (lane 5); Inside control slugs; (lane 6); Inside contaminated slugs; (lane 7); inside control snails (lane 8); Inside contaminated snails (lane 9) negative control (distilled water) | 62 |
| 2); earth worm sample; (lane 3); negative result; (lane 4) positive control | 61 |
| Fig. 3. 1: Black Soldier Fly larvae, 1 st instar | 70 |
| Fig. 3. 2: A Black Soldier Fly adult..... | 70 |
| Fig. 3. 3: Fruit Beetle larvae 3 rd instar | 71 |
| Fig. 3. 4: Fruit Beetle pupae | 71 |
| Fig. 3. 5: Fruit Beetle adults | 71 |
| Fig. 3. 6: <i>Escherichia coli</i> grown on selective chromogenic agar <i>E. coli</i> media isolated from dog faeces and larvae fed on dog faeces; a) HiCrome <i>E. coli</i> agar showing purple/pink coloured colonies, b) Brilliance <i>E. coli</i> /Coliform agar showing blue colonies. | 73 |
| Fig. 3. 7: <i>Salmonella Spp</i> grown on XLT-4 selective <i>Salmonella</i> agar medium..... | 73 |
| Fig. 3. 8: Colonies of bacteria grown on plate count agar medium..... | 74 |
| Fig. 3. 9: Two bacterial isolates showing different external morphology; a) cocci, b) bacillus, observed by SEM..... | 74 |
| Fig. 3. 10: Bacterial numbers in dog faeces grown on; a) plate count medium, b)HiCore <i>E. coli</i> in and c) <i>Salmonella</i> medium after treatment with Black Soldier Fly larvae..... | 75 |
| Fig. 3. 11: Bacterial numbers from dog faeces, bacteria grown on; a) plate count medium, b) HiCore <i>E. coli</i> medium and c) <i>Salmonella</i> medium by Fruit Beetle Larvae. | 76 |
| Fig. 3. 12: Numbers of bacteria; a) <i>E. coli</i> , b) <i>Salmonella enterica</i> and bacterial species grown on plate count medium isolated from gut of BSF larvae | 78 |
| Fig. 3. 13: Numbers of total “heterotrophic bacteria” in gut of FB larvae; a) <i>E. coli</i> b) <i>Salmonella enterica</i> and c) bacterial species grown on plate count medium | 79 |

| | |
|--|----|
| Fig. 4. 1: Untreated dog faeces after 28 days incubation..... | 89 |
| Fig. 4. 2: Modified faeces: a) treated with FBL; | 89 |
| Fig.4. 3: The effect of BSFL activity on indigenous; a) ammonium, b) nitrate, c) sulphate and d) phosphate concentration in dog faeces(* Significantly different from control, p=0.05)..... | 90 |
| Fig. 4. 4: The effect of FBL activity on a indigenous a) ammonium, b) nitrate, c) sulphate and d) phosphate concentration in dog faeces (* Significantly different from control, p=0.05)..... | 91 |
| Fig. 4. 5: The effect of length of incubation on a) ammonium concentration following addition of (10µg/ (NH ₄) ₂ SO ₄ -N g) , b) nitrate concentration following the addition of (10µg/ (NH ₄) ₂ SO ₄ -N g), c) S ⁰ oxidation following the addition of (1% w/w) S ⁰ and d) phosphate solubilisation following the addition of (1% w/w)calcium phosphate to dog faeces modified with BSFL(* Significantly different from control, p=0.05) | 92 |
| Fig. 4. 6: The effect of length of incubation on a) ammonium concentration following the addition of (10µg/ (NH ₄) ₂ SO ₄ -N g), b) nitrate concentration following the addition of (10µg/ (NH ₄) ₂ SO ₄ -N g) , c) S ⁰ oxidation following the addition of (1% w/w) S ⁰ , d) | 93 |
| Fig. 4. 7: Shows a) the seeds planted in untreated dog faeces; from 1 to 3 - turnip seeds, from 4 to 6 is lettuce seeds; b) the seeds planted in compost ; 7 to 9 - turnip seeds, from 10 to 12 is lettuce seeds. | 97 |
| Fig. 4. 8: Shows; a) the lettuce seeds planted in different concentration of treated dog faeces with BSFL No; 1, control dog faeces 2, 3 and 4 are 75, 50 and 25% of dog faeces concentration and 5 is a control compost. From 6 to 10 are the turnip seeds planted at the same concentration b) the lettuce seeds planted in different concentration of treated dog faeces with FBL No; 1, control dog faeces 2, 3 and 4 were 75, 50 and 25% of dog | |

| | |
|--|-----|
| faeces and 5 is a control compost. From 6 to 10 showing turnip seeds planted at the same concentration..... | 97 |
| Fig. 5. 1: The effect of BSFL haemolymph and whole body extract on <i>E. coli</i> , growth curve over 8 h..... | 105 |
| Fig. 5. 2: The effect of FBL larvae haemolymph and whole body extract <i>E.coli</i> , growth curve over 8 h..... | 106 |
| Fig. 5. 3: The effect of BSFL haemolymph and whole body extract on <i>S.enterica</i> growth curve over 8 h. | 106 |
| Fig. 5. 4: The effect of FBL haemolymph and whole body extract on <i>S.enterica</i> growth curve over 8 h..... | 107 |
| Fig. 5. 5: The effect of BSFL haemolymph and whole body extract on <i>S.aureus</i> MRSA, growth curve over 8 h. | 108 |
| Fig. 5. 6: The effect of FBL haemolymph and whole body on <i>Staphylococcus aureus</i> MRSA, growth curve over 8 h. | 109 |
| Fig. 5. 7: The effect of BSFL haemolymph and whole body on <i>Serratia marcescens</i> , growth curve over 8 h. | 109 |
| Fig. 5. 8: The effect of FBL haemolymph and whole body extract on <i>Serratia marcescens</i> , growth curve over 8 h..... | 110 |
| Fig. 5. 9: The effect of larval components on the morphology of <i>E. coli</i> observed by TEM, a) control for BSFL, b) whole body extract of BSFL, c), haemolymph of BSFL; d) control for FBL, e) whole body extracts for FBL and g) haemolymph of FBL. | 112 |
| Fig. 5. 10: The effect of larval components on the morphology of <i>S.aureus</i> observed by TEM, a) control for BSFL, b) whole body extract of BSFL, c), haemolymph of BSFL; d) control for FBL, e) whole body extracts for FBL and g) haemolymph of FBL. | 112 |
| Fig. 6. 1: Plastic bags containing dog faeces bags obtained from Sheffield parks..... | 117 |

| | |
|---|-----|
| Fig. 6. 2: Airtight container contains 200 g of fresh dog faeces, the dog faeces gases were pumped via the tube into the plastic box containing compost planted with lettuce or turnip seeds. | 118 |
| Fig. 6. 3: The presence and population density of <i>E. coli</i> and <i>Salmonella sp</i> in five locations in public areas of Sheffield (location 5 is the uncontaminated grass sample). | 119 |
| Fig. 6. 4: Bacterial species isolated from dog faeces left in polythene bags; a) red colonies of <i>E. Coli</i> , and blue <i>Enterococcus Spp</i> , b) <i>E. coli</i> , c) <i>Salmonella enterica</i> , d) <i>Staphylococcus aureus</i> | 121 |
| Fig. 6. 5: Numbers of <i>E. coli</i> , <i>Salmonella enterica</i> and total heterotrophic bacteria (grown on plate count media) isolated from dog faeces left in plastic bags. | 122 |
| Fig. 6. 6: The effect of gas (odour) emission from dog faeces on vegetable growth; a) control lettuce seeds, b) control turnip seeds c) lettuce treated with dog faeces gasses and d) turnip treated with dog faeces gasses. | 124 |
| Fig. 6. 7: A typical dog waste bin used in public parks in the UK (author’s photograph). | 125 |
| Fig. 7. 1: Fatty acids extracted from; a, BSFL, b) FBL, | 135 |
| Fig. 7. 2: GC–MS chromatogram of a reference mixture of fatty acid representative the total of samples ion taken from dog faeces and BSF and FB larvae. The chromatogram for this figure and all subsequent figures were obtained with Gas Chromatograph using Zebron ZB-S capillary column | 141 |

List of Tables

| | |
|--|-----|
| Table 1. 1: Bacteria that can dissimilate nitrate to ammonium Based on Tiedje (1988) | 24 |
| Table 1. 2: Colourless sulphur oxidising bacteria. (Modified after Kuenen and Beudeker, 1982) | 32 |
| Table 2. 1: PCR reaction mixtures for the amplification of <i>Mycoplasma</i> -DNA fragment. | 48 |
| Table 2. 2: PCR amplification protocol of <i>Mycoplasma</i> DNA fragment | 48 |
| Table 2. 3: Analysis of coloured bacterial strains which require further 16S rRNA tests for their identification confirmation to the species level. | 54 |
| Table 2. 4: Bacterial identification using 16SrRNA | 56 |
| Table 2. 5: Presence of <i>Mycoplasma</i> in various samples | 59 |
| Table 3. 1: Media used to detect and identify <i>E. coli</i> and <i>S. enterica</i> from dog faeces, BSF and FB larvae gut | 72 |
| Table 4. 1: The percentage seed germination and growth planted in different ratios of in modified dog faeces (treated with BSF and FB larvae) and potting compost | 96 |
| Table 5. 1: The effect of the two species of larval haemolymph and whole body secretions on the four species of bacteria. | 111 |
| Table 7. 1: The percentage of lipids in dry weight of 1g samples of faeces and larvae | 136 |
| Table 7. 2: The GC/MS analysis chain composition of a number of fatty acids from dog faeces and larval fats | 139 |
| Table 7. 3: Comparison of fatty acid methyl ester composition of BSFL fat-based biodiesel and rapeseed oil-based biodiesel. | 143 |

| | |
|---|-----|
| Table 7. 4: Comparison of fuel properties of BSFL fat-based biodiesel, rapeseed oil-based biodiesel, and the standard EN14214..... | 144 |
|---|-----|



Chapter One: Introduction



1.1. The environmental problem of canine faeces

The Dog (*Canis lupus familiaris*) is a member of the Canidae family of the mammalian order "Carnivora. Dogs have been adapted for economic usefulness and are loyal and generally protective of humans, being used as guide dogs for the blind and disabled; their keen sense of smell is also used to detect bombs or drugs (Kim, 2008; Murray, 2007). In contrast, dogs are a source of danger to humans through their bites, and because they transmit zoonotic diseases such as rabies, toxoplasmosis, echinococcosis, trypanosomiasis, filariasis, spirocerosis, hydatidosis, larva migrans etc. (WHO, 1959; Oduyemi and Olayemi, 1977; Hill *et al.*, 1985) resulting in death. Canine waste which is not removed from the local environment due to the irresponsible behaviour of dog owners may represent a source of potential pathogens. Pathogenic bacteria can survive in canine faeces for a long period and can be spread by wind and vehicular traffic. Faeces can also be carried inside dwellings via contaminated shoes (Tarsitano *et al.*, 2010). The interactions between plants and other animals including invertebrates in the field may also be an important means of transmission of pathogenic bacteria to crops.

Contamination of the urban environment by dog faeces continues to be a growing problem around the world. While in the UK societal changes in thinking about the dog dirt problem has led more people to pick up the waste products of their pets, our streets continue to be polluted with dog faeces.

1.2. Microbiology of dog faeces

Dog faeces present two major problems. Firstly they smell and are generally offensive, secondly, and more importantly, they spread microbial and parasitic diseases. Since dog faeces often contaminate parks and playing fields, children are likely to be

frequently exposed to the likelihood of catching such damaging and potentially fatal infections. The disease spread to humans by dog faeces include:

Campylobacteriosis and Yersiniosis, dogs which eat raw pork are infected with *Yersinia enterocolitica* which can cause enteritis in dogs and cats; the organism is also shed in the faeces for several weeks after infection, even in the absence of clear symptoms (Fredriksson-Ahomaa, *et al.*, 2001), Dogs may therefore be an obvious source of human infection and this has shown to be high amongst children under six years of age, so dog faeces may provide a transmission link for pathogenic bacteria between pigs and young children. Campylobacteriosis is a bacterial infection causing diarrhoea in humans. Wright *et al.* (1982) found that for dog faeces collected in urban parks *Campylobacter* were isolated from 260 collected samples, while *Salmonella* species were found in only three. Most of the *Campylobacter* were isolated during the warm months of June and July, i.e. they were present when children are most likely to be at play outside.

Salmonellosis, represent a very large group of rod-shaped, gram negative bacteria including more than 2000 known serotypes which belong to the family of *Enterobacteriaceae*. All these serotypes are human pathogens and can cause various symptoms from mild gastroenteritis to severe illness or death. In 1890 more than 30 people out of every 100,000 in the United State died of typhoid caused by *S. typhi*. *Salmonella* can cause food poisoning from eggs, pork, chicken and beef (Jacquelyn, 1999; Blancou *et al.*, 2005) and is the most common bacterial infection transmitted to humans, symptoms include, fever, muscle aches and vomiting and diarrhoea.

Escherichia coli is a gram-negative rod of the family *Enterobacteriaceae* and is found in the gastrointestinal of all warm-blooded animals. All strains of *E. coli* are spread by the faecal-oral transmission route. Many strains can cause gastroenteritis; among these

are the enterotoxigenic (ETEC), enteropathogenic (EPEC), enteroinvasive (EIEL), or enterohemorrhagic (EHEC) *E. coli*. (Maier *et al.*, 2009).

Shigella flexneri is an intestinal bacterium responsible for severe diarrhoea in people and non-human primates (monkeys); it is also found in dogs (Wang *et al.* 1996). About 25,000 cases are reported in people in the United States every year. Many monkeys carry this bacterium without symptoms, while people are also commonly carriers. Small children at child care facilities and people who handle monkeys are most at risk. The diarrhoea produced in people exposed to human or monkey waste is never life threatening but during the two to three weeks the diarrhoea lasts, the victim is quite ill. The chief danger from this disease is dehydration.

Streptococcus and Staphylococci these bacteria are found on all animals that typically associate with humans. In the great majority of cases they cause no disease in the animal. However, in pets, eye infections are occasionally associated with Streptococci and skin infections with Staphylococci. Both bacteria can spread from pets to humans on contaminated hands and objects. Infections are generally limited to the skin and eyes.

Faecal coliforms cause dysentery herpes, typhoid fever and ear infections in humans. Not surprisingly these bacteria are well represented in dog and other animal faeces (Whitlock *et al.*, 2002).

1.3. Parasites and diseases associated with canine faeces

1.3.1. Toxocariasis

Toxocariasis is the main medical problem relating to dog faeces. It is an infection of the round worm *Toxocara canis* (Gillespie, 1988) and is a zoonotic disease spread via unwashed vegetables and dog faeces (Karadam *et al.*, 2008); young people are particularly at risk due to their weaker immune systems and because of their likely increased exposure by ingesting the eggs (Thompson *et al.*, 1986). Puppies, which are a

major source of environmental contamination, can pass up to 15,000 eggs per gram of faeces. Each *T. canis* female can lay up to 700 eggs a day and these are excreted when the dog defecates; they can survive in soil for as long as three years. After two to three weeks of warm weather the eggs develop into an embryo state, containing larvae which are infective to dogs and people. The larvae attempt to migrate throughout the human body like they would do in a dog's, but the human body regards them as foreign and reacts leading to tissue damage (Overgaauw and Nederland, 1997). Two types of toxocariasis exist, namely: visceral larva migrans (VLM) and ocular larva migrans (OLM). In VLM, the larvae reach the liver, causing inflammation and symptoms including abdominal pain and pyrexia; most people however, recover spontaneously. OLM occurs when a migrating larva reaches the eye where it causes the formation of a granuloma on the retina which leads to significant visual impairment and in severe cases, blindness; around 12 new cases of OLM are diagnosed annually in the UK (Despommier, 2003).

1.3.2. Toxoplasma contamination of dog faeces

Toxoplasma gondii is an obligate intracellular protozoan with worldwide distribution (Frenkel 1990, Wallace, 1973), where it can cause blindness in humans. It is found in the Americas, including southern Mexico, Central America, South America, and the West Indies. Infections with *Toxoplasma* are very common in Panama, although most infections are asymptomatic. Antibodies titres to *Toxoplasma* are high in children, notably where cats and dogs are numerous, sanitation is bad and there is high shade and high humidity. Cats are the main hosts of *Toxoplasma*, but the parasite is also found in dogs (Barutzki and Schaper, 2003). Pets become infected by consuming rodents and birds, which are intermediate hosts that contain cysts (bradyzoites) that help to continue the chronic infection. When pets eat infected animals, the bradyzoites develop into the

enteroepithelial stages and the shedding of oocysts in the cat or dog faeces. The oocysts then sporulate in the soil and then lie dormant from up to weeks or months, especially when in moist, shaded areas. Sporozoites within the oocysts are then infectious to humans and other mammals after being passed by hand to mouth, after which tachyzoites and bradyzoites (multiplying asexual stage) continue to induce active infection. Dogs have only recently been considered a factor in the spread of toxoplasmosis (Frenkel, 1996). Interestingly, by eating or rolling in cat faeces, dogs probably play an important role in the mechanical transmission of *Toxoplasma* oocysts (Lindsay *et al.*, 1997) and it is likely that dogs excrete oocysts near human habitats, promoting *Toxoplasma* transmission to humans.

1.3.3. Other protozoan livestock parasite disease related to dog faeces

There is a growing link between two specific protozoan diseases in livestock and the fact that faeces, from infected dogs, is increasingly found on grazing land (Dubey and Lindsay, 2006). The two main diseases in question are:

1.3.3.1. Neosporosis

This disease caused by the parasite *Neospora caninum* is responsible for the highest rate of all cattle abortions reported in the UK. Once this disease occurs in cattle it can remain in the herd as a result of vertical transmission of the parasite between cows and their calves. *Neospora* eggs are produced by infected dogs and then excreted into their faeces. Cattle then become infected when they eat food or drink water contaminated with the eggs. Infection in cattle is common and generally ill effects are not clear, either for the cow or the calf (Williams *et al.*, 2000) The disease becomes obvious when *Neospora* multiplies in the cells of the developing calf and its placenta and causes damage sufficient to bring about abortion or stillbirth. Control of *Neospora* abortion is difficult and there are no drugs available at present to control this disease in cattle or to

cure the infected animals. Similarly, no vaccine is currently licensed in the UK to prevent cattle-neosporosis. Fortunately, current evidence shows that *Neospora* is not a major problem for humans . (Anderson *et al.*, 1995)

The vertical transmission of neosporosis is a main cause of long standing infection within a herd, although spread of the disease between females which are not related only occurs where a dog acts as host to the parasite. The parasite can be picked up by dogs through the consumption of contaminated livestock material, including placentas from newly calved cows, or by being fed contaminated raw meat, faeces from infected dogs then contaminate pasture and also cattle feed, water or bedding (Anderson *et al.*, 1997); (Davison *et al.*, 1999).

Only a small number of infected dogs develop the disease, which produces progressive lameness and paralysis in pups younger than 6 months of age. Infected bitches can pass the parasite to their young during pregnancy by transplacental infection. If dogs do develop symptoms, then the results are usually fatal or lead to euthanasia (Reichel *et al.*, 2007). This disease is very important since it impacts farm economics due to infected cows being more likely to abort and the occurrence of premature culling and reduced milk yields. Since there is no way to prevent (through vaccination), or an effective treatment of neosporosis, a farmer's main defence against the disease is to take action against any likely *Neospora* contamination (Dubey, 2003).

1.3.3.2. Sarcocystosis

This is a disease which is also caused by a parasite, in this case *Sarcocystis* spp, which employs a number of intermediate hosts, including dogs. *Sarcocystis* eggs are produced by infected carnivores and are excreted in their faeces, and sheep become infected when they eat food, or drink water contaminated with *Sarcocystis* eggs. In many cases, infected livestock show no disease symptoms (Traub *et al.*, 2002). The

disease can be transmitted from ewe to lamb during pregnancy, but vertical transmission is not believed to be important. Dogs can pick up the parasite through the ingestion of contaminated material from carcasses, or by consuming contaminated raw sheep meat. Faeces from infected dogs can also contaminate pasture as well as animal feed, water or bedding. In contrast to neosporosis, no transmission of the *Sarcocystis* parasite occurs between bitch and puppy. The link between infected dogs and sarcocystosis in sheep is compelling, but the disease is generally regarded as less of a problem than neosporosis. No vaccine is available against sarcocystosis in sheep and although there are some treatments available, the high cost and practicality of administration of these prevents their spread use. As with neosporosis, the most feasible option for the farmer is to reduce infection risks. In addition to, round worms, giardiasis, tuberculosis, gastroenteritis and cryptosporidiosis and Cystercercosis, is a human disease involving larval tapeworms. Although parasitic infections are important in relation to dog faeces, no attempt was made here in this thesis to study these infectious agents . (Dubey and Williams, 1980)

1.4. Chemical composition of dog faeces

Dog faeces (and those of cats) contains about 0.7% nitrogen, 0.25% phosphate and 0.02% potasium. As a result, dog faeces are not a particularly good plant fertilizer; they are offensive and often oderous and contain pathogens, as well as *Toxoplasma*. In it its unweathered state therefore dog faeces are not a useful, let alone, ideal organic fertilizer.

Currently most dog waste is allowed to breakdown naturally in the environment where it is deposited, and where collected it is usually incinerated. Dog faeces are not ideal additives to composting plants and as a result, their addition to municipal composters is generally avoided. Similarly, because of their low nutrient and high

pathogen content they are not regarded as ideal or safe fertilizers for agricultural, garden or allotment use.

The problems relating to dog faeces and the environmental pollution they cause has been largely overlooked and surprisingly little research work has been published in this area.

Here are some facts about dog ownership and the resultant waste problem relating to the UK: There are around 24 million UK households and, in 2002, the number of households owning dogs was 4.8 million. Some 21% of households with dogs have more than one. There are around 6.8 million dogs in the UK, the highest levels of dog ownership being among the 45 to 54 year-old age group - around 30%. It has been calculated that the UK dog population produces some 900 tonnes of faeces every day and over a ten-year lifetime, a dog can produce up to half a ton of faeces.

There are an estimated 41 million and 60.7 million dogs in Europe and the USA respectively. Australia has one of the highest rates of pet ownership in the world, with almost 36% of Australian households having a dog (Australian Companion Animal Council 2010).

In the UK, the legal position relating to dog fouling is covered by the following statutes:

The Dog (Fouling of Land) Act 1996 in England and Wales. The Dog Fouling (Scotland) Act 2003. The Litter (Animal Droppings) Order 1991 - Made under section 84 (14) of the Environmental Protection Act 1990 and Statutory Instruments Numbers 2762 and 2763. DoE Circular No 18/96 (Welsh Office No 54/96).

These Acts require that the owner should immediately clean up after his or her dog, should it foul what is termed 'designated land'. Designated areas are usually defined as places where dog faeces have the potential to cause a health hazard to people, including children's play parks, public greens and parks, residential areas, cycle paths and

walkways etc. Individual local authorities can use these Acts as a basis to create bylaws (which allow for instant fixed-penalty fines) and nominate the designated areas in the appropriate borough. The fines begin at around £40, rising to a maximum of £1,000. Dog wardens may be employed to patrol these areas and catch those irresponsible dog owners who fail to remove their dog's faeces.

The Government recommends that the dog faeces-related disease problem can be reduced by the following actions:

- a) Poop scooping on each occasion your dog makes a mess.
- b) Dogs should be wormed regularly-every three to six months, using a wormer recommended by a veterinary surgeon.
- c) Dogs should be exercised in dedicated areas of parks where available.
- d) Dogs should be discouraged from parks having children's playgrounds.
- e) Pregnant women and individuals who suffer from impaired immunity should use additional extra precautions when cleaning up the faeces, for example by wearing disposable gloves.

1.5. Transmission of pathogenic bacteria from dog faeces to human food

Because of immunization and the use of litter boxes and flea treatments, the transfer of pet diseases to humans has greatly been reduced over the years, but still some dog related diseases infect people; most troubling of all when people are infected by their pet, they usually are unaware of it. Besides the risk of bites, scratches and allergies, several infections can be transmitted to the human as zoonosis. These pathogens have an oral-faecal transmission cycle and humans can be infected either by faecal contamination of food, water or the environment (gardens, sandpits and playgrounds) or by direct contact (Overgaauw *et al.*, 2009). Outbreaks of diseases caused by infective bacteria as well as parasites have been documented to occur as a result of consumption

of contaminated salad and vegetables. The studies described in this Thesis relating to this problem focus in the potential consumption of contaminated lettuce. Sources of pre-harvest contamination of produce include manure, from livestock operations, and domestic animals (Beuchat, 2006).

Diseases caused by pathogenic bacteria, such as Salmonellosis and Campylobacteriosis are the most frequently reported zoonotic diseases transmitted from animals to humans via food (Norrung and Buncic, 2007). Infections with verotoxin-producing *E. coli* O157 are comparably less frequent but of considerable public health concern as they are associated with life-threatening human diseases such as haemorrhagic colitis (HC) and haemolytic uraemic syndrome (HUS) (Roldgaard *et al.*, 2004). Although a variety of foods may serve as vectors of food borne illness, the farm is the original source of all of these pathogens and there is strong association between prevalence in food production and other animals and post-harvest prevalence on carcasses (Elder *et al.*, 2000).

1.6. Slugs and snails as intermediate host-vectors of pathogenic bacteria

Slugs (Mollusca: Gastropoda) are a major pest of fruits and vegetables (Godan, 1983; South, 1992), with the Grey Garden Slug *Deroceras reticulatum* (Stylommatophora Agriolimacidae) being the most widespread and most serious pest (Wilson *et al.*, 1993). Slugs, by serving as intermediate hosts for many bacterial diseases mainly found in animals faeces (South, 1992), present a risk to plants, animals and humans. Slugs ingest soil bacteria during feeding and as a result, they become hosts of many bacteria including pathogens like such as *E. coli* O157. These can be picked up from dog faeces or from animal faeces which have been spread in open fields (Figure 1.1) (Walker *et al.*, 1999).

Emma *et al.* (2006) conducted research on the Yellow Slug (*Limax flavus*) and the Great Gray Slug (*Limax maximus*), and confirmed that both species can carry *E. coli*

O157, both on the surface and internally. The persistent slug species *Deroceras reticulatum* can carry *E. coli* on its external surface for 14 days (Emma *et al.*, 2006). Slugs which ingest *E. coli* also pass viable bacteria to their faeces and *E. coli* was found to persist for more than 3 weeks in excreted slug faeces (Hogan, 1985). In a similar study Emma *et al.* (2006) showed that *E. coli*, through contact and/or ingestion can survive for many days both internally and externally.




Fig 1. 1: Anticipated transfer pathways of *E. coli* O157 by slugs from an environmental source to vegetable crops.

1.7. Waste composting

Increases in the human population and the expansion of large cities have lead to marked increase in the volume of all kinds of waste. The search for modern approaches to waste management, notably composting has recently gained momentum (Pascual *et al.* 1997; Bhattacharyya *et al.* 2001a and b; Smith and Hughes 2004). Composting is now often a preferred approach to waste management (Lee *et al.* 2004; Sharholly *et al.* 2008) and is defined as the *biological oxidative decomposition of organic matter*

(Stoffella and Kahn 2001) based on the catalytic activity of environmental organisms which are responsible for organic matter decomposition. Under optimum conditions, three stages of traditional composting have been identified: 1) mesophilic, or moderate temperature phase; 2) thermophilic, or high temperature phase and 3) cooling or maturing phase (Kostov *et al.* 1996; Trautmann and Olynciw 2000). The duration of these phases relates to the type of organic matter under compost and its efficiency, which is largely determined by aeration and humidity (McKinley and Vestal, 1985; Strom 1985, Strom *et al.* 1983; Butler *et al.* 2001).

1.8. Composting and recycling of dogs waste

The risk of canine waste accumulation in the urban environment and agro-ecosystem is an increasing problem. Recycling is a sustainable approach for disposing of waste, and composting can be an important component of recycling approaches. The microbes involved oxidize carbon as an energy source for growth and take in nitrogen for protein synthesis (Taylor, 2004). The correct carbon to nitrogen ratio in composting systems is required for the efficient decomposition of wet dog waste contains 0.7% nitrogen (N), 0.25% phosphate, compared to wet cattle manure, dog waste which contains 40% more nitrogen, the same amount of phosphate, and a twentieth of the amount of potash (Hall and Schulte 1979). Dog waste composting reduces the amount of waste being sent to landfills (Sequi, 1996) and also reduces the amount of methane being released into the atmosphere (Peigne and Girardin 2004, Albaladejo *et al.* 2000). Because of the high nitrogen content of dog manure, a rich source of carbon is required for composting, such as wood chips, shavings or sawdust (C:N ratio of 560-641:1) (Rynk 1992, Miller 1996).

1.9. Bio-conversion of putrescent waste using BSFL

After seven years of research, a patented bioconversion process that effects a 95% reduction in the weight and volume of food waste within a matter of just a few hours has been developed using the Black Soldier Fly. This unique approach to bioconversion requires no energy, no electricity, no chemicals, not even the addition of water. It is totally self-contained and does not produce effluent and, while it produces a small amount of carbon dioxide, it does not produce any other greenhouse gases, such as methane (Craig Sheppard *et al.*, 2002).

The Black Soldier Fly BSF (*Hermetia illucens*) is a tropical fly (Craig Sheppard *et al.*, 2002) indigenous to the whole of the Americas, from the south of Argentina to Boston and Seattle, and in World War II, the fly also spread into Europe, India, Asia as well as Australia.

As a result, the currently discussed bioconversion process does not require the introduction of a foreign or exotic species, involving as it does an organism which is indigenous to the Americas, and now Europe. It is also not associated with the transmission of disease. The BSF has the ability to thrive in the presence of salts, alcohols, ammonia and a various food-based toxins and can process food waste as well as swine, human and poultry waste. Upon reaching maturity, the larvae of the BSF migrate out of the bioprocessing unit into a collection bucket without any human or mechanical intervention and thereby provide a self-harvested grub which is rich in nutrients and is food source which rivals in commercial value the finest fish meal. Unlike many other flies, BSF adults do not enter dwellings and since they do not have functional mouth parts, they do not eat waste and cannot (like the House Fly) regurgitate on human food, and therefore are not involved with disease transmission. Black Soldier Fly adults do not, in any way, bite, bother or annoy humans.

1.9.1. The Black Soldier Fly life cycle

Soldier fly adults come together in small numbers near an isolated bush or tree in order to find and choose a mate. After mating, the female finds an ideal place to lay her eggs, and lays about 900 eggs in 5 to 8 days lifespan. Housefly adults, by contrast, often live for 30 days, and during this long period, they eat, and as result actively spread disease.

Male BSF do not go near wastes since they do not lay eggs. The females in fact, prefer to lay their eggs not upon the waste, but either above or to the side of it, thereby allowing the eggs a far better chance of survival. The eggs are relatively slow in hatching (102 to 105 hours). The newly hatched larvae then crawl or fall onto the waste and begin to eat it with unbelievable rapidity. It takes about two weeks for the larvae to become mature a period which may extend to 6 months if the temperature is not right, or if there is not enough food. This ability of BSF larvae to extend its life cycle under conditions of stress is very useful when using it waste bioprocessing.

Black Soldier Fly larvae pass through 5 stages or instars. When mature, pre-pupal larvae are about 25mm long, 6mm in diameter, and weigh around 0.2 grams. These larvae are extremely tough and robust and can survive under conditions of extreme oxygen starvation. They can be also subjected to several 1000 gs of centrifugation without being harmed (Lord *et al.*, 1994).

1.9.2. Bioconversion of dog faeces using Black Soldier Fly larvae (BSFL)

Black Soldier Fly larvae have been used to dramatically reduce food waste and manure, and convert the nutrients from food waste and manure into insect larval biomass containing over 40% protein and over 30% crude fat (Newton *et al.* 2005). As a result, BSF larvae are potentially excellent source of protein and therefore as a high energy meal for chicken, fish and other domestic animals. As an added benefit,

BSF larvae also aerate and dry manure, thereby increasing rates of breakdown and reducing odours. BSF larvae add value to soil composition as a result of digesting organic material and the release of larval faeces into the soil, which promotes plant growth (Jeon *et al.*, 2011).

1.9.3. The Texas Experimental BSF Composting Research Programme

In an experiment conducted in Texas over a period of one year, it was found that BSF larvae are able to digest over 15 kilograms per day of restaurant food waste per square meter of feeding surface area, or roughly 3 lbs per square foot per day, leading to a 95% reduction in the weight and volume of such waste. As a result, for every 100 lbs of restaurant food waste placed into a bio-processing unit, only 5 lbs of a black, friable residue remain. Over 100,000 active larvae can be found in a typical waste disposal unit, and they can eat and digest just about any type of putrescent waste, including meat and dairy products. The instant waste is deposited into the unit, the larvae begin to secrete enzymes into the waste long before it begins to rot and smell. Since thermophilic and anaerobic bacteria play no role in this process, the larvae can conserve and recycle the majority of the nutrients and energy within the waste (Alvarez, 2012).

1.9.4. Rates of bioconversion

Over a period of one year, approximately 20% by weight of the fresh food waste is converted into fresh larvae. This food waste had an average dry matter content of 37%, and the pre-pupae has an average dry matter content of 44%, i.e., on a dry matter basis, the bioconversion of food waste situates at almost 24%. An input of 100 kg of food waste per day can be handled by three 6-foot bioconversion units.

The BSF pre-pupa is composed of:
42.1% crude protein, 34.8% ether extract (lipids), 7.0% crude fibre, 7.9% moisture, 1.4% nitrogen free extract (NFE), 14.6% ash, 5.0% calcium and 1.5% phosphorus.

Live BSF pre-pupae have been successfully fed to bull frogs, tropical fish, reptiles, snakes and many other creatures that have a strong preference for living food, and the value of fresh BSF larvae ranges from \$4 to \$20 /lb. Chickens are especially fond of the live larvae (Amatya, 2009).

1.10. Does winter pose a problem to composting?

In winter, bioprocessing using BSF can be maintained by simply placing a styrofoam sheet on top of the larval residue to retain the heat generated by larval movement. If this heat is not allowed to escape, the temperature on the surface of the residue easily exceeds 35 degrees C. During summer, the conversion rate of fresh food waste into fresh larvae runs as high as 20%, but during winter, this conversion drops to less than 5%, in spite of the fact that the larvae digest roughly the same daily quantity of food waste per unit surface area. Under the right ideal summer conditions, it takes about two weeks for newly hatched larvae to reach the mature pre-pupal form, but during the cold of autumn and winter, this two-week period may last as long as six months. Well-insulated, BSF technology units can be introduced into some of the coldest regions of Earth. During the hot summer months, overcrowding can often occur, a process which leads to relatively high temperatures within the unit; so as to cool the unit down, some actively feeding larvae are forced to exit the unit (Craig Sheppard *et al.*, 2002).

1.11. Slugs as agents of disease transmission

Slugs (Mollusca: Gastropoda) are the major pests of large varieties of vegetables and fruits in fields, home gardens, landscapes, greenhouses, (Godan, 1983; South, 1992). Slugs can also present a risk to plants, animals and humans because they serve as intermediate hosts for many bacterial diseases which predominate in animal faeces (South, 1992). Slugs usually ingest bacteria in soils as an important source of food. Consequently, they become hosts of many bacterial strains, such as *E. coli* O157, either

via direct contact or being contaminated with animal faeces spread in open fields (Walker *et al.*, 1999). Emma *et al.* (2006) showed that both the Yellow Slug (*Limax flavus*) and the Great Gray Slug (*Limax maximus*) carry *E. coli* O157 both on the surface and internally, and that slugs are contaminated with *E. coli* through contact and/or ingestion. Elliot (1969) also suggested that slugs possibly carry *E. coli* O157, and Dawkins *et al.* (1986) showed that four slug species transmit the agent of bacterial soft rot of potatoes (*Erwinia carotovora*) although this was considered to be accidentally rather than obligatory.

1.12. Transfer of bacterial species via slugs and snails to lettuce.

The enteric tract of dogs, like mammals, possesses a complex microbial ecosystem, including several bacteria such as Streptococci, *Bifidbacteria*, *Lactobacilli*, *Bacteroides* and *Clostridium* (Drasar and Hill, 1974; Drasar and Barrow, 1985). *Acinetobacter baumannii* isolated from a number of dogs and cats, is spread (Thierry *et al.*, 2008) nosocomially, a fact which explains the occurrence of several strains of *A. baumannii* in the veterinary hospital environment among dogs.

Some invertebrates may be responsible in transmission of the *E. coli* and other bacteria to field crops either by direct contact or contamination with animal faeces. The greatest concerns to human pathogens on fresh vegetables and fruits is represented by the enteric pathogens (e.g. *E. coli* O157:H7 and *Salmonellae*), which can grow before being eaten.

1.13. Detection and confirmation of Mycoplasma in dog faeces

The first documented occurrence of *Mycoplasma* in dogs was in 1934. During the last 70 years, 15 known species of *Mycoplasma* have been recognised and several have been isolated from or detected in dogs (Chalker, 2005). The isolation of a *Mycoplasma* from a human was first reported in 1937 (Taylor-Robinson, 1996; Kudva *et al.*, 1998).

Four species, *M. hominis*, *M. orale*, *M. pneumoniae*, *M. salivarium* have been isolated in humans as well as dogs (Colaizy *et al.*, 2003).

Culture techniques remain the most commonly used approaches to the detection of the presence of *Mycoplasma* in environmental samples such as canine samples. However, there are now a wide variety of indirect test methods available for *Mycoplasma* detection, including PCR- based kit, DNA fluorochrome staining, autoradiography, ELISA (McGarrity *et al.*, 1985; Lincoln and Gabridge, 1998; Rottem and Barile, 1993)

The most widely used recommended indirect test is DNA fluorochrome staining (McGarrity *et al.*, 1983), an easy and relatively fast procedure which stains DNA using a fluorescent dye. When stained and fixed cells are examined under a UV microscope equipped with the proper filter package, DNA fluoresces brightly and negative control slides should always be used to help interpret staining results. These positive and negative *Mycoplasma* control slides are commercially available. The best overall testing approach is a combination of both methods: direct culture can be provided very high sensitivity while DNA fluorochrome staining can detect any fastidious *Mycoplasma*.

1.14. Nitrification

1.14.1. The Nitrogen Cycle

Nitrogen is essential for life, it is the main component of amino acids which are the building blocks of peptides and protein, and is found in important biological components such as chitin and mucopeptides; it is also an integral part of the genetic material of cells, the nucleic acids. Plant growth in soils throughout the world is often restricted by the supply of available N and, as a result, it is nitrogen supply, more than any other soil nutrient which limits UK and world crop production. Because of this large amounts of nitrogen are applied globally as fertiliser in order to increase crop productivity (Lam *et al.*, 1996)


. In agricultural systems, the need to understand the nitrogen cycle is of extreme importance if maximum crop yields are to be achieved. In natural ecosystems, no additional, fertiliser nitrogen is applied, but the need to understand the soil N-cycle is just as critical.

1.14.2. Ammonification

The great bulk (95-99%) of the soil nitrogen is in organic compounds which are largely unavailable to higher plants (Pate, 1973). When soil microorganisms degrade these compounds, simple amino compounds (R-NH₂-) are formed. Many soil microorganisms are able to deaminate amino acids (e.g. bacteria, actinomycetes and fungi) with the resultant release of ammonia. Any NH₄-N that accumulates in soil represents the quantity of substrate nitrogen in excess of microbial requirements (Richards, 1987). Ammonium production is referred to as ammonification and the fate of the ion varies, depending upon conditions in the soil. Ammonia as a gas is volatile and leaves the soil; however, if dissolved in soil water the ion-NH₄ is formed. Ammonium can be accumulated and utilized by plants and microorganisms and under favourable conditions can be oxidized to nitrate.


The flow of nitrogen in the soil is intimately tied to flow of carbon and the processes involved in the nitrogen cycle bring about changes to the soil environment that also have an influence on other soil processes and cycles

Recently, the definition of nitrification has extended to refer to the biological oxidation of any reduced form of nitrogen to a more oxidised form (Killham, 1994).



1.14.3. Autotrophic nitrification in soil.

It is generally accepted that the major type of nitrification in most agricultural soils is chemoautotrophic, largely carried out by the Gram-negative bacteria *Nitrosomonas* and *Nitrobacter*. (Killham,1994). The reactions carried out are summarised below



In the case of *Nitrosomonas*, the oxidation state of nitrogen is changed from -3 to $+3$, and in the cases of *Nitrobacter* from -3 to $+5$. The energy yields to the chemoautotrophs are approximately 65 kcal (or 8.8 ATP molecules) per mole for *Nitrosomonas* and 18 kcal (or 2.5 ATP molecules) per mole for *Nitrobacter*, energy yields which are somewhat low compared with heterotrophic metabolism. A mole of glucose can for example optimally give an aerobic microbe 280 kcal (or 38 ATP molecules). This partly explains why autotrophic nitrifiers grow relatively slow in the soil and even in

laboratory culture where conditions for growth are optimised. Natural generation times for nitrifying bacteria of the order of 20-40 h, together with their low numbers in most soils however, gives a highly misleading impression of their vital contribution to nitrogen cycling and to soil ecology in general (Killham,1994).

1.14.4. Nitrate reduction

Once nitrate is formed in soil, it is subjected to the following fates;

- 1- It can undergo microbial denitrification to gaseous oxides of nitrogen to dinitrogen-N₂.
- 2- It can be utilized as a source of N for plants and microorganisms. Assimilation of NO₃ is followed by its reduction to NH₄⁺, which is then utilized (i.e. assimilatory reduction) (Alexander, 1977, Paul and Clark, 1989).
- 3- In the absence of O₂, nitrate can be used by microorganisms as an electron acceptor and as a result be reduced to NH₄⁺ (dissimilatory reduction) (Paul and Clark, 1989).
- 4- Being a negatively charged ion, NO₃⁻ is easily leached through soil and into ground water and soil nitrate leaching has several consequences. When nitrate is leached, it reduces the base saturation of a soil and increases exchangeable acidity. High concentrations of nitrate in surface waters can also lead to eutrophication and fatal diseases such as gastric cancers and methaemoglobinaemia (Alexander 1977).

1.14.5. Nitrogen losses from soil

Of all the nutrients required for plant growth, N is by far the most mobile and subject to greatest loss by physical, chemical, and/or biological processes from the soil-plant system (Knowles 1981, 1982). Even under the best circumstances, no more than two thirds of the fertilizer-N is accounted for by crop removal or recovered in the soil at the end of the growing season and nearly one-half of the applied amount can be lost. Five main processes for N loss occur, including microbial denitrification, chemo-


denitrification, NH₃ volatilization, leaching, and erosion. There are two biological processes for reduction of oxidised N forms (e.g., NO₃⁻ and NO₂⁻). One of these, assimilatory NO₃⁻ reduction, the other process is dissimilatory NO₃⁻ reduction, more commonly known as denitrification (Alexander 1977, Cooper, and Smith, 1963, Lynch 1983, Payne 1981).

1.14.6. Assimilatory nitrate reduction

In assimilatory nitrate reduction N is incorporated into cell biomass of plants, bacteria, cyanobacteria, and fungi all of which reduce NO₃⁻ to NH₄. in the biosynthesis of amino acids and proteins (Atlas and Bartha 1993). The process needs energy and several enzyme systems, including nitrate and nitrite reductases to form ammonia, which is subsequently converted into amino acids. As a result, this process is regulated by the availability of nitrogen, and nitrate utilization occurs when energy exceeds the concentration of ammonium or organic-nitrogen compounds (Atlas and Bartha 1998).

1.14.7. Dissimilatory nitrate reduction

Dissimilatory nitrate reduction is a process, in which the N is not utilized and in the absence of O₂. Nitrate ions acts as a terminal electron acceptor. The process is also known as nitrate respiration, or dissimilatory nitrate reduction (Atlas and Bartha 1998). Respiratory denitrification is usually the major dissimilatory process occurring in soils which reduces nitrate and under anaerobic conditions nitrate-respiring bacteria reduce nitrate to nitrite. Facultative anaerobes are often involved and many of these can also further reduce nitrite to ammonium:



The following genera are involved:

Table 1.1: Bacteria that can dissimilate nitrate to ammonium Based on Tiedje (1988)

| Genus | Typical habitat |
|------------------------------|------------------------|
| <i>Clostridium</i> | Soil, sediment |
| <i>Desulfovibrio</i> | Sediment |
| <i>Selenomonas</i> | Rumen |
| <i>Veillonella</i> | Intestinal tract |
| <i>Wolinella</i> | Rumen |
| Facultative anaerobes | |
| <i>Citrobacter</i> | Soil, wastewater |
| <i>Enterobacter</i> | Soil, wastewater |
| <i>Erwinia</i> | Soil |
| <i>Escherichia</i> | Soil, wastewater |
| <i>Klebsiella</i> | Soil, wastewater |
| <i>Photo bacterium</i> | Seawater |
| <i>Salmonella</i> | Sewage |
| <i>Serratia</i> | Sediment |
| <i>Vibrio</i> | sediment |
| Microaerophiles | |
| <i>Campylobacter</i> | Oral cavity |
| Aerobes | |
| <i>Bacillus</i> | Soil, food |
| <i>Pseudomonas</i> | Soil, water |
| <i>Neisseria</i> | Mucous membranes |




Fig 1. 2: The Nitrogen Cycle

Abbreviations; d, denitrification; dan, dissimilatory and assimilatory nitrate reduction to ammonium; I, immobilisation; m, mineralisation; n, nitrification and subsequent leaching; p, plant; r, root exudation and turnover (Killham, 1994)

1.15. The Sulphur cycle

Sulphur is an essential element which in plants, it is an important component of amino acids methionine and cysteine. Plants contain as much sulphur as phosphorus, and sulphur is as important in the formation of protein. Despite this, sulphur has traditionally been regarded as a plant nutrient of secondary importance to nitrogen, phosphorus and potassium.

1.15.1. Forms of S in soil

It is generally accepted that well over 90% of the sulphur in most non-calcareous, non-tropical, surface soil is in organic forms, about half in the form of sulphate esters and esters with C-O-S linkage (Tisdale, Nelson and Beaton 1985); about 20% of the sulphur directly bonded to carbon such as S-containing amino acids (Biederbeck 1978), and the remainder in a variety of largely inert organic compounds.

1.15.2. Biological and biochemical S-mineralisation

The mineralisation of organic sulphur in soil occurs by two main processes, biological and biochemical. Carbon-bonded sulphur is mineralised biologically during the oxidation of carbon by soil organisms to provide energy, whereas non-carbon bonded organic sulphur is mineralised through enzymatic catalysis outside the cell (Killham, 1994). An example of this latter biochemical release of sulphur, the form of sulphates-catalysed by cleavage of sulphate esters, Sulphur is released into the inorganic pool in various oxidation states from sulphide (oxidation state-2) to sulphate (oxidation state in ⁺⁶).




Fig 1. 3: The Sulphur Cycle

Abbreviations: I, immobilisation; m, mineralization; p, plant uptake; r, root exudation and turnover; so, oxidation and subsequent, leaching; sr, reduction (Killham, 1994).

1.15.3. Soil ecology and S-mineralization

The rate of S-mineralisation in soil is influenced by similar environmental factors which control N-mineralisation, including water potential, temperature, and pH, the presence of plants, drying/heating cycles and the form and quantity of organic sulphur have the most important influence on rates of soil S-mineralisation.

In most soils, the inorganic S-pool is very small and biological uptake, both as microbial immobilization and plant uptake depends upon an adequate rate of mineralisation. Some soils cannot meet the sulphur demands of all crops in this way,

and this S-deficiency is sometimes not offset by fertilizer, pesticide and atmospheric S-inputs (Hoque and Killham 1987).

The great similarities between the cycling of sulphur and nitrogen (notably with regard to mineralisation from organic matter) suggest a fundamental involvement of soil animals in S-mineralisation. Until further information becomes available, it can be assumed that the quantitative involvement of soil animals in the cycling of sulphur is approximately similar to that for the cycling of N. The atmosphere contains considerable amounts of sulphur released by the burning of fossil fuels and by microbiological sulphate reduction. Plants meet most of their sulphur needs from sulphate, but they may obtain some sulphur directly from the atmosphere. When plants and animals are incorporated into the soil their proteins are hydrolysed to form amino acids, which together with other sulphur containing compounds are further oxidized by microorganisms to form sulphate, while in anaerobic soils H_2S is formed partly from sulphate reduction and partly from the mineralization of organic sulphur. A number of intermediates are produced during S-oxidation, including thiosulphate, polythionates and sulphate. These ions do not generally persist however, and their concentration in nature is as a result, usually low.

1.15.4. Microorganisms involved in the S-Cycle

Microorganisms are responsible for:

- (1) The mineralization of organic sulphur to sulphate.
- (2) The oxidation of reduced forms of inorganic sulphur to sulphate.
- (3) The aerobic reduction of sulphate to sulphides.
 - (a) The immobilization of sulphate as organic sulphur.

1.15.5. S-Mineralization

Sulphur mineralization is the conversion of organic sulphur into inorganic forms, notably sulphate; it provides a substantial source of soil sulphate, notably in forest ecosystems Johnson (1932). Sulphur mineralization is therefore an important way in which sulphate is mobilised in soils (it also provides a source of H⁺ ions) (Tabatabai 1985). Sulphur mineralization increases in the presence of oxygen, temperature (in the mesophilic range, Tabatabai and Al-Khafaji (1980), moisture level and the addition of lime to acid soils (Williams 1967).

1.15. 6. Sulphate reduction


Bacteria of the genus *Desulfuvibrio* are the main microorganisms concerned with the reduction of sulphate. They use sulphate as an electron acceptor for growth. While few microorganisms are able to reduce sulphate to sulphide many bacteria, actinomycetes and fungi can reduce partially reduced inorganic sulphur compounds such as thiosulphate, tetrathionate and sulphite to sulphide (Alexander, 1977).

1.15.7. Sulphur oxidation

The oxidation of reduced sulphur in soil is generally regarded as a microbial process (Wainwright, 1978 Burns 1967), although some non-biological oxidation of the element has been shown to occur in autoclaved soils (Wainwright and Killham 1980, Nor and Tabatabai 1977).

During the microbial decomposition of organic sulphur compounds, sulphides and other incompletely oxidised substances are formed, such as elemental S, thiosulphate, and polythionates; these reduced substances are the available for oxidation. The oxidation of some sulphur compounds, such as sulphites (SO₃²⁻) and sulphides (S²⁻) can

occur by strict chemical reactions (Brady and Weil 1974) and such abiotic oxidation can occur to a limited extent in soils, but microbial reactions are clearly dominant:



The microorganisms involved in S-oxidation can be divided into: chemoautotrophs (lithotrophs), including species of the genus *Thiobacillus*

1- Photoautotrophs, including species of purple and green sulphur bacteria, and

2- Chemoheterotrophs (organotrophs), including a wide range of bacteria and fungi

1.15.8. Heterotrophic sulphur-bacteria

The ability of heterotrophic bacteria to oxidise sulphur was initially studied by Guittonneau (1927) and confirmed by Starkey (1934) who showed that soil heterotrophic bacteria can oxidise thiosulphate to sulphate via tetrathionate. A wide range of heterotrophic bacteria, fungi and actinomycetes (Yagi, 1971) can oxidise a variety of reduced forms of sulphur *in vitro*. Bacterial species of the genera *Arthrobacter*, *Achromobacter*, *Bacillus*, *Beggiatoa*, *Flavobacterium*, *Micrococcus*, *Mycobacterium*, *Pseudomonas* and *Sphaerotilus* can oxidize sulphur; unlike the S-oxidising autotrophs, heterotrophic bacteria do not seem to obtain energy from the process (Trudinger, 1967, Schook and Berk, 1978).

1.15.9. Chemotrophic sulphur bacteria

The chemotrophic sulphur bacteria vary in both morphology and physiology, ranging from specialist obligate chemolithotrophs through facultative chemolithotrophs which can grow mixotrophically, to specialist heterotrophs, some of which may not benefit directly from the oxidation of reduced sulphur compounds (Kuenen and Beudeket, 1982; Table, 1.2); the most widely studied in this group belong to the genus *Thiobacillus*. The

thiobacilli are rod shaped organisms which can obtain energy from oxidizing inorganic sulphur compounds, using oxygen as an electron acceptor, while CO₂ or bicarbonate supplies the carbon for chemoautotrophic growth (London and Rittenberg 1967). Additionally, they can be subdivided into those growing on neutral pH and those, which live at acidic pH. They can also grow both at acidic and alkaline pH values. Vitols and Swaby (1969) pointed out that Thiobacilli are important sulphur oxidisers only at pH values below 7, while heterotrophs are the primary sulphur oxidisers in neutral to alkaline soils (pH 6.0-7.5). The reduction in pH resulting from sulphuric acid formation by thiobacilli may also control some diseases of plants including potato scab (Brown, 1982).

Table 1. 2: Colourless sulphur oxidising bacteria. (Modified after Kuenen and Beudeker, 1982)

| Obligate chemolithotrophic S-bacteria | | Facultative chemolithotrophic S-bacteria |
|--|-----------------------|---|
| (a) Aerobic bacteria | | |
| <i>Thiobacillus thiooxidans</i> | | <i>T. novellus</i> |
| <i>T. neapolianus</i> | | <i>T. intermedius</i> |
| <i>T. ferrooxidans</i> | | <i>T. acidophilus</i> |
| <i>T. kabobis</i> | | <i>T. organoparus</i> |
| <i>T. tepidarius*</i> | | <i>Sulfolobus acidocaldarius</i> |
| <i>Thiomicrospira pelophila</i> | | <i>Sulfolobus brierleyi</i> |
| (b) Facultative anaerobic bacteria | | |
| <i>T. denitrificans</i> | | <i>Thiobacillus A2</i> |
| <i>T. thioparus</i> | | <i>Thermothrix thiopara</i> |
| <i>Thiomicrospira denitrificans</i> | | <i>Paracoccus denitrificans</i> |
| | | <i>Thiosphaera pantotropha**</i> |
| Chemolithoheterotrophs | Heterotrophs | Unclassified |
| <i>T. permetabolis</i> | <i>Beggiatoa</i> | <i>Thiovulum</i> |
| <i>Pseudomonas sp</i> | <i>Pseudomonas sp</i> | <i>Thiophysa</i> |
| | | <i>Thiothrix</i> |
| | | <i>Thiospira</i> |
| | | <i>Thioploca</i> |

1.15.10. The phototrophic sulphur bacteria

The phototrophic sulphur bacteria are found in anaerobic environments, (e.g. in H₂S rich mud, and stagnant waters), which remain exposed to light; since they are not found in most agricultural soils they need not be discussed further here.

1.16. Phosphorus cycling

Phosphorus is second only to nitrogen as an inorganic nutrient required by plants and microbes and is important for the accumulation and release of energy during cell metabolism. Phosphorus is added to the soil as a fertilizer, or is produced during the breakdown of plant residues or animal remains. Microorganisms are involved in a number of transformations of the element including:


- (1) Solubilization of inorganic insoluble phosphorus compounds.
- (2) Mineralization of organic compounds to form inorganic phosphate.
- (3) Immobilization of inorganic phosphate into cell components.
- (4) Oxidation and reduction of inorganic phosphorus compounds.

Some 15-85% of phosphorus in soil is organic, notably as minerals existing as insoluble phosphates. These insoluble inorganic compounds are largely unavailable to plants. Microorganisms including fungi and bacteria can solubilize this insoluble phosphate. These phosphate-solubilizing fungi include species of *Penicillium*, *Sclerotium*, *Fusarium* and *Aspergillus* (Alexander 1977). This process involves the production of organic acids and/or chelating agents (Wainwright 1981). The mineralization of organic phosphorus is an important soil process due to the existence of a large reservoir of non-plant available organic phosphorus which is converted to inorganic forms by enzymes collectively known as phosphatases. The microbial immobilization and assimilation of phosphorus may depress crop yields and the resultant phosphorus shortages can be overcome by adding phosphorus fertilizers to the soil. Fungi are also able to oxidize

reduced phosphorus compounds eg. phosphite to phosphate and can use these compounds as sole phosphorus source. There is the possibility of a reductive phosphate pathway occurring in soils with phosphate being reduced to phosphate and hypophosphate; fungi however, have not been implicated in this pathway (Alexander, 1977).

1.16.1. P-Mineralization and immobilization

The maintenance of soluble phosphate in the soil solution depends to some extent on the magnitude of the two opposing processes



Phosphorus mineralization is an enzymatic process. As a group, the enzymes involved, called phosphatases, catalyze a variety of reactions which release phosphate from organic phosphorus compounds.


1.16.2. P-solubilization

Phosphate solubilising microorganisms (fungi and bacteria) include species of *Pseudomonas*, *Mycobacterium*, *Micrococctts*, *Bacillus*, *Flavobacterium*, *Penicillium*, *Sclerotium*, *Fusarium*, and *Aspergillus* (Alexander, 1977). Many common microorganisms, including species of *Pseudomonas*, *Achromobacter*, *Flavobacterium*, *Streptomyces*, and especially *Aspergillus* and *Arthrobacter*, are able to solubilize insoluble inorganic phosphates in soil. The rhizosphere often has a particularly high proportion of such organisms, for example, having found that 20 - 40% of the bacteria, actinomycetes and fungi isolated from the rhizospheres of many plants are able to dissolve hydroxyapatite, compared with 10 - 14% from non-rhizosphere soil.


1.17. Aims of the Work Reported in this Thesis

The overall aim of the work reported in this Thesis is to attempt to correct this deficiency by providing research data relevant to:

- a) The composition of bacterial populations found in dog faeces
- b) The distribution of bacteria in playing field soils exposed to dog faeces
- c) The fertilizer potential of dog faeces and its potential to be used as additive to potting composts.
- d) The question of whether slugs can act as vectors of bacterial disease, transmitting pathogenic bacteria from dog faeces to lettuce.
- d) The potential use of Black Soldier Fly and Fruit Beetle larvae to increase the rate of composting of dog faeces.
- e) The ability of slugs, snails and earth worms to carry and transfer Mycoplasmas to lettuce from dog faeces.
- f) The potential use of dog faeces as a source of biofuels.



**Chapter Two: Transfer of Bacteria by Invertebrate
Slugs from Canine Faeces to Lettuce**



2.1. Introduction

Several bacterial species, by being associated with a variety of invertebrates, can infect humans, plants and animals; either via vectors (such as mosquitoes, the malaria disease carrier) or through infecting the food chain with bacteria, notably vegetables crops and fresh fruits.

Food poisoning outbreaks are frequently associated with cross contamination from insects, such as houseflies, to meat products; however, direct contamination of foods, notably of vegetables and lettuce in the field is also highly. Many field crops are prone to be contaminated with remains of faeces from farm and domestic animals which are in contact with manure contaminated soil or dust, and this is the major source of pre-harvest contamination. Indirect sources of contamination also result from the interaction between vegetables and fruits and phytophagous birds, mammals, and insects.

Pets and farm animals have been shown to be effective vectors of pathogenic bacteria, either in the visceral or on their remains and animal faeces have been shown to contain bacteria such as *Salmonella*, *E. coli* O157, and *Listeria monocytogenes* (Beuchat, 1996).

Dog faeces are likely to be a major a source of bacterial contamination of vegetable and lettuce crops especially where these are produced in allotments, small-holdings, or agriculture land located close to housing estates where dogs are common. The canine enteric tract, like most mammals, has a complex microbial ecosystem, which includes several bacteria such as species of *Streptococci*, *Bifidobacteria*, *Lactobacilli*, *Bacteroides* and *Clostridium* (Drasar and Hill, 1974; Drasar and Barrow, 1985). *Acinetobater baumannii* has also been isolated from a number of dogs and cats, and according to Thierry *et al.* (2008) the molecular typing of samples with limited polymorphisms of ribosomal DNA provided confirmation of nosocomial spread of this

pathogen as well as for the occurrence of several strains of *A. baumannii* in dogs in veterinary hospitals.

Slugs (Mollusca: Gastropoda) are major pests of large varieties of vegetables and fruits in fields, domestic gardens, allotments, landscaped areas and greenhouses, (Godan, 1983; South, 1992). Slugs present a risk to plants, animals and humans because they often serve as intermediate hosts for many bacterial diseases found in animals' faeces (South, 1992). Slugs usually ingest bacteria in soils as an important component of their food. Consequently, they become hosts to many bacteria, such as *E. coli* O157, either via direct contact or by being contaminated with animal faeces spread in open fields (Fig. 1.1), (Walker *et al.*, 1999). Emma *et al.* (2006) carried out a study on both yellow slug (*Limax flavus*) and great gray slug (*Limax maximus*), and confirmed that both species can carry *E. coli* O157 both on the surface and internally.

Emma *et al.* (2006) also showed that once a slug has become contaminated with *E. coli* through contact and/or ingestion, the bacterium can survive for many days internally and externally, thereby providing adequate time for successful transmission. Elliot (1969) also suggested that slugs possibly carry *E. coli* O157. However, the laboratory study of Dawkins *et al.* (1986) demonstrated that in four slug species i.e. *Arion hortensis*, *Deroceras reticulatum*, *Milax budapestensis* and *Limax maculatus* tested to see if they transmit bacterial soft rot of potatoes *Erwinia carotovora*, the association between the pathogen and its slug vectors was considered to be accidental, rather than obligatory. Invertebrates then may be responsible for the transmission of the *E. coli* O157 and other bacterial strains to fruits and vegetables either by direct contact, or following contamination with animal faeces. Human pathogens present on fresh fruits and vegetables, notably enteric pathogens (e.g. *E. coli* O157:H7 and *Salmonella*), have the ability to grow before being eaten and bacterial pathogens have been isolated from a large number of vegetables and fruits. Many strains of food poisoning bacteria

have also been isolated from soils including *Vibrio cholera*, *E. coli*, *Campylobacter jejuni* and *Shigella*. In addition, *Enterococci* have been found in soils, and their distribution is generally associated with animal faeces.

The aim of the work reported in this Chapter was to determine if bacteria can be transferred from canine faeces to vegetable crops (in this case lettuce) by slugs indigenous to the UK.

2.2. Materials and Methods

2.2.1. Slug culturing

One hundred slugs were used in this research. They were divided into small groups; sets of twenty slugs were placed in five sterilised plastic boxes (of size 20cm × 30cm × 10 cm). The grey garden slug *Limax maximus* was obtained from Blades Biological Ltd. In order to maintain the slugs alive, wet tissues were placed in the bottom of the boxes and changed every 3 days. Fresh lettuce was introduced to the slugs twice a week as their only food source. All slug-containing boxes were left at room temperature.

2.2.2. Isolation of bacteria from dog faeces

Samples were collected from fresh dog faeces. Portions (1g) were added to 99 ml of distilled water and were shaken at 70 g for 15 min. Aliquots of the shaken samples were spread onto the surface plates containing Chromagar medium, and left overnight in an incubator at 37°C (Samra *et al.*, 1998).




Fig. 2. 1: Bacterial colonies isolated from dog faeces using Chromagar medium

2.2.3. Isolation of bacteria from slugs

The slugs were starved for 2 days and then exposed to fresh dog faeces (which was collected locally) as their sole food source. Bacteria were then isolated from both the outside and inside of the animal.

Samples were collected from inside slugs in sterile deionised water as follows: Slugs were killed using formalin (90% v/v in a Petri-dish), and then dissected. Portions of 2 ml of (dH₂O) were directly injected into the digestive track of the slugs, and the samples were collected using sterile syringes (Fig.2.2). Samples (1 ml) were then added to 9 ml of distilled water and were shaken at 100 g for 15 min, Surface samples were obtained directly from the surface secretion of the slugs. Both collected samples were spread onto the surface of Chromagar media (e.g. Orientation media); and the inoculated plates were incubated at 37°C for overnight.



Fig. 2. 2: Section of digestive track of slug from where the samples were taken




Fig. 2. 3: Isolation of Bacteria from slugs: (a) colonies inside slugs, and (b) colonies from the outside of slugs.

2.2.4 Isolation of bacteria from lettuce

Bacteria were initially isolated from fresh lettuce leaves as controls. Slugs previously fed on dog faeces were then fed lettuce. After two days, the suspension was prepared from both control and contaminated lettuce leaves, by immersing small pieces of lettuce leaves in 50 ml of liquid Nutrient Agar. The samples were incubated overnight at 25°C under continuous shaking (250 g). In order to obtain different isolation batches, four dilution series (e.g. 10^{-1} , 10^{-2} , 10^{-3} and 10^{-4}) were used in this experiment, followed by spreading the samples (100µl) onto Chromagar plates which were then incubated for 18-24 hours.

2.2.5 Bacterial identification

2.2.5.1 Identification of bacteria using Chromagar medium

Chromagar Orientation medium was prepared as recommended by the Manufacturer's instructions (Chromagar Company, Paris, France). The medium was made up as follows: (per litre) 15 g of agar, 16 g of peptone, (1.3 g of chromogen mix) meat extract, and yeast extract, pH 7. The species of bacteria were presumptively identified on the basis of the colour of the colonies produced when growing on Chromagar. The colour is produced by the reaction of enzymes, produced by a specified genus or species, with certain chromogenic substrates. Chromagar Orientation media contains two chromogenic substrates used for the detection of B-D glucosidase and B-D-glucosidase enzyme activity, together with 1- pyrolidonyl-B-naphtylamide hydrolysis (PYR) which is described for the identification and differentiation of Enterococci from other bacteria (Merlino, 1997). See appendix two (Fig. 2. 2)

2.2.5.2. Extraction of Genomic DNA

Genomic DNA (gDNA) was extracted from the bacterial strains (JS1, JS2, JS3 and JS4) which were grown on LB nutrient solid medium for overnight at 37° C, DNA was isolated using Key Prep-Bacterial DNA Extraction kit by following procedure. 4 ml of culture grown overnight in LB medium was centrifuged at 6,000×g for two minutes at room temperature. The supernatant was decanted completely. The cell pellet was then washed with TE buffer and completely suspended in 100µl of buffer R1 by pipetting it up and down. Then 10µl of lysozyme (50mg/ml) was added to the cell suspension and mixed thoroughly. After 20 minutes incubation at 37°C, the digested cells were collected by centrifugation at 1000×g for 3 minutes and the supernatant was decanted immediately. The collected pellet was then re-suspended in 180µl of Buffer R2 and 20µl of Proteinase K, mixed thoroughly and incubated at 65°C for 20 minutes in a shaking water bath. In order to obtain a RNA-free DNA, 20µl of RNase A (DNase-Free, 20mg/ml) was added and followed by incubation at 37°C for 5 minutes. Homogeneous solution was obtained by adding 2volumes of Buffer BG to the previous mixture and mixed thoroughly by inverting tube several times then incubated at 65°C for a further 10 minutes, immediately 200µl of absolute ethanol was added and mixed carefully. The samples were transferred into a column assembled in a clean tube and centrifuged for 1 minute at 10,000×g, supernatants were discarded. The column was then washed with 750µl of the wash buffer and centrifuged twice for 1 minute at 10,000×g in order to remove the residual ethanol, supernatants were also discarded. Finally, the column was placed into a clean microcentrifuge tube and 100µl of preheated Elution Buffer, TE buffer or sterile water was added directly onto column membrane and left for 2 minutes which then centrifuged at 10,000×g for 1 minute to elute DNA. To verify DNA presence and purity, 10µl of each resultant DNA was mixed with 2µl of Blue/Orange 6x

loading dye and electrophoresed using 1% agarose gel. The checked DNA was stored at -20°C

2.2.5.3. Agarose preparation

To separate DNA, RNA or protein. Nucleic acid molecules are separated by size. This is achieved by moving negatively charged nucleic acid molecules through an agarose matrix with an electric field (electrophoresis) where shorter molecules move faster and migrate farther than longer ones due to the sieving effect of the gel. The presence, size and quantity of the DNA were checked and determined by electrophoresis in 1% agarose gel which was prepared as follow: 0.5g of molecular biology grade agarose was dissolved in 1ml of 50x TAE buffer and 40ml distilled water. The agarose was completely dissolved by heating in a microwave oven. The solution was mixed gently and allowed to cool to 55°C, and then 2.5µl ethidium bromide was added. After mixing, the solution was poured into a sealed gel rack and a comb was inserted at one side of the gel and allowed to stand in room temperature for 20 minutes. Then the comb and seal were removed and gel was placed into an electrophoresis tank and submerged in 1x TAE buffer. 10µl of the DNA sample was mixed with 2µl of Blue/Orange 6x loading dye and loaded into the wells. 6µl of Hyper Ladder was loaded as well into one of the wells as a reference. The samples were electrophoresed for 40 minutes at 80V to allow DNA to migrate toward the anode. The DNA fragments were visualised under a UV transilluminator and the images were captured using a connected digital camera

2.2.5.4. PCR amplification

The 16S rRNA gene was amplified with the universal bacterial forward primer (5` CCG AAT TCG TCG ACA ACA GAG TTT GAT CCT GGC TCA G 3`) and universal reverse primer (5` CCC GGG ATC CAA GCT TAC GGC TAC CTT GTT ACG ACT T

3') according to Weisburg *et al.*, (1991). A typical PCR mixture (50 µl in volume) contained the following components: 39µl sterile distilled water, 5µl 10× Buffer, 2.5µl 50 mM MgCl₂, 0.5µl forward Primer, 0.5µl Reverse primer, 1.0µl dNTPs, 1µl genomic DNA and 0.5µl Biorline Taq. The PCR reaction mixture, after incubation at 94°C for 3 minutes as an initial denaturation, were cycled 30 times through the following temperature profile: denaturation for 1 minute at 94°C; annealing for 1 minute at 60°C; and elongation for 5 minutes at 72°C with final incubation for 5 minutes at 75°C, after which 10 µl of each PCR amplification mixture was mixed with 2µl of Blue/Orange 6x loading dye and analysed by 1% agarose gel electrophoresis. In addition, 6µl of 1 Kb Hyper ladder I loading was used to confirm the correct sized product (Fig. 2.4). All PCR products were analyzed on agarose gels to check for the successful amplification 16S rRNA gene in the samples

2.2.5.5. Phylogenetic identification of unknown bacteria

All samples sent to Sheffield University Medical School Core Genetics Unit for sequencing. BLAST was used to compare the sequence. Finch TV software was used to correct the sequence then compared it with BLAST to look for the closest matches (Fig. 2.5)

2.2. 6. *Mycoplasma* identification

2.2.6.1. Samples description

A variety of samples were tested here for the presence of *Mycoplasma spp*, namely: a) Earthworms, b) both the internal organs and outer slime of slugs and snails) dog faeces, d) control and contaminated lettuce.

The earthworms were placed in polystyrene boxes together with a sterilized soil. A number of earthworms (10) were then placed in 40 ml of distilled water and then

centrifuged at 5000 g for 10 minutes. A 1 ml of the suspension was then taken to test for *Mycoplasma*. Slugs and snails were collected from the local area.




Fig. 2. 4: Standard hyperladder I with 14 lanes indicates higher intensity bands, 1000 and 10,000 and each lane (5 μ l) provides 720ng of DNA (BIOLINE supplier).




Fig. 2. 5: FinchTV software that manually adjusts errors of consensus sequences before BLASTn NCIMB database

2.2.6.2. Detection of *Mycoplasma*

The followed procedure to detect the presence of *Mycoplasma* or not was using EZ-PCR *Mycoplasma* Test Kit (Gene flow Ltd, Cat No.20-700-20).

2.2.6.3 Test samples preparation for PCR

A quantity of 1.0 ml of sample was transferred to 1.5 ml sterile Eppendoff tube and centrifuged at 10000 g for one 1 minute to sediment the *Mycoplasma*. The supernatant was transferred to a fresh 1.5 ml tube and centrifuged at 15000 g once again for 10 minutes to sediment the *Mycoplasma*. The supernatant was carefully discarded and removed then re-suspended the pellet (not always visible) in 50µl of buffer solution and mixed thoroughly using a micro- pipette samples were heated at 95°C for 3 minutes then stored at -20 °C.

2.2.6.4 PCR amplification

PCR amplification of *Mycoplasma* DNA was performed as described by the manufacturer. For the test sample, positive and negative control a reaction mixture of the following was prepared (no ice in PCR tube) by addition of the PCR mixture containing sdH₂ O, reaction mix and the test sample (Table 2.1). The tube was placed in a thermal cycle and the *Mycoplasma* programme was run. After PCR amplification, the PCR tube moved from the thermal cycler and 4 µl of 5x loading dye was added to each tube. For the positive control, the DNA template control (270bp) was provided by manufacturer whereas for negative control, the DNA template was replaced with double-distilled water.

Table.2. 1: PCR reaction mixtures for the amplification of *Mycoplasma*-DNA fragment.

| Reaction/Component | Volume |
|--|--------|
| Sterile deionized water (sdH ₂ O) | 35 µl |
| Reaction Mix | 10 µl |
| Test sample | 5 µl |
| Positive control | |
| Negative | |

The reaction procedure (Table 2.2.) consisted of the initial denaturation step at 94°C for 30 sec followed by 35 cycles of denaturation at 94°C for 30 sec., primer annealing at 60°C for 120 sec, and extension at 72°C for 60 sec, and was ended by 1 cycle of denaturation at 94°C for 30 sec., primer annealing at 60sec for 120 sec., and extension at 72°C for 5 min.

Table.2. 2: PCR amplification protocol of *Mycoplasma* DNA fragment

| Step | Time | Temperature | Number of cycle |
|------------------------|--------|-------------|-----------------|
| Initial denaturation | 30 sec | 94 °C | 1 |
| Denaturation | 30 sec | 94 °C | 35 |
| Annealing | 2 min | 60 °C | |
| Extensions/ Elongation | 1 min | 72 °C | |
| Final denaturation | 30 sec | 94 °C | 1 |
| Final annealing | 2 min | 60 °C | |
| Final extension | 5 min | 72 °C | |
| Hold | | 4 °C | |

2.2.6.5. Agarose gel electrophoresis

The amplified products (10µl), including positive and negative control, (2µl) 5x loading dye were separated in a 2% agarose gel in TAE for 45 minutes at a constant voltage of 80 V. Amplified products were visualized by UV transillumination with ethidium bromide staining. A 200 bp Ladder (Bioline, UK) was used as a molecular size standard.

2.2.7. Detection and identification of fungi

2.2.7.1. Samples isolation on PDA and Czapek-Dox media

To determine fungal species in dog faeces samples (0.1 g) of fresh dog faeces were inoculated onto PDA (Potato Dextrose Agar), while 0.1 g inoculated into CDA (Czapek Dox Agar) then incubated at 25 °C for 3 to 5 days. (Figs, 2.6 a, 2.6 b). Identification of fungi was achieved by extraction of genomic DNA and using 18S rRNA primers. Samples were 0.5- 1 ml of an overnight Potato Dextrose Broth culture.

2.2.7.2. Extraction of gDNA from pure cultures of fungi for PCR using 18S rRNA

Fungi /Yeast Genomic DNA Isolation Kit, Model 27300 from Sigma was used to extract genomic DNA from fungal isolates using the following procedure:

1. Lysate preparation; 5 ml of pure culture grown 3 days on PDA broth medium was centrifuged at 14,000 (g) for 1 minute to pellet the cell. The supernatant was decanted then 500µl of Lysis Solution added to the cell pellet. The suspended pellet was mixed with gentle vortexing, RNase 20µl was added to the cell suspension then the mixture transferred to a provided Bead Tube and vortexing again for 5 minutes at maximum speed. The bead Tube was incubated with lysate at 65C° for 10 minutes, the lysate mixed 2 times during incubation by inverting the tube. The lysate including cell debris was transferred to DNase-free microcentrifuge tube by pipetting, and the tube centrifuged for 2 minutes at 14,000 ×g. The supernatant transferred carefully to a new

DNase-free microcentrifuge tube without disturbing the pellet. An equal volume of 96% -100% ethanol was added to the lysate collected above (1µl of ethanol: 1µl of lysate) then vortexed to mix, 300µl of binding solution was added briefly to the suspension which was mixed by vortexing.

2. Binding of nucleic acid. Lysate (650 µl) was applied with ethanol to the column and centrifuged for 1 minute at 8,000 ×g, the spin column was reassembled with the collection tube, while the flow was discarded. This step was repeated with remaining lysate.

3. Column wash. 500µl of wash solution was next added to the column and centrifuged for 1 minute at 8,000 ×g.

4. Nucleic Acid elution. The column was transferred into a fresh 1.7 ml Elution tube which is provided with the kit and 100µl of elution buffer was added to the column and centrifuged for 2 minutes at 8,000 ×g; again the column was spin at 14,000 ×g for 1 minute.

5. Storage of DNA; the purified nucleic acid was stored -20⁰C for a few days, while it is recommended that samples be placed at -70⁰C for long term storage.

2.2.7.3. PCR amplification of fungal gene

The 18S rRNA gene was amplified with the ITS1 fungi forward primer (5` TCCGT AGGTGAACCTGCGG 3`) and ITS4 reverse primer (5` TCCTCCGCTTATTGAT ATGC3`) (see appendix). A representative PCR mixture 50 µl included the following reagents: 35 µl sterile distilled water, 5 µl of PCR buffer (10 mM Tris-HCl, pH 8.3, 50 mM KCl, 1.5 Mm), 2.5µl of MgCl₂, , 1µl forward Primer (ITS1), 1 µl Reverse primer (ITS4), 1.0 µl dNTPs, 4µl genomic DNA and 0.5 µl of Taq polymerase . The following thermal cycling conditions were performed in a thermal cycler BIO Rad 57BR0200; an initial denaturation step at 95⁰C was conducted for 2 minutes and a

thermocycle for 30 cycle, where each cycle consisted of 30 second at 95°C followed by 30 second at 58°C for annealing , and 30 second at 72°C elongation; and a final extension cycle of 7 minutes at 72°C. The amplified product (10µl) was added to 2 µl 6× loading dye visualized on a 2% (w/v) agarose gel made up in Tris-Borate- EDTA buffer (0.09 M Tris base, 0.09 M boric acid, 2 mM EDTA pH 8.0) containing 0.5 mg ethidium bromide ml⁻¹ to confirm the presence of amplified target DNA in the PCR mixture. In addition, 6µl of Hyper Ladder was loaded into one of the wells as a reference. Samples were run at a constant voltage of 80 V for 45 minutes. Twenty microlitres of the remaining PCR product was used for the electrochemical assay (Muir et al., 2011)

2.2.8. Identification of unknown fungi

All samples were sent to the Medical School Core Genetics Unit for sequencing. BLAST was used to compare the sequences and Finch TV software was used to correct the sequence with BLAST and look for the closest matches.




Fig. 2. 6: Fungi isolated from dog faeces grown on a) PDA medium b)




Fig. 2. 7: Purification of fungi species; a) green mould; b) black mould on PDA medium

2.3. Results and Discussion

2.3.1. Transfer of bacteria by slugs to lettuce

All isolated bacteria were grown on Chromagar (Orientation) media and incubated at 37°C for overnight. Cultures shown in (Fig. 2.1) showed different colonies grown on Chromagar media. Chromagar information sheets (reference) were used to identify genera and species of the grown bacterial colonies. There was a consistent colour reaction observed for some species or genus, according to colony colour. For instance, *Enterococcus sp*, from all isolations, appeared as blue (turquoise) colonies, while, other colonies from different isolates showed a variety of colours as shown in Table 2.3. Previous studies on slugs have shown no relationship between *E. coli* transfer and animals faeces (Elliot, 1969) or have failed to detect the presence of pathogens (Shrewsbury and Barson, 1947). On the other hand, field studies by Sproston *et al.* (2006) provided evidence that *E. coli* can be transferred to vegetables from sheep faeces via contaminated slugs.

2.3.2. Bacteria isolated from dog faeces

A number of colonies developing colours on Chromagar media were isolated from dog faeces based on colony-colour blue turquoise, beige, metallic blue, pink-red, cream is shown in Table (2.3) Based on the colour identification coding catalogues for Chromagar (Orientation) media, the genera and species identification of the isolates are: *Enterococcus sp*, *Salmonella sp*, *Staphylococcus lentus*, *E. coli*, *Proteus sp* and *Acinetobacter sp*.

2.3.3. Bacteria isolated from slugs

2.3.3.1. Bacteria isolated from control slugs

Blue bacterial colonies grew from cultures of both the internal and external parts of slugs; these were identified as an *Enterococcus sp.* Bacterial colonies of a metallic blue colour were found to grow from the outside secretion of slugs and these were identified as *Enterobacter amnigenes*, a finding which was confirmed using 16S rRNA sequence gene analysis (i.e.. 99% confirmed).

Table.2. 3: Analysis of coloured bacterial strains which require further 16S rRNA tests for their identification confirmation to the species level.

| | Colonies colour on Chromagar medium | Presence of bacteria species in different Samples | | | | | | |
|-------------------------------|-------------------------------------|---|---------|----|---------------|----|---------------------|--------------|
| | | Dogs faeces | Slugs | | | | Vegetable (Lettuce) | |
| | | | Control | | Fed on faeces | | Control | Contaminated |
| | | | In | Ex | In | Ex | | |
| <i>Enterococcus spp</i> | Blue Turquoise | + | + | + | + | + | + | + |
| <i>Enterobacter amnigenus</i> | Metallic blue | - | - | + | - | - | - | - |
| <i>Staphylococcus lentus</i> | Metallic blue | + | - | - | + | - | - | + |
| <i>Acinetobacter sp</i> | Beige | + | - | - | + | + | - | - |
| <i>Comamonas sp</i> | Blue | - | - | - | - | + | - | - |
| <i>Delftia sp</i> | Beige | + | - | - | + | | - | + |
| <i>Proteus sp</i> | Beige with brown Halo | + | - | - | + | - | - | - |
| <i>E. coli</i> | Pink-red | + | - | - | + | - | - | + |
| <i>Salmonella sp</i> | Creamy | + | | | | + | - | - |

In = Internal samples Ex= External samples

Grey shading indicates bacteria that were transferred to lettuce

2.3.3.2. Bacteria isolated from slugs fed on faeces

The results showed that five species of bacteria were grown from the inside/ outside of slugs fed on dogs faeces. All five species were isolated from the internal secretions and two out of five species grown from the surface of slugs. *Staphylococcus lentus* exhibited a metallic blue colony colour, while *Acinetobacter sp* was beige and identify confirmed (100%) using 16SrRNA analysis. *Salmonella sp* and *Proteus sp* appeared as cream and beige with a brown halo, respectively, and *E. coli* as a light red colour. Externally isolated species were identified as *Acinetobacter sp* (99%) using 16SrRNA (Table 2.3) and *Comamonas sp* showing a blue colony was also confirmed by 16S rRNA analysis is (91%).

2.3.3.4. Fungi isolated from dog faeces

Several species of fungi and yeasts were isolated from dogs faeces using PDA and Czapek Dox media (Fig 2.5), some of colonies were purified on a number of occasions (Fig 2.6). Results of g DNA and PCR-18SrRNA analysis are respectively shown in gel Fig.2.12 and Fig. 2.13. One fungal isolate was identified using 18SrRNA analysis, namely *Trichoderma asperellum* (99% similarity).

2.3.4. Transmission of bacteria from dog faeces to lettuce

Based on the colour of the developed colonies on Chromagar media, three groups of bacteria dominated the contaminated lettuce leaves, i.e. blue turquoise, metallic blue and beige, whereas only one bacterial species (i.e. *Enterococcus*) was grown from the control lettuce leaves. *Enterococcus* Spp. bacteria were regularly isolated using Chromagar from the control lettuce leaves. Contaminated lettuce samples gave two colonies which were coloured differently from *Enterococcus*., The colonies were initially identified using colony-colour references as *E. coli* and *Salmonella sp* (Table

2.3); these were not however, confirmed by 16SrRNA analysis. The results of genomic DNA and PCR amplification (Figs 2.8 and 2.9) showed that *Enterobacter amnigenus*, *Acinetobacter sp*, *Comamonas sp* and *Acinetobacter sp* were isolated from contaminated lettuce (samples, JS1, JS2, JS3 and JS4 respectively (Table 2.4) in addition to *Delftia sp*, *Staphylococcus lentus* and *E. coli* (samples MS1, MS2 and MS3) were also genomic DNA and PCR amplification achieved (Figs 2.10 and 2.11).

Table.2. 4: Bacterial identification using 16SrRNA

| S/No | S/ source | Species | S % |
|------|---------------------------------------|-------------------------------|------|
| JS1 | From outside control slugs (C) | <i>Enterobacter amnigenus</i> | 99% |
| JS2 | From inside slugs fed on faeces (C) | <i>Acinetobacter sp</i> | 99% |
| JS3 | From outside slugs fed on faeces (B) | <i>Comamonas sp</i> | 91% |
| JS4 | From outside slugs fed on faeces (C.) | <i>Acinetobacter sp</i> | 100% |
| MS1 | From contaminated lettuce (B) | <i>Delftia sp</i> | 99% |
| MS2 | From inside slugs fed on faeces (MB) | <i>Staphylococcus lentus</i> | 100% |
| MS3 | From inside slugs fed on faeces (P) | <i>E. coli</i> | 100% |

C = Cream or Beige colony, B= Blue colony, MB= Metallic blue colony, P= Pink-red colony




Fig.2.8: Extraction of genomic DNA of isolated bacterial species; (lane JS1) *Enterobacter amnigenus*; (lane JS 2) *Acinetobacter sp*; (lane JS3) *Comamonas sp*; (lane JS 4) *Acinetobacter sp*(lane L); hyper ladder and(lane JS5) *Delftia sp*




Fig. 2. 9: PCR- 16Sr RNA, amplification products of four isolated bacterial species analyzed by electrophoresis in agarose gel lanes represent; (lane L); hyper ladder;(lane JS1) outside control slugs (Metallic blue); (lane JS 2) inside slugs fed on faeces (B); (lane JS3) outside slugs fed on faeces(B); (lane JS 4) outside slugs fed on faeces (C); (lane 5) contaminated lettuce(Pink-red).




Fig. 2. 10: Extraction of genomic DNA of isolated bacterial species; (lane L); hyper ladder and (lane MR1) *Delftia sp* ; (lane MR 2) *Staphylococcus lentus*; (lane MR3) *E. coli*




Fig. 2. 11: PCR-16Sr RNA, amplification products of isolated bacterial species analyzed by electrophoresis in agarose gel lanes represent; (lane L); hyper ladder;(lane MR1), contaminated lettuce (Blue); (lane MR 2) Inside slugs fed on faeces (Metallic blue); (lane MR3), Inside slugs fed on faeces (Pink-.red)

2.3.5. *Mycoplasma* detection

2.3.5.1. Attempts to isolate *Mycoplasma* from various samples

Samples were examined for the presence of *Mycoplasma* contamination using the EZ- Polymerase Chain Reaction (PCR) assay. The samples were obtained from the inside and outside of slugs and snails, whole body of earthworm, dog faeces samples and lettuce control and following exposure to slugs fed on dog faeces (Table 2.5).

Table.2. 5: Presence of *Mycoplasma* in various samples

| Samples | Symbol | Reaction |
|----------------------------|--------|----------|
| Outside slug | SR1 | Positive |
| Inside slugs | SR2 | Negative |
| Earthworm | SR3 | Positive |
| Control lettuce | AZ1 | Negative |
| Contaminated lettuce | AZ2 | Positive |
| Control snail | AZ3 | Negative |
| Outside contaminated slug | AZ4 | Negative |
| Inside contaminated slug | AZ5 | Positive |
| Inside contaminated snail | AZ6 | Positive |
| Outside contaminated snail | AZ7 | Negative |
| Dog faeces | AZ8 | Positive |

The PCR technique (Figs 2. 14, 2.15 and 2.16) was successful at detecting *Mycoplasma* species from dog faeces and from slugs and earthworm collected from local areas of Sheffield. The main finding of the work reported here is that *Mycoplasma* was transmitted from dog faeces to slugs or snails which they, ingested or carried, and then in turn transferred to lettuce. Earthworms were also shown to carry *Mycoplasma* from soil on or in their bodies, a fact which presumably reflects contamination from animal faeces, mainly dogs.




Fig. 2. 12: Extraction of genomic DNA fungi species ; (lane L); hyper ladder and (lane 6 and 7) fungi isolated from dog faeces




Fig. 2. 13: PCR-18SrRNA, amplification products of fungal species analyzed by electrophoresis in agarose gel lanes represent; (lane L); hyper ladder; (lane5) *Trichoderma asperellum* (lane 6) no result.




Fig. 2. 14: Polymerase chain reaction (PCR) detection of *Mycoplasma* species in Earth worm samples, EZ-PCR-*Mycoplasma* test analyzed by electrophoresis in 2% agarose gel; the lanes represent (lane M); 1kb hyper ladder, (lane 1); negative control(distilled water). ; (lane 2); earth worm sample; (lane 3); negative result; (lane 4) positive control




Fig. 2. 15: Polymerase chain reaction (PCR) detection of *Mycoplasma* species in Dog faeces and lettuce samples, EZ-PCR- *Mycoplasma* test analyzed by electrophoresis in 2% agarose gel; the lanes represent (lane L); 1kb hyper ladder, (lane 1); negative control (distilled water).; (lane 2); negative result; (lane 3); dog faeces samples ; (lane 4) control lettuce; (lane 5); contaminated lettuce




Fig. 2. 16: Polymerase chain reaction (PCR) detection of *Mycoplasma* species in Dog faeces and lettuce samples, EZ-PCR- *Mycoplasma* test analyzed by electrophoresis in 2% agarose gel; the lanes represent (lane L); 1kb hyper ladder, (lane 1); Outside control slug (negative result) (lane 2); Outside contaminated slug; (lane 3); Outside control snails (lane 4) Outside contaminated snails (lane 5); Inside control slugs; (lane 6); Inside contaminated slugs; (lane 7); inside control snails (lane 8); Inside contaminated snails (lane 9) negative control (distilled water)

2.3.5.2. Comments on the detection of *Mycoplasma* using PCR procedure

In this study the EZ-PCR technique was used to detect culturable-independent *Mycoplasma* DNA. This kit can detect a range of *Mycoplasma* species such as (*M. fermentans*, *M. hyorhinism*, *M. arginini*, *M. orale*, *M. salivarium*, *M. hominis*, *M. pulmonis*, *M. arthritidis*, *M. bovis*, *M. pneumoniae*, *M. pirum* and *M. carpricum*) in addition to *Acholeplasma* and *Spiroplasma*, all with a high degree of sensitivity and specificity. The main aspect of this method is that rRNA gene sequences of prokaryotic, including *Mycoplasma*, are well conserved, whereas, the sequences and length of the spacer region in the rRNA operon (e.g. the region between 16S and 23S gene) differ from species to other. Two primers were used to amplify the conserved 16S ribosomal DNA (rDNA) coding region. The primer sequences allow for the detection of *Mycoplasma* DNA but not any other bacterial DNA that may contaminate sample preparations or solutions used for the PCR. Therefore, amplification of the gene sequence with PCR using this primer set enhances not only the sensitivity, but also the specificity of detection. Amplified products are then detected by agarose gel electrophoresis. Although our study was focused on domestic dogs, several species of *Mycoplasma* found in dogs have also been found in humans and other animals which means the potential of transition several species of pathogenic *Mycoplasma* from canine and amongst animal species (e.g. *M. arginini*) has been isolated from a range of hosts which are associated with canines, including goats, sheep, cattle, healthy cats (Tan *et al.*, 1974 and 1977), and camels suffering from pneumonia.

**Chapter Three: Reduction in Bacterial Numbers in
Dog Faeces Using Black Soldier Fly and Fruit Beetle
Larvae**

3.1 Introduction

Animal faeces is known to contain pathogens such as *Escherichia coli* and *Salmonella spp*, recent research has shown that waste production from animals is increasing annually as a result of population growth and human activities (Mawdsley *et al.*, 1995; Pell, 1997). Around a million tons of hen waste is produced annually worldwide (Turnell *et al.*, 2007), while approaching up to one billion tons of cow wastes is produced annually in the United State alone (Islam *et al.*, 2005). This type of waste leads to the pollution of water sources, soils, crops and the air by animal wastes. Factors such as temperature, moisture, aeration and pH value, in addition to manure composition and animal health together influence the population of pathogens present prior to, and during manure storage (Erickson *et al.*, 2004). *Escherichia coli* normally grow over the range of pH 5 to 9 but under laboratory condition can survive several hours at pH 2 to 3 (Small *et al.*, 1994). In order to manage animal wastes a variety of experimental treatments have been attempted, e.g. the mixing of animals manure with carbon sources in order to accelerate the biological oxidation of wastes (Kashmanian and Rynk, 1996). Such a treatment makes the resultant manure more effective for use in agricultural and also kills pathogenic bacteria by the production of high temperature (40-55⁰C) during the primary stages of composting (Jiang *et al.*, 2003). However, failure to maintain this heat for long periods may allow bacteria to survive (Pryor Williams *et al.*, 2006; Nemiroff and Patterson, 2007). Consequently, the possibility exists that insufficiently treated manure will lead to bacterial contamination of soil and agricultural crops (Cekmecelioglu *et al.*, 2005). As well as biological treatments, chemical approaches can successfully reduce pathogens in manure, examples include: gassing with ammonia and the application of sodium carbonate (Himathongkham, 1999; Park and Diez-Gonzalez, 2003).

Invertebrate larvae have also been evaluated in the treatment of animal wastes and manures; saprophagous or polyphagous larvae can take up as important elements during benefaction of animal waste and increase their protein and carbohydrate contents as a result; in this way, not only is the manure improved for fertilizer use, but the nutrient content of the larvae is increased, making them ideal as food sources for, for example poultry chicken and egg production (Erickson *et al.*, 2004). Black Soldier Fly (BSF) larvae, *Hermetia illucens* L. (Diptera:Stratiomyidae) have been widely evaluated for this purpose. The life cycle of BSF has four stages; egg, larva, pupa and adult. The larvae range in size from (3– 9 mm) and usually have five larval instars. Black Soldier Fly larvae have been used to bioremediate livestock manure in studies by Sheppard *et al.*, (1994) and Sheppard and Newton, (2001) and a number of studies have shown that BSF larvae can reduce the pathogenic bacteria in waste, e.g. *E coli* and *Salmonella Spp* in chicken manure (Erickson *et al.*, 2004; Jeon *et al.*, 2011). Black Soldier Fly larvae can also dry and aerate manure in addition to reducing odours by eating and ingesting of bacterial and fungicidal compound, as a results BSF larvae can modify the microflora of manure and reducing the number of harmful species (Jeon *et al.*, 2011).

The studies reported here were aimed at using BSF larvae and Fruit Beetle grubs (Sun Beetle larvae, FB) *Pachnoda marginata peregrina* (Coleoptera: Scarabaeidae) to reduce the bacterial population of dog waste and also to improve texture and fertilizer nutrient content in the hope that treated dog faeces might be converted from an offensive waste into a useful compost material.

3.2. Materials and Methods

3.2.1 Methods for bacterial isolation and identification

Several broth and agar-based media and selective media based on colony coloration were used:

HiCrome *E. coli* (TM) Agar is a selective medium obtained from Fluka-Sigma-Aldrich Company (see media in Appendix). The medium was used to isolate and enumerate *E. coli* as recommended by Hansen and Yourassowsky, (1984). Most of the *Escherichia coli* strains can be differentiated from other coliforms by presence of the enzyme glucuronidase which is highly specific for *E. coli* (see medium composition in Appendix). The chromogenic agent X-glucuronide used in this medium detects glucuronide activity. *Escherichia coli* cells absorb X-glucuronide and the intracellular glucuronidase splits the bond between the chromophore and the glucuronide. The released chromophore then gives a blue colouration to the colonies (Fig. 3.6 a) of *E. coli*. Brilliance *E. coli*/Coliform agar from Oxoid, on the other hand, gave a purple colour with *E. coli* (Fig.; 3. 6 b).

A highly selective XLT-4 Agar CM 1061 (Xylose Lactose Tergitol™ 4) agar medium obtained from Oxoid was used to isolate and identify *Salmonella* sp., as recommended by Mallinson *et al.*, (2000). Differentiation on this medium is facilitated by the fermentation of xylose, lactose and sucrose as well as the decarboxylation of lysine. *Salmonella* appear as black or red colonies with a black centre (Fig. 3. 7), due to their ability to reduce thiosulphate to hydrogen sulphide, which then causes the colony to blacken (as there are ferric ions present in the medium). The makers instruction for the preparation of this medium are as follows: Suspend 59g of XLT-4 Agar Base in 1 litre of distilled water, then 4.6ml of XLT-4 SR0237 selective supplement was added to the medium and brought to the boil. The medium was not autoclaved. After cooling

to approximately 50°C, the medium was poured into sterile Petri-dishes (Tate *et al.*, 1990).

A range of media including Luria-Bertani (L.B) , Plate Count Agar (Fig. 3.8) and Mueller Hinton (M.H) and Nutrient Broth or Agar were also prepared for routine isolation and growth studies. Bacteria were also identified using 16s rRNA analysis; DNA was extracted according to the instructions of the Anachem Key prep-Bacterial DNA Extraction kit, and then bacterial 16S rRNA gene was amplified and sequenced using PCR. After checking, the purified PCR products were sent to the Medical School, (University of Sheffield) to be sequenced. 16SrRNA gene sequences were adapted using the Finch TV software and then exported into The Basic Local Alignment Search Tool (BLAST), available from the website of National Centre for Biotechnology Information (NCBI)<http://www.ncbi.nlm.nih.gov>, to identify matches with existing characterized reference sequences

3.2.2. Dog faeces and insects larvae collection

Dog faeces (from an Alsatian) were collected with the cooperation of a dog owner in Sheffield. The fresh dog faeces samples were distributing onto two small boxes (30×20×15 cm) and placed at laboratory condition whereas the first box treated with Black Soldier Fly larvae (Fig.3.1). One hundred larvae per 100g dog faeces approximately were used, as suggested by Li *et al.*, (2011). The second box included 100g of dog faeces (without larvae) as the control. Fruit Beetle larvae were treated in the same manner, except that 10 FB larvae (Fig.3.3) were added to 100 g of dog faeces; controls were also set up. The larvae were purchased online from Ricks LiveFood. Black Soldier flies were reared to the adult stage (Fig.3.2) and Fruit Beetle stages from larvae (Fig. 3.3) pupa (Fig. 3.4) and adults (Fig. 3.5) to the next generation.

3.2.3. Bacterial isolation from dog faeces

Bacteria were isolated at weekly intervals over a 28 day incubation period. Samples 1g were diluted in 9 ml of sterilized water then a serial dilution from 0.1 to 10^6 was performed; the 0.1 ml of each diluted samples were then spread on *E. coli* agar, XLT4, L.B and count plate media, and incubated between 18 and 24 h at 37°C. In this case, both the control and treatment dog faeces were sampled from control and treatment (with larvae) boxes in triplicate at each sampling day (0, 7, 14, 21 or 28)

3.2.4. Bacterial isolation from the larval gut

Bacteria were isolated from BSF and FB larvae gut and surface after being fed on faeces. For internal isolation, the outer surface was rinsed three times in sterilized water. The larvae were then anaesthetized using a piece of cotton wool wetted with chloroform (100%), placed with the larvae in Petri-dishes for few minutes. The anterior end, near the cephalopharyngeal skeleton, was opened and the abdominal contents were squeezed gently to collect 1g of the gut contents. A serial dilution was then prepared for BSF and FB larvae to determine the number and type of bacteria, and 0.1 ml of each diluted samples was taken for both control and treatment larval gut and spread onto the *E. coli* agar, XLT4, and count plate media and incubated between 18 and 24 h at 37°C; triplicates were used throughout.

3.2.5. Determination of pH

The pH of the dog faeces was determined as follows; 5 g of dog faeces were dissolved in 50 ml deionised water (Erickson *et al.*, 2004); suspension were shaken for 15 min on a reciprocal shaker at 100 g until the suspension became homogenous, The pH of the final solution was measured weekly using a calibrated Jenway 3310 pH meter (Bogdanov *et al.*, 2002).

3. 2. 6. Statistical analyses

All observations are presented as Mean \pm SD (Standard deviation). Bacterial populations were converted to log CFU/g before statistical analysis. Statistical analyses were performed on experimental datasets using the t-Test and three way ANOVA at >0.05 probability level compared significance of means of three replication. Results were analysed using Sigma Plot 11.0[®] software.



Fig. 3. 1: Black Soldier Fly larvae, 1st instar



Fig. 3. 2: A Black Soldier Fly adult



Fig. 3. 3: Fruit Beetle larvae 3rd instar




Fig. 3. 4.:Fruit Beetle pupae



Fig. 3. 5: Fruit Beetle adults

3.3. Results and Discussion

3.3.1. Identification of bacteria associated with dog faeces and larvae

Several species of bacteria were isolated from dog faeces and intestinal tract (gut) of both the Black Soldier Fly and the Fruit Beetle FB larvae using the chromogenic media, Hicrome *E. coli* and Brilliance *E. coli*/ Coliform agars. XLT-4 agar was used to isolate and identify *Salmonella Spp.* The results (Table 3.1) show that *E. coli* produced a blue or pink/purple colony (Fig. 3.6), while *Salmonella enterica* produced a black colony on these media (Fig. 3.7). Bacteria growing on Plate Count Agar were not identified (Fig. 3.8). Scanning Electron Microscope (SEM) confirmed that the morphology of bacteria isolated from dog faeces and gut of larvae agreed with *E. coli* and *Salmonella enterica* (Fig. 3.9 a) while some of bacteria species were cocci, e.g. MRSA(Fig. 3.9 b). Molecular identification of gDNA using 16S rRNA results confirmed the species isolated as *Escherichia coli* and *Salmonella enterica*. The samples were sent to Medical School Core Genetics Unit for sequencing, and then we used (BLAST) to compare the sequence. Finch TV software was used and compared with BLAST to look for the closest matches. The results show the species isolated from dog faeces were: *E. coli* and *Salmonella enteric* 100% and 99% respectively see appendix 2 bacteria sequences

Table. 3. 1: Media used to detect and identify *E. coli* and *S. enterica* from dog faeces, BSF and FB larvae gut

| Sample | Bacterial species | Appearance of the colony | SEM / Cell morphology | Media name | Gram nature | DNA Similar |
|------------|--------------------|--------------------------|-----------------------|-----------------------|---------------|-------------|
| Dog faeces | <i>E. coli</i> | Blue or purple | <i>Bacillus</i> | HiCrome or Brilliance | Gram negative | 100% |
| BSFL gut | <i>E. coli</i> | Blue or purple | <i>Bacillus</i> | HiCrome or Brilliance | Gram negative | 100% |
| FBL gut | <i>E. coli</i> | Blue or purple | <i>Bacillus</i> | HiCrome or Brilliance | Gram negative | 100% |
| Dog faeces | <i>S. enterica</i> | Black | <i>Bacillus</i> | XLT-4 | Gram negative | 99% |
| BBSFL gut | <i>S. enterica</i> | Black | <i>Bacillus</i> | XLT-4 | Gram negative | 99% |
| FBL gut | <i>S. enterica</i> | Black | <i>Bacillus</i> | XLT-4 | Gram negative | 99% |




Fig. 3. 6: *Escherichia coli* grown on selective chromogenic agar *E. coli* media isolated from dog faeces and larvae fed on dog faeces; a) Brilliance *E. coli* /Coliform agar showing blue colonies, b) HiCrome *E. coli* agar showing purple/pink coloured colonies



Fig. 3. 7: *Salmonella Spp* grown on XLT-4 selective *Salmonella* agar medium




Fig. 3. 8: Colonies of bacteria grown on plate count agar medium




Fig. 3. 9: Two bacterial isolates showing different external morphology; a) cocci, b) bacillus, observed by SEM.




Fig. 3. 10: Bacterial numbers in dog faeces grown on; a) plate count medium, b) HiCore *E. coli* in and c) *Salmonella* medium after treatment with Black Soldier Fly larvae (Means of triplicate (±) SD log CFU/g (*significantly different from control))




Fig. 3. 11: Bacterial numbers from dog faeces, bacteria grown on; a) plate count medium, b) HiCore *E. coli* medium and c) *Salmonella* medium by Fruit Beetle Larvae.

(*Significantly different from control)

Figure 3.10a shows that the “total heterotrophic bacterial count” in the faeces treated with BSFL were markedly and statistically reduced. The same trend was found in relation to numbers of *E.coli* (Fig.3.10b) and *Salmonella* (Fig.3.10c). Figure 3.11a shows that the total number of heterotrophic bacteria in faeces treated with FBL, as was the case with BSFL declined compared to the control, at least for the first 14 days; the numbers in the control then declined rapidly so that numbers in the treated faeces then exceeded the control at days 21 and 28. This marked decline in bacterial numbers following BSF treatment was also seen in relation to treatment with *E.coli* and *Salmonella* (Figs.3.11b,c).

The overall trend following BSF larval treatment was a reduction in numbers for the first 14 days, followed by an increase over the control for days 21 and 28. The increase in numbers after 14 days may however, have been anomalous due to a rapid and unexplained fall in numbers in the control. The results show that BSFL can be used to reduce the numbers of bacteria, including some important pathogens in dog faeces. The findings relating to the use of FBL are in agreement with this trend up to day 14 but increases in bacterial population then occur which, if the control’s rapid decline is not anomalous is worrying in relation to the potential use of FBL as a compost additive or fertilizer.




Fig. 3. 12: Numbers of bacteria; a) *E. coli*, b) *Salmonella enterica* and bacterial species grown on plate count medium isolated from gut of BSF larvae

Means of triplicate (\pm) SD log CFU/g (*Significantly different from control).




Fig. 3. 13: Numbers of total “heterotrophic bacteria” in gut of FB larvae; a) *E. coli* b) *Salmonella enterica* and c) bacterial species grown on plate count medium
Means of triplicate (\pm) SD log CFU/g (*Significantly different from control).

Figure 3.12 shows the number of bacteria, including pathogens inside the gut of BSFL following feeding on dog faeces. The overall trend is obvious, namely feeding leads to a statistically significant increase in the number of bacteria inside the BSFL gut. The same trend is seen in relation to dog faeces fed FBL (Fig 3.13). This trend of increasing bacterial numbers in larvae fed on dog faeces is particularly worrying in relation to the potential feeding of these larvae to animals-post exposure to faeces. Black soldier Fly larvae in particular are currently being produced for use as animal feed (mainly to chickens) following feeding on restaurant food waste. It is unlikely then, considering the high pathogen content of larvae fed on dog faeces that these could be safely used as animal feed following feeding on dog faeces. The only reference seemingly available on the effect of feeding BSFL on animal faeces was reported by Erickson *et al.* (2004), who found that they reduced the pathogenic bacteria population, including *E. coli* and *Salmonella* in chicken and hogs manure.

Chapter Four: Modification of Canine Faeces by Black Soldier Fly and Fruit Beetle Larvae in relation to Plant Nutrient Release and Potential Use of Modified Faeces as a Compost Additive

4.1 Introduction

The digestive tract of animals contains a wide range of microorganisms and as a result, their faeces may also contain these organisms which include a variety of pathogenic and saprophytic faecal bacteria (Mawdsley *et al.*, 1995; Pell, 1997).

The accumulation of dog faeces in the urban environment poses a major threat to animals, plants, human health and water sources (Rippy *et al.* 1997). In the United States alone, dog faeces contamination is estimated to be around ten million tons annually (Brinton and Storms 2004) and the disposal of dog faeces is a major problem in regions of high-density dog populations. Dog populations vary worldwide; in the city of Quebec for example, approximately 25% of residents own at least one dog (Marketing, 2002). The estimated dog population of Quebec is 742,728, with 432,000 being found in Montreal alone (Institute de la statistique Quebec 2005a). A European survey showed that 44,000,000 domestic animals, including 6,900,000 dogs, were living in Italy in 2002. In 2007 a random study in the UK revealed that dogs were owned by 26% of households, while the 2006 dog population was around 10.5 million. The average daily faecal production of a dog is approximately 100 grams and open spaces, public gardens, pathways, arcades, pedestrian precincts and roadways are the sites with the highest pollution (Tarsitano *et al.*, 2010). Canine faeces, are often not removed from the urban environment, due to the bad habits of the owners and they clearly represent a source of potential pathogens in addition to being an unsightly inconvenience (Nemiroff and Patterson, 2007).

In the past, series of treatments have been used to reduce the pathogenic bacteria content of animals manures; the mixing of animals manure with various carbon sources being the most common method used to activate the biological oxidation process. (Kashmanian and Rynk, 1996). In this treatment method, pathogenic bacteria are killed by the high temperature (40-550C) produced in the primary stages of the composting

process (Jiang *et al.*, 2003). However, the failure to maintain such high temperatures for long periods may allow pathogens to survive (Pryor-Williams *et al.*, 2006, Nemiroff and Patterson, 2007) Consequently, the possibility for bacterial contamination of soil and agricultural crops resulting from the of inefficiently composted manure is a concern (Cekmecelioglu *et al.*, 2005). In addition, chemical treatments that have been used with success to reduce pathogens in manure (including gassing with ammonia and the application of sodium carbonate (Himathongkham, 1999) may introduce further environmental risks.

Many insect species depend completely on animals manure as organic materials as their main food source, so that a number of insects (Saprophagous) can be used to help recycle and improve the fertilizer quality of animal manures (Erickson *et al.*, 2004).

Recently, a range of studies on animal waste benefaction have been conducted in China, United States, Mexico, Eastern Europe, Australia and Central and South America, aimed at treating animals faeces with insects to produce manure and animal feed in the form of larvae; insect larvae were found to reduce the nutrient concentration and mass of the manure residue, thereby reducing the pollution potential of the waste by 50-60% or more. (Newton *et al.*, 2005)

A novel approach involves the use of black Soldier Fly (BSF) larvae *Hermetia illucens* L. (Diptera: Stratiomyidae). The life cycle of BSF has four stages; egg, larva, pupa and adult. The larvae range in size from (3– 9 mm) and usually have five larval instars. Female adults do not need to feed and lay around 900 eggs in their short life of 5 to 8 days; adults surviving on the large fat body stored from the larval stage. The black Soldier Fly is not recognized as a pest since the adult Fly is not attracted to human habitation or foods, nor does it bite or sting (Newton *et al.*, 2005)

The larva of the black Soldier Fly is a common sight in some countries in decaying organic matter including kitchen waste, spoiled feed and the larval stage of BSF and its

use has been suggested as a means of reducing livestock manures (Craig Sheppard *et al.*, 1994; Sheppard and Newton, 2001) a number of studies have demonstrated the ability of BSF larvae to recycle animals manure and reduce the number of pathogenic bacteria which they contain; *E coli* and *Salmonella spp* populations have, for example, been reduced in chicken manure (Erickson *et al.*, 2004). Previously, several studies have also reported on the possibility of recycling the dog faeces using physical and chemical approaches whereas some have utilized BSF larvae to reduce the bacterial population from animals manure such as hogs, cows and poultry. The work discussed in this Thesis focuses on using the BSF larvae to treat dog faeces. In addition, Fruit Beetle larvae FBL *Cotinis mutabilis* (Scarabaeidae: Coleoptera) were investigated and compared with BSFL in relation to their ability to reduce populations of *E. coli* and *Salmonella Spp.*

The aim of this study therefore was also to investigate changes in the concentration of ammonium and nitrate, and sulphur oxidation and phosphate solubilisation during larval treatment of dog faeces with of BSF and FB larvae in the hope of converting the biomass of these wastes dog faeces from a toxic, unsightly waste-product, into a safe and useable fertilizer and compost additive.

4.2. Materials and Methods

4.2.1. Sample collection

4.2.1.1. Samples collection from canine waste

Samples of fresh dog faeces were collected from a single dog (Alsatian) owned by a local family. The collected samples were distributed into eight plastic boxes (15x 20x 10 cm), each box containing 100 g of dog faeces with or without larvae. Three replicates were used and the results compared using T-Test at 0.05 probability level and ANOVA to determine levels of significance using Sigma Plot 11.0.

4.2.1.2. Samples collected from treated dog faeces

After 28 days, the treated dog faeces (now compost) were collected, weighed and then divided into control and treatment without added larvae. Samples (100g) of the treated and control faeces for each element were then placed in polythene bags. Treated samples were amended by addition ammonium, nitrate, sulphur and insoluble phosphate reagents while the control no elements reagents were added to the samples. Samples were taken weekly to measure the elements.

4.2.1.3. Effect of larvae on the nutrient content of dog faeces.

A number of first instar Black Soldier Fly (BSF) and Fruit Beetle (FB) larvae were bought from a commercial company (Ricks Live-Food) in UK. Dog faeces (100g) were placed in the plastic boxes and 100 larvae of BSF and 10 larvae of FBL were added; a control with no larvae was also set up. The treatments were left for 28 days under laboratory conditions and sampled at 7 day intervals for the following plant-fertilizer ions: ammonium, nitrate, sulphate and phosphate.

4.2.3. Nutrient transformations in modified dog faeces amended with ammonium nitrate, sulphur and insoluble phosphate

Samples of dog faeces which had been modified by treatment with either BSFL or FBL and the larvae were removed. The two types of modified samples (100g) were then amended, in triplicate (in polythene bags) with ammonium (5ml of a 100ug per ml solution of NH_4SO_4) and ammonium and nitrate was measured; elemental sulphur (1g) and sulphate was measured; insoluble calcium phosphate ($\text{Ca}_3\text{O}_8 \text{P}_2$, 1g) and soluble phosphate was measured. Controls lacking amendment were also set up. Then 20 ml of water was added to each bag, which were closed to allow a small hole to allow for gas exchange; the samples were then incubated at 25° C for 7, 14, 21 and 28 days elements determined as follows.

4.2.3.1. Determination of ammonium-N ($\text{NH}_4^+\text{-N}$) in dog faeces

Ammonium was extracted from both control(without larvae) and treatment (with larvae) dog faeces as well as to the modified control and treatment dog faeces (dog's compost) by adding solution of (15% w/v) of KCL in the ratio;(1g) of samples : (10 ml) KCL. The dog faeces were shaken for 30 min at 70 g then, in order to determine the concentration of ammonium N($\text{NH}_4^+\text{-N}$) 2 ml of filtrate was mixed with (1ml) of EDTA (6% w/v), then (7ml) of distilled water, (5ml)of phenolate reagent (see appendix), and (3ml) of sodium hypochlorite solution (10%v/v). The mixture was then incubated at 25C⁰ for 20 min in the dark. The volume was made up to 50 ml and mixed then the concentration of the indophenol-blue ammonium complex was measured at 630 nm (Wainwright and Pugh, 1973), the concentration of ammonium intensity was determined by reference to a standard curve(10-100µg $\text{NH}_4^+\text{-N}$) prepared from a standard solution of ammonium sulphate (see Appendix). Urea (0.5g) was added to the treatment samples (amended) dog faeces then while control was set-up lacking urea.

4.2.3.2. Determination of Nitrate – N (NO₃⁻-N)

Samples were extracted (1g) of each and shaken for 15 min with 20ml of water then the samples were filtrated through Whatman No 1 filter paper. Nitrate was determined using the method of (Sims and Jackson, 1971). Chromotropic acid: 7ml was mixed with 3ml of the filtrate then incubated at 40C⁰ in water bath for 45 min; the producer yellow colour was measured at 410 nm and the concentration of nitrate was determinate by reference to a standard curve of nitrate concentration.

4.2.3.3. Determination of Sulphate-SO₄²⁻- S

Samples (1g) were shaken with 10 ml of water then shaken at 70 g using an orbital shaker for 15 min the samples were filtrated through Whatman No.1 filter paper. Sulphate-S was determined using turbidimetric method to analysis the sulphur element (Hesse, 1971) as follows: Filtrate (10 ml) was transferred into a 250 ml volumetric flask, 1g of barium chloride BaCl₂, and 2 ml of gum acacia (0.25% w/v) were added and mixed well then the volume increased up to 25 ml of water. The white suspension resulting from precipitation of BaCl₂ was measured at 470 nm using a spectrophotometer and the concentration if of SO₄²⁻-S was determined by reference to a standard curve (0- 100µg sulphate ml) prepared from a standard solution of Na₂SO₄. (see Appendix).

4.2.3.4. Determination of phosphate solubilisation PO₄³⁻-P

Samples (10g) were placed into screw capped glass bottle containing 100 ml of 0.5 N NaHCO₃ and all the bottles were shaken for 30 min at 70 g, using an orbital shaker, the contents were then filtered through Whatman No. 1 filter paper. Phosphate ions were determined as described by Falih, (1995) as follows: Filter (3 ml) was mixed with 7 ml of work solution (see Appendix) then the mixture incubated at 37C⁰ for 1 hour. A

serial dilution were taken up to $\log 10^{-3}$ for the blue colour reaction then was measured at 820 nm using a spectrophotometer and the amount of $\text{PO}_4\text{-P}$ were determined by reference to a calibration curve ($0\text{-}8 \mu\text{g PO}_4\text{-P ml}^{-1}$).

4.2.4. Measurement of the pH of dog faeces samples

The pH of the dog faeces was determined each from zero time until 28 days, using a 1:1 distilled water suspension, shaken for 15 min on a reciprocal shaker (100 revolution min^{-1}), and a pH meter fitted with a glass electrode.

4.2.5. Determination of the effects of modified dog faeces as a compost additive on plant seed germination and growth

The two types of modified dog faeces (BSFL and FBL) mixed with a local grassland soil, which had not recently been fertilized, in the ratio 25%, 50% and 75%. Controls consisted of modified and unmodified dog faeces. Samples (100g) of each mixed proportions were planted with turnip or lettuce. The treatments were set up in triplicate and watered twice a week and left under laboratory condition and plant growth was observed.

4.3. Results and Discussion

4.3.1. Effect of larvae on the texture of dog faeces

Untreated dog faeces were greasy and tightly packed (Fig.4.1) and possessed an offensive odour. Treatment of fresh dog faeces with Black Soldier Fly and Fruits Beetle larvae reduced two 100g sample to 38g and 43g of modified dog's faeces within 4 weeks (62% and 57% reduction using BSF and FB larvae respectively. The modified dog faeces were more friable and loosely textured and had lost their faecal structure and their offensive odour (Fig.4.2).




Fig. 4. 1: Untreated dog faeces after 28 days incubation




Fig. 4. 2: Modified faeces: a) treated with FBL;
b) treated with BSFL after 28 days




Fig. 4. 3: The effect of BSFL activity on indigenous; a) ammonium, b) nitrate, c) sulphate and d) phosphate concentration in dog faeces(* Significantly different from control, $p=0.05$).




Fig. 4. 4: The effect of FBL activity on a indigenous a) ammonium, b) nitrate, c) sulphate and d) phosphate concentration in dog faeces (* Significantly different from control, $p=0.05$)




Fig. 4. 5: The effect of length of incubation on a) ammonium concentration following addition of ($10\mu\text{g}/(\text{NH}_4)_2\text{SO}_4\text{-N g}$), b) nitrate concentration following the addition of ($10\mu\text{g}/(\text{NH}_4)_2\text{SO}_4\text{-N g}$), c) S^0 oxidation following the addition of ($1\% \text{ w/w}$) S^0 and d) phosphate solubilisation following the addition of ($1\% \text{ w/w}$) calcium phosphate to dog faeces modified with BSFL (* Significantly different from control, $p=0.05$)




Fig. 4. 6: The effect of length of incubation on a) ammonium concentration following the addition of ($10\mu\text{g}/(\text{NH}_4)_2\text{SO}_4\text{-N g}$), b) nitrate concentration following the addition of ($10\mu\text{g}/(\text{NH}_4)_2\text{SO}_4\text{-N g}$), c) S^0 oxidation following the addition of ($1\% \text{ w/w}$) S^0 , d) phosphate solubilisation following the addition of ($1\% \text{ w/w}$) of calcium phosphate to dog faeces modified with FBL (* Significantly different from control, $p=0.05$)

Fig.4.3 shows the effect of BSFL activity on the concentration of the indigenous dog faeces-related plant nutrients: ammonium, nitrate, sulphate and phosphate. Fig (4.3a) shows that there was a significant reduction in ammonium concentration over the entire incubation period, while (Fig.4.3b) shows a similar decrease in nitrate concentration up to week three, followed by a significant increase at week 4. Sulphate concentration in the dog faeces also decreased following BSFL treatment over the entire incubation period (Fig.4.3c), while phosphate concentrations were significantly reduced up to week three and then significantly increased at week 4 (Fig.4.3d)

Fig.4.4 shows the effect of FBL on the indigenous plant nutrients in the dog faeces. Ammonium concentrations were reduced to week 2, but significantly increased at weeks 3 and 4 (Fig.4.4a). Nitrate concentrations, in contrast were significantly reduced over the entire incubation period (Fig.4.4b), as were sulphate concentrations; phosphate concentrations were significantly reduced up to week 3, but significantly increased at week 4 (Fig.4.4d).

The effect of amendment with BSFL modified dog faeces with a source of plant nutrients (ammonium, sulphate and insoluble phosphate) is shown in Fig.4.5. The concentration of all of the released nutrients (ammonium, nitrate, sulphate and phosphate) was generally significantly increased over the control, over the entire incubation period. The same trend was seen in nutrient-amended faeces modified by the addition of FBL (Fig.4.6).

As a generalization, the addition of both types of larvae to dog faeces significantly reduced the concentration of indigenous plant nutrients over the entire four week incubation period; exceptions to this were nitrate and phosphate concentrations in BSFL treated faeces, where significant increases were seen at week 4 and 3 respectively and in faeces treated with FBL, where ammonium concentrations were significantly increased at weeks 2-4, and phosphate at week 4. While the addition of both larvae therefore

initially decreased levels of indigenous plant nutrients there was a trend in some of the nutrients to increase the longer the incubation went on. This suggests that perhaps a longer term exposure of dog faeces to the two larvae might have led to increase in ammonium, nitrate, sulphate and phosphate concentrations. While this would generally be advantageous to the fertility of soils surrounding dog faeces, the potential long term increase in nitrate could at first sight be considered detrimental as nitrate can be readily leached to ground waters (where, when present in drinking water, it can cause blue baby disease and gastric cancer in humans). However, the relatively small amounts of dog faeces which are present, spread over large areas of soil, would likely make such increased nitrate contributions to drinking water relatively insignificant.

The addition of ammonium, elemental sulphur and insoluble phosphate to dog faeces which had been modified by the two larvae led to significant increases in nitrate, sulphate and plant-available phosphate, results which shows that dog faeces contains the indigenous microflora required for the transformation of these amendments (which simulate fertilizer addition). The increased friability and therefore increased aeration of the dog faeces following larval modification is also likely to result in enhanced rates of nitrification, S-oxidation and phosphate solubilisation. Clearly the microflora of dog faeces can act to modify added fertilizers and convert them to the plant available form.

4.3.5. Determination of the effects of modified dog faeces as a compost additive on plant seed germination and growth

Table 4. 1 shows that none of the seeds used germinated or grew in raw, untreated dog faeces (Control DF), but all grew in untreated compost (control compost). Of the seeds used only turnip germinated/grew in potting compost: modified dog faeces (FBL) at a ratio of 25:75. At a ratio of 50:50 lettuce seed failed to grow in BSFL modified faeces, but 80% of turnip seeds grew in this mix; 20 % of lettuce seeds grew in 50:50 FBL: compost and 40% of turnip seeds. At a ratio of 25:75 BSFL modified dog faeces: compost 20% of lettuce seeds grew and 100 % of turnip seeds grew. The growth figures for FBL modified faeces were relatively, lettuce 40%, and turnip, 80%. The results given in Table 4.1 are shown visually in Figs. 4.7 and 4.8

Table. 4. 1: The percentage seed germination and growth planted in different ratios of in modified dog faeces (treated with BSF and FB larvae) and potting compost

| Planted seeds | Control Modifies DF | 75% DF | 50% DF | 25% DF | Control compost |
|--|---------------------------|-----------|-----------|-----------|--------------------|
| Lettuce seeds planted in dog faeces treated with BSFL | 0% | 0% | 0% | 20% | 80% |
| Turnip seeds planted in dog faeces treated with BSFL | 0% | 0% | 80% | 100% | 60% |
| Lettuce seeds planted in dog faeces treated with FBL | 0% | 0% | 20% | 40% | 60% |
| Turnip seeds planted in dog faeces treated with FBL | 0% | 40% | 40% | 80% | 80% |

DF= Dog faeces




Fig. 4. 7: Shows a) the seeds planted in larval treated dog faeces; from 1 to 3 - turnip seeds, from 4 to 6 is lettuce seeds; b) the seeds planted in compost ; 7 to 9 - turnip seeds, from 10 to 12 is lettuce seeds.



Fig. 4. 8: Shows; a) the lettuce seeds planted in different concentration of larval treated dog faeces with BSFL No; 1, control dog faeces 2, 3 and 4 are 75, 50 and 25% of dog faeces concentration and 5 is a control compost. From 6 to 10 are the turnip seeds planted at the same concentration b) the lettuce seeds planted in different concentration of treated dog faeces with FBL No; 1, control dog faeces 2, 3 and 4 were 75, 50 and 25% of dog faeces and 5 is a control compost. From 6 to 10 showing turnip seeds planted at the same concentration

Clearly, by increasing the ratio of compost to modified faeces the inhibitory effect of raw dog faeces on plant growth was “diluted out”, thereby suggesting the possibility that larval modified dog faeces could be used as a compost additive fertilizer, or perhaps even be used as an agricultural soil fertilizer. These possibilities are however, limited by the fact that dog faeces are widely separated in the environment, so that it would be difficult to economically collect large amounts, making unviable its industrial use. Domestic dog pounds and dog racing kennels might on the other hand provide a sufficiently large source of faeces to make such use economic.

Chapter Five: The Antibacterial Activity of the Haemolymph and Whole Body Extracts from BSFL and FBL against Four Species of Bacteria

5. 1 Introduction

Like many other organisms, insects exhibit an efficient immunity response allowing them to survive in environment and it is known that insects have various levels of internal defence against microbial invasion (Cociancich *et al.*, 1994), including the ability, in some species, to externally secrete antibacterial compounds. Internal components of Green Blow Fly maggots for example, have been used with success, since 1930 (Simmons, 1935) to treat wounds infected with bacteria (Thomas *et al.*, 1999). In medicine, larvae of the Green Blow Fly (*Lucilia sericata* Meigen) are well-known for having the ability to reduce and eradicate bacteria in human wounds (Thomas *et al.*, 1996); they can both ingest and digest such bacteria or may produce active antimicrobial substances. A study by Murry and Hinckley (1992) showed that earthworms (*Eisenia foetida*) can also reduce the population of *Salmonella enterica* in horse manure.

The published work above shows a variety of results relating to the antibacterial activity of larval extracts against several species of bacteria. Bexfield *et al.*, (2004) established that *Lucilia sericata* excretions/secretion inhibit growth of *E. coli*. Similarly, they showed that *Lucilia sericata* extracts inhibit the growth of *P. aeruginosa* over a 24-hour period, whereas Jaklič *et al.*, (2008) reported a prolonged lag phase of more than 5-hours, and Thomas *et al.*, (1999) found only limited inhibition.

More recently, Mumcuoglu *et al.*, (2001) found green fluorescent protein (GFP)-producing *E. coli* in the alimentary canal of Fly larvae and found a decreasing intensity of this marker during passage through the digestive tract. A battery of defence proteins synthesized by insects in response to bacterial challenge may be responsible for this decrease. Cecropins, and lytic proteins for example, have been isolated from larvae of blow flies (*Calliphora vicina*) (Crowley and Houck, 2002). Another antibacterial substance, *p*-hydroxycinnamaldehyde, isolated from the larvae of the Saw Fly,

Acantholyda parki S., has also been found to have a broad antibacterial spectrum against both gram-negative and gram-positive bacteria (Leem *et al.*, 1999).

The Black Soldier Fly, *Hermetia illucens*, is a beneficial insect because its larvae feed on organic material, including the remains of plants, animals and humans. Such larvae can degrade large amounts of waste quickly and efficiently than any other known species of fly due to their potent mouthparts and digestive enzymes (Tomberlin *et al.*, 2002). The effect of larval excretion/ secretion (ES) against some common species of bacteria found in dog faeces was studied in relation to determining how larvae can resist bacterial invasion. Based on these findings, similar antibacterial substances might be present in BSF and FB larvae.

The investigations reported in this Thesis were aimed at determining the ability of Black Soldier Fly and Fruit Beetle larvae to reduce the number of pathogenic bacteria, such as *E. coli*, *Salmonella enterica* and *Staphylococcus aureus* MRSA, in dog faeces. Few studies have reported the ability of Black Soldier Fly or Fruit Beetles to demonstrate immunity or the ability of their internal and external secretion, as antibacterial agents.

The aims of the work reported in this Chapter was to investigate the presence of antibacterial activity in the body extracts of BSF and FB larvae and their inhibitory effects against four species of bacteria.

5.2. Materials and Methods

5.2.1. Culture of larvae

The two species of insects (larvae of Soldier Fly and Fruit Beetle) used in this study were obtained from a commercial company (Ricks, Livefood.Co.UK), the third instar of BSF larvae was purchased, while large sized FB larvae were used.

5.2.2. Antibacterial properties of BSF and FB larvae

5.2.2.1. Collection of excretion/secretion (ES) from larval whole-body and haemolymph

Samples were placed separately in small boxes (10×15×30 cm) under laboratory conditions. In order to collect the whole body and haemolymph extraction, samples 20g of each larval species (approximately 200 larvae of BSF and 10 larvae of FB) were left unfed for 4 days until they were free from gut residues. For whole body extracts and larval secretions, larvae were washed with sterile deionised water (dH₂O) and placed in Petri-dish at -4°C for a few days. The larvae were then ground manually, using a mortar and pestle, until they became homogenous. Haemolymph was collected from larvae by clipping the anterior end near the cephalopharyngeal skeleton. The abdomen was then squeezed gently to force the haemolymph to flow from the wound. The resultant suspensions were then filtered (0.2µm) to remove any large particles and bacteria. The excretion/secretion ES from each group of insect larvae were dissolved in 0.1 µg/ml protein in 0.1% trifluoroacetic acid (TFA) 1: 10 v/v. The suspension was placed on ice for 6 hours and centrifuged four times at 14000 g for 30 min. The supernatant was then harvested and separated into 15 ml centrifuge tubes; samples were used fresh or kept frozen at -80°C for a few days until required (Sahalan *et al.*, 2006; Huberman *et al.*, 2007).

5.2.2.2. Antibacterial assay

Antibacterial activity was determined as follows: Viable counts: the effect of BSF and FB extract on the viability of *E. coli*, *Salmonella enteric*, *Serratia marcescens* and *S. aureus* MRSA cells were monitored by inoculating 1ml of an overnight 5.0 x10⁶ cfu/ml, into 9 ml nutrient agar with and without both extracts and incubating with shaking at 37°C (250g). Samples (100 µl) were removed at intervals, diluted serially and 100µl

of the diluted samples were placed on nutrient agar plates and incubated at 37°C for 24 hours. The colonies were then counted after overnight incubation and the results expressed as colony forming units (CFU/ml). Turbidimetric analysis: bacterial cells were incubated the extracted haemolymph and whole body; the final optical density of the treated and untreated cultures were then measured at 600 nm.

5.2.3. Time killing curves for testing the bacterial activity

The time kill assays were prepared by inoculating one or two colonies of overnight test bacteria into 9 ml of Mueller–Hinton broth and incubating for 18-24 hours at 37°C with shaking, adjusting to a 0.5 McFarland 5×10^6 CFU ml⁻¹. Haemolymph and whole body were prepared separately by adding 4ml of haemolymph to 16 of ml of Mueller–Hinton broth in 100 ml flasks, solutions lacking haemolymph acted as controls. Treatments and control were incubated with shaking at 37°C and samples were transferred to nutrient agar at 0, 2, 4, 6 and 8 hours. Viable counts were performed by serially diluting each sample 10-fold in PBS and spreading 100 µl volumes from the appropriate dilutions onto nutrient agar. After incubation at 37°C, for 20–24 h, viable counts (CFU ml⁻¹) were determined to provide a quantitative determination of bacterial growth colony forming units. To confirm the sterility of samples of ES prior to bacterial inoculation, a loop (10 µl) of haemolymph and whole body secretions from each larvae, TSB, and sterile dH₂O were spread separately onto nutrient agar and incubated at 37°C for 24 h. This procedure was repeated at four, eight and 24 h to ensure the media used to prepare dilutions for viable counts was contamination-free.

5.2.4. Bacterial culture

Four species of bacteria were used in this study. Three species were Gram negatives (*Escherichia coli*, *Salmonella enterica* and *Serratia marcescens*) were obtained from previous works, whereas the fourth, *Staphylococcus aureus* was a Gram positive. The organisms were obtained from the Departmental Culture Collection.

Bacteria were maintained on nutrient agar at 4°C. New subcultures were prepared as required (Alexander and Strete, 2001). In order to test that the colonies were viable, the plates were incubated overnight at 37°C to allow growth.

5.2.5. Transmission Electron Microscopy of Bacteria treated with haemolymph or whole body secretions of the two larvae

For transmission E/M, harvested cells were fixed in 3% glutaraldehyde and 0.1M phosphate buffer overnight at 4°C, and then washed twice times with 30 min intervals at 4°C in 0.1M phosphate buffer. Cells were subjected to a secondary fixation with 2% osmium tetroxide aqueous for 2 hour at room temperature and the previous wash step was repeated. After this, cells at room temperature were serially dehydrated with 75%, 95%, and three times using 100% ethanol, the last one was dried over anhydrous copper sulphate for 15 min. The cells were then placed in an intermediate solvent, propylene oxide, for two changes of 15 min duration. Infiltration was accomplished by placing the cells in a 50/50 mixture of propylene oxide/Araldite resin. The cells were then left in this 50/50 mixture overnight at room temperature. Once this incubation was finished the cells were transferred into full strength Araldite resin and left for 6-8 hours at room temperature after which time they were embedded in fresh Araldite resin for 48-72 hours at 60°C. Semi-thin sections approximately 0.5 µm thick were cut on a Reichert Ultracut E ultramicrotome and stained with 1% toluidine blue in 1% borax. Ultrathin sections, approximately 70-90nm thick, were then cut using a Reichert Ultracut E

ultramicrotome and stained for 25 mins with 3% uranyl acetate in 50% ethanol followed by staining with Reynold's lead citrate for 25 min. The sections were examined using a FEI Tecnai Transmission Electron Microscope at an accelerating voltage of 80Kv.

5.2.6. Statistical analysis

All data were presented as means \pm SD. (Standard deviation). The data was analyzed by SigmaPlot[®] 11.0. $P < 0.05$ was considered as significant.

5.3. Results and Discussion

Figures 5.1 and 5.2 show the effects of the haemolymph and whole body extract of the BSFL and FBL respectively on the growth of *E. coli*. Both the BSFL extracts had only a minor effect on the growth of *E. coli* from 2-8h (Fig 5.1). After 8h however, BSFL haemolymph had a marked effect on *E. coli*. In contrast the addition of the haemolymph and whole body extracts of the FBL produced an increase in numbers of *E. coli* compared to the control, from hour 2-8 (Fig.5.2).




Fig. 5. 1: The effect of BSFL haemolymph and whole body extract on *E. coli*, growth curve over 8 h.




Fig. 5. 2: The effect of FBL larvae haemolymph and whole body extract *E.coli* , growth curve over 8 h.




Fig. 5. 3: The effect of BSFL haemolymph and whole body extract on *S. enterica* growth curve over 8 h.

Both the haemolymph and whole body extract of the BSFL inhibited the growth of *S. enterica* at hour 2 of the incubation period, although by hour 4, there was little impact of both extracts on bacterial numbers; the inhibition trend for both bacteria then resumed from hours 6-8 (Fig5.3).

Haemolymph of the FBL had a general slight stimulatory impact on numbers of *Salmonella enterica* over the entire incubation period, with the effect being at its lowest in the middle of the 8 hours. The whole body secretion in contrast, reduced bacterial numbers over the entire incubation period (Fig.5.4).




Fig. 5. 4: The effect of FBL haemolymph and whole body extract on *S.enterica* growth curve over 8 h.




Fig. 5. 5: The effect of BSFL haemolymph and whole body extract on *S.aureus* MRSA, growth curve over 8 h.

Both of the extracts from the BSFL had a general inhibitory effect on the growth of *S. aureus*, the effect being most marked with the haemolymph, which inhibited growth over the entire incubation period (Fig.5.5). The haemolymph extract from the FBL inhibited the growth of MRSA between 4 and 8 hours (Fig.5.6), while the whole body extract increased numbers at 6hours.




Fig. 5. 6: The effect of FBL haemolymph and whole body on *Staphylococcus aureus* MRSA, growth curve over 8 h.




Fig. 5. 7: The effect of BSFL haemolymph and whole body on *Serratia marcescens*, growth curve over 8 h.

Both the haemolymph and whole body extract of BSFL inhibited the growth of *S. marcescens* over the entire incubation period, the effect of the haemolymph extract being most marked at the end of the growth period ((Fig.5.7). The whole body extract of the FBL had less effect on the growth of *S. marcescens* over the entire incubation period, while the haemolymph had a slight, and very transient stimulatory effect, at the 6h point only; the haemolymph increased numbers at the end of the incubation period (Fig.5.8).




Fig. 5. 8: The effect of FBL haemolymph and whole body extract on *Serratia marcescens*, growth curve over 8 h.

Table. 5. 1:The effect of the two species of larval haemolymph and whole body secretions on the four species of bacteria.

| Bacterial strains | Gram strain | BSF larvae | | FBL larvae | |
|----------------------------|-------------|------------|------------|------------|------------|
| | | Haemolymph | Whole body | Haemolymph | Whole body |
| <i>E. coli</i> | Gram -ve | + | + | - | - |
| <i>Salmonella enterica</i> | Gram-ve | - | - | - | + |
| <i>S. aureus MRSA</i> | Gram +ve | + | - | + | - |
| <i>Serratia marcescens</i> | Gram -ve | + | + | - | - |

Table. 5.1 shows that the effect of the extracts on bacteria varied with different species of larva and with different types of extracts used. Clearly, no overall generalization can be made about the effects of BSFL and FBL extracts on bacteria can be observed.




Fig. 5. 9: The effect of larval components on the morphology of *E. coli* observed by TEM, a) control for BSFL, b) whole body extract of BSFL, c), haemolymph of BSFL; d) control for FBL, e) whole body extracts for FBL and g) haemolymph of FBL.




Fig. 5. 10: The effect of larval components on the morphology of *S. aureus* observed by TEM, a) control for BSFL, b) whole body extract of BSFL, c), haemolymph of BSFL; d) control for FBL, e) whole body extracts for FBL and g) haemolymph of FBL.

The effects of the two larval extracts on *E.coli* and *S. aureus* as observed under the transmission electron microscope are respectively shown in Figs. 5.9 and 5.10. The haemolymph extract of BSF is the only one which has a readily observed, and marked, effect on the morphology of *E.coli* (Fig.5.9c); note that the cells are seen to lose some of their electron dense structure, but that cell lysis is not evident. In the same way, the haemolymph of BSFL is the only fly body extract which has an effect on the cell structure of *S. aureus*; again there is no obvious evidence of cell lysis (Fig.5.10c).

Interest in the possible antibacterial activity of the body extracts of the two flies used here was based on two possibilities. Firstly that an antibacterial agent, possibly a protein might be obtained from either or both of the extracts and secondly that BSFL or FBL might be used in infected wounds in the same way that larvae of the Green Blow Fly (*Lucillia sericata*) are used in maggot therapy (Boxfield *et al.*,2004). The latter possibility is perhaps unlikely based on the large size of the two fly larvae compared with maggots, a fact which would restrict their application to open wounds. Maggot therapy is extremely effective, so it is unlikely that the two larvae studied here would replace this approach to wound treatment. Similarly while extracts from the two larvae have been shown to inhibit the growth of bacteria, their effectiveness is unlikely to be comparable to antibiotics like penicillin.

Conclusion: In conclusion, whole body extracts and the haemolymph of the two fly species used here do have antibacterial effects, but such inhibitory effects are not consistent, both in relation to the individual fly species, or in relation to the type of bacterium investigated. The addition of the fly larvae to faeces is likely however, to lead to reduction in overall bacterial numbers. It seems unlikely that BSFL or FB larvae could be usefully applied to the control of bacterial infections in human or animal wounds.

Chapter Six: The Number of Bacteria in Grassed Soil of Public Parks Contaminated with Dog Faeces and Effect of Faeces-Volatiles on Plant Growth.

6. 1. Introduction

The number of dogs in the UK has increased dramatically in recent times especially in cities, and their waste represents a considerable source of pollution in relation to humans and the environment (Carvalho *et al.*, 2009). Dog faeces are known to carry a large and various microbial and parasite population which can cause diseases and public-health problems (Baxter and Leck, 1984). Dog waste is linked with more than 60 zoonotic diseases and provides a serious public-health problem (Rinaldi *et al.*, 2006). Irresponsibility on the part of dog owners in relation to the disposal the dog waste represents a source of potential pathogens in public areas such as playgrounds, parks, gardens, public squares and sandpits (Schantz, 1994; Rubel and Wisnivesky, 2005); the risks to young children being particularly obvious. Children may, by accident consume contaminated soil or grass, or touch their mouths or eyes with hands contaminated with dog faeces or otherwise handle bags left hanging outdoors on trees branches and in parks (Fig.6.1) ; people using hand activated wheelchairs and active sports players may also be at risk (Jackson, 1995). Viable pathogens present in dried canine faeces can also be spread by the wind and are carried into dwellings and workplaces on soiled shoes.

The aim of the work reported in this Chapter was to study faecal contamination accumulation in a local public park, a problem which is likely to be a risk to the health of young children and parks visitors. In addition, the work focuses on the extremely undesirable habit of some dog owners to remove dog faeces from the environment and leave them in a plastic bag on the parks, a habit which contributes to a potentially bigger problem than does exposed faeces. The effect of dog faeces volatiles on plant growth will also be touched upon.

6.2. Materials and methods

6.2.1. Samples collection from public parks

The following protocol was used to determine the presence of *E. coli* and *Salmonella enterica* in addition to other heterotrophic bacteria (as determined using Plate Count Agar) in soil samples collected from a public park in Sheffield UK (a grassed area of playing field behind the former Lodge Moor Hospital). Four samples of soil were taken (0 - 3 cm) from under surface grassed soil contaminated with dog faeces (samples were taken directly from soil which had long exposure to faeces and not from under fresh material); the last sample was taken from an area not contaminated with faeces, i.e. an uncontaminated control. Samples (1 g) were suspended in (9 ml) autoclave-sterilised water in sterile 15 ml tubes and shaken at 70 g for 30 min. A serial dilution was performed, then dilutions from 10^3 to 10^6 were taken and (100 ml) of this suspension was then spread onto the surface of selective media for *E. coli* and *Salmonella* Spp. XLT4 and Plate Count Agar; the plates were then incubated in triplicate at 37°C overnight. Presumptive isolation was based on the use of selective media such as *E. coli* and XLT4 *Salmonella* Spp. agar media, followed by use of the Gram stain. In addition, the 16s rRNA gene dependant technique was used; DNA being extracted according to the Anachem Key prep-Bacterial DNA extraction kit; bacterial 16SrRNA gene was then amplified and sequenced.

6.2.2. Effect of incubation of dog faeces in plastic bags left under ambient outside conditions

A variety of fresh dog faeces were collected from dog bins from Weston Park. Samples were placed in black plastic bags and left exposed to ambient weather conditions on the roof of the Firth Court building from the middle of May until the middle of June, 2012. Samples (1 g) of the faeces were diluted in (9 ml) in sterilised

water. Then a serial dilution from 10^3 to 10^7 was performed and (100 μ l) of the final suspension was then spread onto plates contained the selective media: HiCrome *E. coli* agar and *Salmonella Spp* XLT4, as well as Plate Count Agar. Plates were incubated in triplicate at 37°C for 18-24 h.



Fig. 6. 1: Plastic bags containing dog faeces bags obtained from Sheffield parks

6.2.3 The effect of gasses (odour) from dog faeces on plant growth

Two plastic containers were prepared, one contained compost soil divided to two parts each one include 200 g of compost into which was planted lettuce seeds, and the other turnip; controls not receiving faeces-gasses were also set up. The gasses were transferred, using a pump, from a large Erlenmeyer flask containing 200g dog faeces. The gasses then passed through holes in the bottom of the plant growth container and passed over the growing seedlings (Fig. 6.2). Seeds (50 each of lettuce and turnip) were planted and watered twice a week and left at room temperature in the light. The experiment was continued for two weeks when the growth was visually assessed, i.e. for four weeks. The samples were aerated and suspended with unoccupied space under natural conditions.




Fig. 6. 2: Airtight container contains 200 g of fresh dog faeces, the dog faeces gases were pumped via the tube into the plastic box containing compost planted with lettuce or turnip seeds.

6.3 Results and Discussion




Fig. 6. 3: The presence and population density of *E. coli* and *Salmonella sp* in five locations in public areas of Sheffield (location 5 is the uncontaminated grass sample).

6.3.1. Occurrence of *E. coli* and *S. enterica* in dog faeces contaminated grassed-soil

Escherichia coli and *Salmonella enterica* were isolated from all four sites (Fig. 6.3), while no isolates were obtained from the fifth location which was uncontaminated with dog faeces. The number of *E. coli* varied from site to site (from 10^4 to 10^5 CFU/g); *E. coli* was isolated in highest numbers. The isolation of these pathogenic bacteria from playing fields frequented by children is obviously a major public health concern since

pathogenic bacteria could be picked up from these soils, either on shoes, or by transfer from contaminated skin to the ears, eyes or mouths of infected children (and of course, people in general) from where they could cause disease.

Several studies have concluded that there is a high potential risk to human health of contaminated soil and animal faeces, and direct or indirect contact with dog faeces has been implicated in several human infections with *E. coli* and *Salmonella enterica* (Kudva *et al.*, 1998). The risk of the transfer pathogenic bacteria to young children from dog faeces in public areas-notably gardens, parks and playing-field probably is critical because people do not always know that they have been exposed to contamination and can touch their eyes or mouth with faeces without realising it. Contaminated garden and park soils could possibly influence the health of large number of adults and children and it is therefore important that research and development is carried out on risk assessment and also to develop measures to mitigate such contamination (Alloway, 2004).

6.3.2 Effect of leaving dog faeces in plastic bags on bacterial numbers

An undesirable and unhygienic habit has recently developed amongst some dog owners in Sheffield (and presumably other parts of the country) of picking up their dog waste, transferring it to polythene bags and then leaving the filled bags in the environment as litter, rather than depositing them in dedicated dog-waste containers.

A wide range of potentially pathogenic bacteria were isolated from dog faeces left at ambient conditions in plastic bags (Fig.6.4). The results presented in Fig.6.5 show that numbers of *E.coli* and *S. enterica* increased in the dog faeces, left in sealed plastic bags, over the 28 day incubation period; as did the number of total heterotrophic bacteria.







Fig. 6. 4: Bacterial species isolated from dog faeces left in polythene bags; a) red colonies of *E. coli*, and blue *Enterococcus Spp*, b) *E. coli*, c) *Salmonella enterica*, d) *Staphylococcus aureus*.




Fig. 6. 5: Numbers of *E. coli*, *Salmonella enterica* and total heterotrophic bacteria (grown on plate count media) isolated from dog faeces left in plastic bags.

This study shows that the numbers of “total bacteria”, as well as those of two important human bacterial pathogens increase in dog faeces left under ambient conditions in sealed plastic bags. It would have been interesting to continue the exposure period for longer than the one month used here , since numbers of pathogenic bacteria are likely to have increased even further under such circumstances.

6.3.3 The effect of the gasses (odour) released from dog faeces on the plant growth

Fig.6.6 shows that gasses from dog faeces have a marked detrimental effect on the growth of lettuce and turnip from seed, with seedling growth obviously visually reduced in treated samples when compared to the controls. The question could be asked- why

was the effect of dog faeces volatiles on plant growth determined, and not the direct effect of the faeces themselves? The answer is that it would have been difficult to separate the likely toxic effects of dog faeces, on plant growth, from the positive fertilizer effect resulting from the fact that the faeces are rich in plant nutrients like ammonium and nitrate. Although the nature of the dog faeces gases and volatiles was not determined, the observed inhibitory effects presumably result from the inhibition of seed germination or shoot growth, following the production of a cocktail of toxic gases, including ammonia in high concentration (Van der Eerden, 1982). Several studies have discussed emissions from animals and researchers have obtained varying results; clearly research is needed to discover the impact of gases, emitted from animal manures (such as ammonium) on plants (Kebreab *et al.*, 2006). The effect of ammonia on vegetation has been noticed since 1896, when König observed injuries to the surrounding vegetation near a soda factory where ammonia was accidentally released. Garber (1935) appears to have been the first to expose plants to NH_3 and to prove its phytotoxic effects (Fangmeier *et al.*, 1994). Ammonia can cause various types of injury, including necrosis, growth reduction, growth stimulation and increased frost sensitivity. Several plant species have been assessed for sensitivity to ammonia. Some conifer species were relatively sensitive to low concentrations in the long term; some cultivars of cauliflower and tomato were relatively sensitive to somewhat higher concentrations for a short term. Special attention has been paid to plant injury around intensively managed livestock. The emission from these sources consists of a large number of compounds, ammonia proving to be the main toxic component (Van der Eerden, 1982).




Fig. 6. 6: The effect of gas (odour) emission from dog faeces on vegetable growth; a) turnip treated with dog faeces gasses and, b) lettuce treated with dog faeces gasses c) control turnip seeds d) control lettuce seeds.

Dog faeces represent a potential source for spreading pathogenic bacteria such as *E. coli* and *Salmonella enterica*, bacteria which represent an obvious risk human health. Bacterial pathogens from dog faeces could infect young children and others playing, or using, parks etc (including of course dog owners). Dog faeces have also been shown

here to produce gasses which are injurious to plant germination and/ or growth, and are also likely to be directly inhibitory to plant growth.

There has been a welcomed trend of late for dog owners to act responsibly by picking up waste from their dogs, placing this in plastic bags and than putting these bags into special local council run dog-waste containers, from where they are transported and incinerated (Fig.6.7).



Fig. 6. 7: A typical dog waste bin used in public parks in the UK (author's photograph).

Unfortunately there is another increasing trend, namely the discarding of such bags filled with dog faeces into the local environment. The results presented here show that this practise potentially increases the problem of dog faeces because the sealed polythene bags provide an environment where pathogenic bacteria (and presumably other pathogens, like viruses) can rapidly increase. Such an increase in pathogen numbers was seen during a relatively cool UK summer and could be much greater where summer temperatures are higher. In addition, the possibility exists that the environment inside the bags could become anaerobic and select for, or otherwise encourage the growth of anaerobic, pathogenic bacteria, such as species of *Clostridium*. Clearly this habit should be discouraged. In fact it would be better to encourage dog

owners to leave dog faeces in the open air, where they can break down naturally (with pathogens being outcompeted by saprophytic microbes) than to allow them to discard the faeces in sealed bags which, when broken (by for example a child sliding on them) could release an increased, and potentially more dangerous, pathogen load.

The survival of bacteria in animals waste under environmental condition has been studied by, amongst others, LaGoy, (1987), who showed that *E. coli* in cattle manure survived for 42 to 49 days at 37°C, for 49 to 56 days at 22°C, and for 63 to 70 days at 5°C (LaGoy, 1987) Another study of *E. coli* in manure commutation revealed that the pathogen may survive for up to 47 days, 4 months, and 21 months in bovine, aerated ovine, and non-aerated ovine manure, respectively (Kudva *et al.*, 1998). Himathongkham *et al.*, (1999) based on their research findings, recommended that cow manure should be survive for 105 days at 4°C or 45 days at 37°C to achieve a 5-log₁₀ reduction of both *E. coli* and *Salmonella enterica* serovar (Himathongkham *et al.*, 1999). These results indicate that *E. coli* can continue in bovine faeces for an extended period of time and that bovine faeces are a potential vehicle for transmitting the pathogen to cattle, food, and the environment (Jiang *et al.*, 2003). However, pathogenic bacteria such as *E. coli* and *Salmonella Spp.* can survive in animals manure or soil up to 70 days under 14⁰F (McLaughlin, 2002). A study by Kudva *et al.*, (1998) reported that *E. coli* survived in the manure for 21 months, and the numbers of bacteria recovered ranged from <10² to 10⁶ CFU/g at different times over the course of the experiment.

Conclusion: In conclusion dog faeces represent an important means of transfer of bacterial pathogens from dogs to humans via the soil. The reprehensible recent habit amongst some dog owners of leaving faeces in sealed plastic bags is likely to greatly exacerbate this problem.

Chapter Seven: Production of Biodiesel from Dog Waste by BSF and FB Larvae- A Potential Solution To The Dog Faeces Problem

7.1. Introduction

Biodiesel is receiving increased attention as a renewable source of fuel. Biofuels can be made from a variety of feedstocks including starch, vegetable oil, or animal fats as an fuel alternative to help reduce consumption of petroleum. Alternatively cheap waste, such as dairy manure can be used to produce bioethanol (Liao *et al.*, 2008; Predojevic 2008) and is hoped that, as well as producing biofuels, a reduction can be made in the amount of waste which has to be disposed of to the environment (Ann *et al.*, 2002). In addition to biodiesel, other fuels can be obtained from wastes as is discussed below.

7.1.1. Fuels from wastes

Many fuels can be derived from waste refuse can be combusted directly in an incinerator to provide power and chicken litter can (Ferrer *et al.*, 2005) , like dog faeces, be dried and directly combusted (Li *et al.*, 2011).

7.1.2. Anaerobic digestion to methane

The anaerobic fermentation of animal fats, vegetable oil, starch and sugar can be achieved using conventional technology and results in the production of methane, i.e. biogas (Demirbas, 2008). Digestate, a solid by-product which can be used as a fertilizer is also produced. Dog faeces can fuel an anaerobic digester and produced methane. This might be done in a large centralized facility, but it is more likely that it will be done on a more local scale. Methane is of course a more potent green house gas than carbon dioxide and when burnt its releases CO₂; any treatment will however solve the direct problem of dog waste removal. In the US, and increasingly world-wide, dog waste is being turned into a local energy source (Demirel *et al.*, 2010). Dog owners throw the waste into a feeding tube, turn on a hand crank, so that the anaerobic biogas digester can decompose the faeces and make it into a burnable methane gas. The gas is then used to

power the local street. i.e. the so-called “Park Spark Project”. A criticism of this approach is that during methane production, pathogens are not removed completely a potentially harmful product will remain after digestion; such residues can however, be treated in the conventional sewage treatment system (Demirel *et al.*, 2010).

Biofuel or agrofuel is obtained from biomass and may be produced in a solid, liquid or gaseous form. The utilization of waste biomass as a source of energy can decrease problems relating to waste management , pollution, greenhouse gaseous emissions and, as a result, the burning of fossil fuels. Some 19 million tons of oil equivalents could be derived from biomass, with some 46% being obtained from bio-wastes like farm waste, agricultural waste, municipal solid waste and other biodegradable waste streams (Demirbas., 2008)

7.1.3. Landfill methane

Landfill waste undergoes anaerobic digestion to landfill gases (LFG). Such gases can be burned and looked up to as a source of renewable energy (Themelis and Ulloa., 2007). The LFG is made of around 50% methane and since it is equivalent to natural gas can be used to generate electricity for public consumption or alternatively it can be burned for heat. If LFG is not used properly, it would be released to the atmosphere. and since methane is a greenhouse gas, it has a global warming potential of 23 times carbon dioxide, i.e. one ton of methane results in the greenhouse gas equivalent of 23 tons of carbon dioxide. Harvesting and burning LFG reduces the global warming potential by a factor of 23 and energy is provided for heat and power. A typical landfill power plant can supply power to 1900 homes and avoid 6000 tons of methane per year from entering the atmosphere. As well as eliminating 18,000 tons per year of carbon dioxide by fossil fuel replacement (De Montfon, 2012)

7.1.4. Biodiesel from waste cooking oils

Each week the UK population uses in excess of 1,500 tonnes of cooking oil. Large scale producers of waste oil like restaurants, chip shops and industrial food producers have their waste oil removed by licensed collectors, and around 100,000 tonnes of recoverable waste oil produced each year within the UK (Phan and Phan., 2008). Currently most of this recovered oil is treated and added to animal feed. The EU however, its use for pig feed due to several food scares related to vegetable oil and a similar ban may soon apply to cattle also. Such oil can however, be converted into biodiesel (Silvis, 2006). The waste vegetable oil currently used to produce UK biodiesel is mostly rapeseed oil, whose trans-esterification produces RME (Rape Methyl Ester), or more properly, FAME (Fatty Acid Methyl Ester) which is the correct chemical term for biodiesel produced from any oil or fat using methanol as the alcohol for the production process (Zhang *et al.*, 2003).

7.1.5. Methane from anaerobic digestion

Anaerobic digestion (or gasification) is the use of bacteria to decompose organic matter in the absence of oxygen to produce a gas containing 60% methane, 40% carbon dioxide together with a mix of solid and liquid fertiliser, and since anaerobic digestion does not produce anymore carbon dioxide than would be produced by the natural decomposition of the waste, it regarded as being 'carbon neutral' In this process, waste is pumped once a day into the digester, and remains there for about 10 to 40 days, when the internal of the digester must reach between 30 and 70 degrees C. Any mix of organic waste mater can be used, including animal and human sewage, crop residues, newspaper, abattoir waste and agricultural and food processing waste. Since the resultant product is a gas this fuel is more applicable to electricity generation, rather

than a transport fuel; although methane could be liquefied for vehicular use (Chynoweth *et al.*, 2001).

7.1.6. Oil from tyre pyrolysis

Around 40 million tyres (440,000 tonnes) per year are produced world-wide. Currently the UK recovers some value from 70% of the total used tyres produced, with the rest going to landfill. In 2006 a ban on tyre-landfill was introduced, so tyres are increasingly being treated by pyrolysis, a process of thermal degradation in the absence of oxygen at elevated temperatures (430°C) and pressures (Shulman, 2004). The products of pyrolysis (from organic waste) are gases, small quantities of liquid, and a solid residue containing carbon and ash. The gases produced in the process can then be re-used to provide the heat required to continue the process (Makarov and Drozdovskiĭ., 1991).

7.1.7. Biodiesel production

Currently, biodiesel is generally used as a diesel fuel additive to minimize particulate release and the production of carbon monoxide, and hydrocarbons. The relatively high production cost is the main reason why biodiesel is not used as a primary fuel. Three quarters of the cost of biodiesel is derived from the feedstock, such as soybean oil, rapeseed oil and sunflower oil. In addition, the use of limited food supplies for the production of biodiesel is not feasible for developing countries like China. As a result, non-food feedstocks such as *Jatropha curcas*, Chinese tallow and microalgae are being developed as sources of biofuels; the use of these sources however, is certainly not without problems. On the other hand, organic wastes such as animal wastes, residential wastes (e.g., household), commercial and institutional wastes are generated in large quantities in developing countries and present a cheap form of organic wastes which would otherwise present an environmental pollution problem and so can act as a useful resource for biodiesel production (Ma and Hanna., 1999)

7.1.8. Biodiesel production by BSFL

Animal faeces waste can be used for biodiesel production, with the resultant biodiesel being comparable to rapeseed based biodiesel, and met the European biodiesel standard, EN14214. Among the organic wastes evaluated, chicken manure appears to be the best one for achieving high rates of BSFL biomass production. About 30% of the waste-grown BSFL biomass can be extracted as crude fat, which can be used for the production of biodiesel; the resultant fuel properties of the BSFL fat-based biodiesel, including density (885 kg/m³), viscosity (5.8 mm²/s), and cetane number (53) were found to be similar to those of rapeseed oil based biodiesel (Li *et al.*, 2011a). To date there appear to be no reports of the use of dogs faeces to produce biodiesel either alone or in conjunction with BSFL or FBL. Among the organic wastes evaluated chicken manure was found to be the best for maximal BSFL growth (327.6 g), which gave 8.5 g crude fat production after petroleum ether extraction. An optimized two step conversion process was performed, yielding 91.4 g of biodiesel with a biodiesel yield of 93% from the crude fat contents (Li *et al.*, 2011c). Cattle, pig and chicken manure can also act as a food source for the production of high fat, nutrient-rich larvae. Fat-rich BSFL can be used to degrade organic waste and produce biodiesel as well as providing foods from animals notably chickens (Li *et al.*, 2011b).

The results of recent studies have shown that BSFL-produced fatty acids could be a valuable feedstock for biodiesel production having two advantages over crop oil-based biodiesel, namely:

- 1) It does not compete with food resources or land use
- 2) It uses “waste nutrients” for insect growth.

The aim of the work reported in this Chapter was to determine if the larvae used here can convert dog faeces into a potentially viable source of biodiesel

7.2. Materials and Methods

7.2.1. Samples collection

Black soldier fly and fruit beetle larvae were obtained from LiveFood Co., while dog faeces samples were collected from local dog waste bins. The faeces were distributed in two boxes, each containing 100g of dog faeces. Ten BSFL were then added to the waste (1000 larvae per kg dog waste) according to Li *et al.*, (2011) and ten FB larvae were similarly added to the dog faeces. The treated faeces were incubated under laboratory conditions for 28 days; samples were obtained in triplicate at weekly intervals.

7.2.2. Extraction of lipids

In order to extract lipids from dog faeces and larvae, the larval whole body were ground and then weekly samples (1 g) of all samples were added to Eppendorf tubes 1.5 ml and frozen at -80°C overnight, then freeze dried (lyophilized) for 24-48 hours. To estimate the weight of biomass the Eppendorf tubes were re-weighed. Samples were transferred to centrifuge tubes (50 ml), 20 ml of methanol/ chloroform (2:1 v/v) were added; the contents were then sonicated for 1 min on ice. Samples were centrifuged at 5000 g for 5 mins and transferred to a centrifuge tubes in order to determine the volume of supernatant. Then 2:1 methanol: chloroform, and chloroform and 1% NaCl (1 g NaCl in 100 ml) was added to give 2:2:1 methanol: chloroform:1%NaCl. (13.33ml methanol:13.33 chloroform:6.7 1%NaCl) (Fig 7.1). Samples were then centrifuged for 2 mins at 5000 g. The centrifuge tubes were labelled and weighed then transferred to a chloroform phase into pre-weighed sample were left in fume cupboard with tops open to evaporate until dry. The centrifuge tubes were then re-weighed and the weight of lipids calculated. (Lu *et al.* 2008; Chiu *et al.* 2009)

7.2.3.Detection and measurement the fatty acid using a combined mass spectrometer and gas chromatographic GC/MS

The harvested lipids (10mg/ml) of dog faeces and BSF and FB larvae before and after being fed on dog faeces were dissolved in chloroform. The samples were then subjected to gas chromatography-mass spectrometry GC/MS. The GC/MS analyses were performed using a Perkin Elmer Turbo mass spectrometer detector (software version 5.4. Perkin Elmer). Injector and oven temperatures were set as above with a transfer line temperature, 260°C. The injection volume was 1.0µl of dog faeces and larvae and was injected in a capillary column (Zebran ZB-S, 30 m length x 0.25 mm diameter x 0.25µm phase thickness) (see appendix 4). The oven temperature was programmed from 60°C and increased at the rate of 10°C/min to a final temperature of 260°C, which was held at 260°C for 10 min. High purity helium was used as carrier gas at 1ml/min, split ratio and 1:25; Electron ionization (EI) spectra were obtained at 70 eV; the scan range was 50-450 m/z at 30 min. The identity of each compound was determined by comparison of its retention index (RI) as well as of its total ion chromatogram with the NIST mass spectral library version 5.4.2 (Albano *et al.*, 2011; Sivasamy *et al.*, 2011).

7.3. Results and Discussion

7.3.1. Extraction of biodiesel from larvae

A visual representation of the fact that lipids could be extracted from the faeces on which larvae were grown and from the larvae themselves is shown in Fig.7.1. Table 7.1 shows the percentage of lipids produced in the faeces fed on the larvae and in the larvae themselves




Fig. 7. 1: Fatty acids extracted from; a, BSFL, b) FBL, c) dog faeces with BSFL, d) Dog faeces with FBL

Table. 7. 1: The percentage of lipids in dry weight of 1g samples of faeces and larvae

| Time/ week | Dry weight/ g SD± | Total lipids/ g SD± | % of lipids in dry weight |
|--|------------------------------|--------------------------------|--------------------------------------|
| Zero time | | | |
| Untreated dog faeces (without larvae) | 0.21 ± 0.01 | 0.15 ± 0.02 | 71% |
| BSFL before fed on dog faeces | 0.26 ± 0.01 | 0.22 ± 0.02 | 85% |
| FBL before fed on dog faeces | 0.20 ± 0.01 | 0.10 ± 0.02 | 50% |
| Week 1 | | | |
| Dog faeces with BSFL | 0.70 ± 0.05 | 0.10 ± 0.00 | 14% |
| Dog faeces with FBL | 0.44 ± 0.04 | 0.12 ± 0.01 | 27% |
| BSFL fed on dog faeces | 0.28 ± 0.01 | 0.23 ± 0.01 | 82% |
| FBL fed on dog faeces | 0.25 ± 0.02 | 0.14 ± 0.03 | 56% |
| Week 2 | | | |
| Dog faeces with BSFL | 0.90 ± 0.02 | 0.10 ± 0.02 | 11% |
| Dog faeces with FBL | 0.80 ± 0.02 | 0.14 ± 0.01 | 18% |
| BSFL fed on dog faeces | 0.50 ± 0.02 | 0.15 ± 0.02 | 30% |
| FBL fed on dog faeces | 0.60± 0.01 | 0.14 ± 0.04 | 23% |
| Week 3 | | | |
| Dog faeces with BSFL | 0.90± 0.03 | 0.23± 0.03 | 26% |
| Dog faeces with FBL | 0.80± 0.02 | 0.22± 0.02 | 28% |
| BSFL fed on dog faeces | 0.50± 0.02 | 0.16± 0.04 | 32% |
| FBL fed on dog faeces | 0.70± 0.02 | 0.13± 0.01 | 19% |
| Week 4 | | | |
| Dog faeces with BSFL | 0.80 ± 0.01 | 0.23± 0.01 | 29% |
| Dog faeces with FBL | 0.84± 0.02 | 0.20 ± 0.02 | 24% |
| BSFL fed on dog faeces | 0.62± 0.04 | 0.16 ± 0.02 | 26% |
| FBL fed on dog faeces | 0.82± 0.02 | 0.14 ± 0.01 | 17% |

Untreated dog faeces (without larvae) possessed a lipid content of 71% Table 7.1). The unfed larvae, BSFL and FBL, contained 85 and 50% dry weight of lipids. This suggests that both types of dog faeces and the two larvae could be used independently as a source of lipids for the production of biofuels. The next step was to determine if feeding the larvae on dog faeces would increase, over time, their lipid content and the lipid content of the faeces on which they were feeding.

The results of the week 1 analysis shows that the lipid content of the dog faeces treated with BSFL and FBL fell from 71 to 14 and 27% respectively, while the lipid content of the two larvae remained broadly the same as before feeding on faeces. By week 2, the lipid content of the dog faeces treated with BSFL and FBL declined, as did the concentration of lipids in both types of larvae (Table 7.1); this trend was also seen at weeks 3 and 4 (Table 7.1).












These results show that the decline in dog faeces lipid content was not mirrored by an increase in the lipid content of the two larvae. Presumably, the faeces-lipids were, over time, metabolized by the larvae, ultimately to carbon dioxide, i.e. the larvae were using the faecal lipids as a food source. The results show that no advantage can be gained in relation to total lipid production by treating dog faeces with BSFL and FBL. However, Fig.7.2 shows that the type and concentration of fatty acids (as seen by referring to peak retention times and peak height respectively of the exhibited chromatograms) varied markedly depended on time of exposure of the larvae to dog faeces and the type of larvae used. This is shown more clearly in Table 7.3. Dog faeces without larvae and unfed FBL contained only palmitic and stearic acids, while unfed BSFL contained these fatty acid together with lauric acid.

The type of fatty acid produced in the larvae and in the dog faeces on which the larvae fed was then seen to vary over time. Oleic acid, for example was only produced in the BSFL fed on dog faeces and then only at week 2 (Table 7.2). Clearly, the type of

fatty acid varied with the larvae used and in the faeces treated with the different larvae, all of which varied over time. This finding suggests that BSFL and FBL might be used to produce individual fatty acids, either within their bodies, when fed on dog faeces, or in the dog faeces itself. In this way a single, or desired mixture of fatty acids, might be produced. Thus, if oleic acid alone was needed as a biofuel feedstock, or for some other biotechnological purpose then feeding BSF on dog faeces for 2 weeks and the harvesting the larvae and extracting the fatty acid may provide a source of this individual fatty acid.

Table. 7. 2: The GC/MS analysis chain composition of a number of fatty acids from dog faeces and larval fats

| Fatty acid | Lipids Number | Time | | | | | | | | | | | | | | | |
|------------------|---------------|----------------------|--------------|-------------|-----------------|----------------|--------------------|-------------------|------------------|----------------|--------------------|-------------------|------------------|----------------|--------------------|-------------------|---|
| | | Week 0 | | | Week 1 | | | Week 2 | | | Week 3 | | | Week 4 | | | |
| | | Dog faeces no larvae | Control BSFL | Control FBL | Dog faeces+BSFL | Dog faeces+FBL | BSFL fed on faeces | FBL fed on faeces | Dog faeces +BSFL | Dog faeces+FBL | BSFL fed on faeces | FBL fed on faeces | Dog faeces +BSFL | Dog faeces+FBL | BSFL fed on faeces | FBL fed on faeces | |
| Capric acid | C10:0 | | | | + | | | | | + | + | | | | | | |
| Undecylic acid | C11:0 | | | | | + | | | | | | | | | | | |
| Lauric acid | C12:0 | | + | | | | + | | | | | | | + | | | + |
| Myristic acid | C14:0 | | + | + | | | + | + | | + | | | | + | | | |
| Palmitic acid | C16:0 | + | + | + | | | + | | + | + | | + | + | + | + | + | + |
| Stearic acid | C18:0 | + | | | | | | | | | | + | | | | + | + |
| Palmitoleic acid | C16:1 | | | | | | | | | | | | | | | | |
| Oleic acid | C18:1 | | | | | | | | | + | | | | | | | |






Fig. 7. 2: GC-MS chromatogram of a reference mixture of fatty acid representative the total of samples ion taken from dog faeces and BSF and FB larvae. The chromatogram for this figure and all subsequent figures were obtained with Gas Chromatograph using Zebron ZB-S capillary column

The use of BSF to treat domestic waste has been pioneered by the New York-based Ecosystem Corporation which is asking the US Department of Energy for \$1.75 million to prove the concept. This Company is also seeking a matching grant from the Dog's Biomass Research and Development Initiative to build a \$3.5 million demonstration project that would use 24,000 tons of food waste from an Ohio waste transfer station to feed a "bioreactor" full of flies (Otis et al., 1980). The pilot plant could produce between 150,000 to 195,000 gallons of oil per year; some 26 million tons of food scraps are dumped by Americans into landfills each year, and only about 3 percent of that is being recovered today. Ecosystem claim that with only one-quarter of all those food scraps it could make 100 million gallons of oil every year; add in livestock manure and other agriculture wastes and that amount could increase – particularly if the company can be paid to take it off waste handlers' hands, a possibility which forms an important part of many waste-to-biofuel business plans (Vaughn, 2009)

Restaurant wastes are also an ideal foodstock for BSFL biodiesel production. The major methyl ester components of the biodiesel derived from BSFL were oleinic acid methyl ester (27.1%), lauric acid methyl ester (23.4%), and palmitic acid methyl ester (18.2%) (Table 7.3, 7.4). Table 7.3 also shows that BSFL biodiesel contains a far richer mix of fatty acid methyl ester compounds than does biodiesel obtained from rapeseed. Certainly the BSFL product provides a better source of fatty acids for other possible industrial/food-based products than does rapeseed oil biodiesel.

Table. 7. 3: Comparison of fatty acid methyl ester composition of BSFL fat-based biodiesel and rapeseed oil-based biodiesel.

| Composition | BSFL fat-based biodiesel^a (%) | Rapeseed oil-based biodiesel^b (%) |
|--------------------------------------|---|---|
| Capric acid methyl ester | 3.1 | n/a |
| Lauric acid methyl ester | 35.6 | n/a |
| Myristic acid methyl ester | 7.6 | n/a |
| Palmitoleic acid methyl ester | 3.8 | n/a |
| Palmitic acid methyl ester | 14.8 | 3.5 |
| Oleic acid methyl ester | 23.6 | 64.4 |
| Linoleic acid methyl ester | 5.8 | 22.3 |
| Linolenic acid methyl ester | Nd | 8.2 |
| Stearic acid methyl ester | 3.6 | 0.8 |
| Noadecanic acid methyl ester | 1.4 | n/a |

(See EN 14214 in Refs. Nd stands for not detected).

Most of the properties of this biodiesel met the specifications of the standard EN 14214, including density (860 kg/m^3), viscosity ($4.9 \text{ mm}^2/\text{s}$), flash point ($128 \text{ }^\circ\text{C}$), cetane number (58) and ester contents (96.9%) (Table 7.4).

Table. 7. 4: Comparison of fuel properties of BSFL fat-based biodiesel, rapeseed oil-based biodiesel, and the standard EN14214.

| Properties | EN14214 | BSFL biodiesel^a | Rapeseed biodiesel^b |
|--|----------------|---------------------------------------|---|
| Density (kg/m³) | 860–900 | 885 | 880 |
| Viscosity at 40 °C (mm²/s) | 1.9–6.0 | 5.8 | 6.35 |
| Sulfur content (wt.%) | 0.05 | Nd | <0.01 |
| Ester content (%) | 96.5 | 97.2 | n/a |
| Water content (mg/kg) | <0.03 | 0.03 | 0.03 |
| Flash point (°C) | 120 | 123 | n/a |
| Cetane index | 48–60 | 53 | 45 |
| Acid number (mg KOH/g) | <0.8 | 1.1 | 0.3 |
| Methanol or ethanol (m/m) | 0.2% | 0.3% | n/a |
| Distillation (°C) | n/a | 360 | 352 |

nd stands for not determined

(See EN 14214 in Refs. Nd stands for not detected).

Results such as these show that BSFL obtained from larvae grown on fat wastes could potentially be used as a non-food feedstock for biodiesel production, as well as significantly reducing the large quantity of a solid waste.

It has also been shown in the literature (REF) that larval biomass of BSFL fed on animal manure can efficiently produced biodiesel. Approximately 15.8 g biodiesel was produced from about 1200 black soldier fly larvae when fed on dairy manure over a 21 days period.

Conclusion

The results reported in this Thesis show that dog faeces and BSFL and FBL fed on such faeces can provide a source of fatty acids which could form the basis of biodiesel production. Again however, the main limitation of using dog faeces for any process, be it agricultural or industrial soon becomes apparent and relates to the economics of collecting sufficient quantities of dog waste. When one compares the vast world-wide production of rapeseed oil with the large, but widely dispersed sources of dog faeces it soon becomes obvious that the former has such a huge competitive advantage which makes it impossible that bulk dog faeces-biodiesel could ever compete with other sources of this product.



Chapter Eight: Final Discussion



8.1 Final Discussion

It is perhaps surprising, considering the vast amounts of dog faeces which reach the environment that relatively few scientific studies have been made on the bioremediation of this potentially toxic waste which is major source of disease-causing microorganisms in the urban environment. The obvious distasteful nature of dog faeces doubtless contributes to the lack of interest amongst researchers in this waste product. This lack of interest means that there exists relatively few publications available which are directly relevant to the scientific study of dog faeces.

The broad aim of this work reported in this Thesis was to study a variety of aspects of dog faeces in relation to public health, their fertilizer potential and the possibility of them being remediated using larvae, ultimately to provide a source of biodiesel.

Dog faeces were shown to be a source of pathogenic bacteria, notably *E.coli* and *Salmonella*. These bacteria were shown to be transferred to the soil of a local playing field by direct, *in situ*, transfer from dog faeces undergoing weathering. The potential danger to public health is obvious, especially since children play in parks like these, and are therefore at risk of picking up pathogens from contaminated soil, or more directly from the faeces themselves. Dog faeces are known to transmit bacterial pathogens and, perhaps more worryingly, potentially deadly parasites.

It was shown here that “common or garden” slugs can transfer potentially pathogenic bacteria from dog faeces to lettuce. Again this is potentially damaging to public health, especially since slugs frequent agricultural areas. The following genera and species of bacteria were isolated from slugs: *Enterococcus sp*, *Salmonella sp*, *Staphylococcus lentus*, *E. coli*, *Proteus sp* and *Acinetobacter sp*. *Enterobacter amnigenes* was isolated from the outside secretion of slugs and their identification was confirmed using 16S rRNA sequence gene analysis (i.e.. 99% confirmed). Externally isolated species were identified to *Acinetobacter sp* (99%) using 16SrRNA and *Comamonas sp* was also

confirmed by 16S rRNA analysis is (91%), *Enterobacter amnigenus*, *Acinetobacter sp*, *Comamonas sp* and *Acinetobacter sp* were isolated from lettuce contaminated with slugs fed on dog faeces. *Mycoplasma* was transmitted from dog faeces to slugs or snails which they, ingested or carried, and then in turn transferred to lettuce. Earthworms were also shown to carry *Mycoplasma* from soil on/or within their bodies, a fact which presumably reflects contamination from animal faeces, mainly dogs. Perhaps the greatest potential risk relates to gardens and allotments where slugs may be common, and not successfully controlled. The consumption of pathogen contaminated lettuces, or other salad vegetables, is an obvious potential problem, which would presumably only be solved by vigorous washing of locally grown produce. There is no doubt more research effort has been devoted to the potential risk to human and animal of parasites in dog faeces, rather than bacterial and virus pathogens. Viruses have been isolated from dog faeces (Carmichael and Binn, 1981)

Dog faeces were shown to have potential inherent fertilizer content, the nutrients present being released over a time period mimicking the natural weathering of dog faeces in the environment. The fertilizer potential of dog faeces is not however, great and their offensive nature and potentially rich pathogen content means that un-weathered, or non-composted canine faeces are rarely used as an agricultural fertilizer. Dog faeces were also shown to contain an indigenous microflora capable of mineralizing organic nitrogen to ammonium and oxidizing added ammonium to nitrate; elemental sulphur to sulphate and were able to solubilise insoluble phosphate to plant available, soluble phosphate. As a generalization, the addition of both types of larvae to dog faeces significantly reduced the concentration of indigenous plant nutrients over the entire four week incubation period; exceptions to this were nitrate and phosphate concentrations in BSFL treated faeces, where significant increases were seen at week 4 and 3 respectively and in faeces treated with FBL, where ammonium concentrations

were significantly increased at weeks 2-4, and phosphate at week 4. While the addition of both larvae therefore initially decreased levels of indigenous plant nutrients there was a trend in some of the nutrients to increase the longer the incubation went on. This suggests that perhaps a longer term exposure of dog faeces to the two larvae might have led to increase in ammonium, nitrate, sulphate and phosphate concentrations. While this would generally be advantageous to the fertility of soils surrounding dog faeces, the potential long term increase in nitrate could at first sight be considered detrimental as nitrate can be readily leached to ground waters (where, when present in drinking water, it can cause blue baby disease and gastric cancer in humans). However, the relatively small amounts of dog faeces which are present, spread over large areas of soil, would likely make such increased nitrate contributions to drinking water relatively insignificant.

The addition of ammonium, elemental sulphur and insoluble phosphate to dog faeces which had been modified by the two larvae led to significant increases in nitrate, sulphate and plant-available phosphate, results which shows that that dog faeces contains the indigenous microflora required for the transformation of these amendments (which simulate fertilizer addition). The increased friability and therefore increased aeration of the dog faeces following larval modification is also like to result in enhanced rates of nitrification, S-oxidation and phosphate solubilisation. Clearly, the microflora of dog faeces could act to modify added fertilizers and convert them to the plant available form. Dog faeces represent a potential source for spreading pathogenic bacteria such as *E. coli* and *Salmonella enterica*, bacteria which represent an obvious risk human health. Bacterial pathogens from dog faeces could infect young children and others playing, or using, parks etc (including of course dog owners). Dog faeces have also been shown here to produce gasses which are injurious to plant germination and/ or growth, and are also likely to be directly inhibitory to plant growth.

The potential for using fly larvae for the bioremediation of dog faeces was investigated. Black Soldier Fly (BSFL) and Fruit Beetle (FBL) Fly larvae were shown to dramatically improve the physical nature of canine faeces, even after only a short exposure period, giving a bioremediated product which is markedly improved in terms of texture, reduced odour and overall reduced offensiveness.

The feeding of BSFL on faeces led to a statistically significant increase in the number of bacteria inside the BSFL gut and the same trend was seen in relation to dog faeces fed FBL. This trend of increasing bacterial numbers in larvae fed on dog faeces is particularly worrying in relation to the potential feeding of these larvae to animals- post exposure to faeces. Black soldier Fly larvae in particular are currently being produced for use as animal feed (mainly to chickens) following feeding on restaurant food waste. It is unlikely then, considering the high pathogen content of larvae fed on dog faeces that these could be safely used as animal feed following feeding on dog faeces. The bioremediated dog faeces produced was also found to be suitable as potting compost when “diluted” with proprietary potting compost.

The haemolymph and total body extracts of BSFL and FBL were shown to be antibacterial.

The potential for using dog faeces and dog faeces which had been treated with BSFL and FB was determined. It was shown that potential biodiesel precursors (mainly fatty acids) were present both in the raw dog faeces and in faeces which were treated with the two different larvae. Whether or not dog faeces, treated or otherwise, could be used economically as a source of biodiesel remains to be seen. Although vast amounts of dog faeces are produced annually all over the world problems relating to such collection are obvious. The increasing social trend towards dog owners collecting dog faeces and placing them in council collection containers means that an increasing amount of this potential bioresource is being collected together, at least in the urban environment. The

quantities collected will never however, approach that of other animal faeces, such as cattle feedlot waste. As a result, it might be better to use the relatively small amounts of dog faeces which are collected in any one location for localised methane generation, rather than biodiesel production.

The number of bacteria, including pathogens declined inside the gut of BSFL following feeding on dog faeces. The feeding of BSFL on faeces led to a statistically significant increase in the number of bacteria inside the BSFL gut and the same trend was seen in relation to dog faeces fed FBL. This trend of increasing bacterial numbers in larvae fed on dog faeces is particularly worrying in relation to the potential feeding of these larvae to animals- post exposure to faeces. Black soldier Fly larvae in particular are currently being produced for use as animal feed (mainly to chickens) following feeding on restaurant food waste. It is unlikely then, considering the high pathogen content of larvae fed on dog faeces that these could be safely used as animal feed following feeding on dog faeces. The bioremediated dog faeces produced was also found to be suitable as potting compost when “diluted” with proprietary potting compost. Increasing the ratio of compost to modified faeces “diluted out” the inhibitory effect of raw dog faeces on plant growth suggesting the possibility that larval modified dog faeces could be used as compost additive fertilizer, or perhaps even be used as an agricultural soil fertilizer.

Escherichia coli and *Salmonella enterica* were isolated from all four sites (Fig. 6.3nb) while no such isolates were obtained from the fifth location which was uncontaminated with dog faeces. The number of *E. coli* varied from site to site (from 10^4 to 10^5 CFU/g); *E. coli* was isolated in highest numbers. The isolation of these pathogenic bacteria from playing fields frequented by children is obviously a major public health concern since pathogenic bacteria could be picked up from these soils, either on shoes, or by transfer from contaminated skin to the ears, eyes or mouths of

infected children (and of course, people in general) from where they could cause disease. The number of bacteria present in dog faeces disposed of in plastic bags dramatically increased over exposure to the UK summer, when temperatures were recorded between 10-27°C. The following bacteria were isolated from the enclosed faeces (Fig. 6.4) a), *Enterococcus spp* and *E. coli* in Chromoagar media b), *E. coli* in HiCrome agar media c), *Salmonella enterica* in XLT-4 media while d) was *Staphylococcus aureus* with MRSA selective media respectively. However, the faecal sample counts were:(range 3.5×10^5 to 4.6×10^6 , 9.7×10^3 to 1.0×10^5 and 4.2×10^3 to 2.9×10^5) for *E. coli*, *Salmonella enterica* and plate count respectively. This study has demonstrated that the numbers and the diversity of bacteria can increase when dog faeces are enclosed over a month in plastic bags.

There has been a welcome trend of late for dog owners to act responsibly by picking up waste from their dogs, placing this in plastic bags and than putting these bags into special local council run dog-waste containers, from where they are transported and disposed off.

8.2 Future work

The lack of published work on all aspects of the environmental and agricultural impact of dog faeces means that there is considerable potential to continue and extend the work reported in this Thesis. This Thesis reports preliminary work on all of the individual research questions investigated and all of the separate chapters could be extended to form the basis of separate theses.

1) The work described here on biofuels could be extended, particularly in relation to the type of lipids produced by fly larvae growing on faeces. However, since there exist more readily available and technically more readily useable biofuel substrates available

it would seem economically pointless to devote too much attention to this area of green technology.

2) No work was done here on viruses, both in relation to their presence in faeces and transmission to salads and vegetable. This is clearly an opportunity for workers with the experience and methodology in relation to working with viruses.

3) Considerable interest is currently being devoted to the use of BSFL, produced from waste vegetables and other waste organic matter, as a source of poultry feed. It would be of interest to develop this line of research. Problems related to pathogen transfer mean that it is unlikely however, that such animal feeds could be safely produced from dog faeces.



References



- Albano, S. M., Lima, A. S., Miguel, M. G., Pedro, L. G., Barroso, J. G. And Figueiredo, A. C.** (2011). Antioxidant, Anti-5-lipoxygenase and anti-acetylcholinesterase activities of essential oils and decoction waters of some aromatic plants. *Record of Natural Products* **6**, 35-48.
- Alexander, M.** (1977). *Introduction to Soli Microbiology*. (2nd ed). Wiley, New York.
- Alexander, S. K. and Strete, D.** (2001). *Microbiology: a photographic atlas for the laboratory*. San Francisco: Benjamin Cummings, Addison Wesley Longman, Inc
- Alloway, B.** (2004). Contamination of soils in domestic gardens and allotments: a brief overview. *Land Contamination and Reclamation* **12**, 179-187.
- Alvarez, L.** (2012). The Role of Black Soldier Fly, *Hermetia illucens* (L.)(Diptera: Stratiomyidae) in Sustainable Waste Management in Northern Climates.
- Amann, R., F. Gloëckner and Neef, A.** (1997). Modern methods in subsurface microbiology: in situ identification of microorganisms with nucleic acid probes. *FEMS Microbiology Reviews* **20**, 191-200.
- Amatya, P.** (2009). *Economics of Black Soldier Fly (Hermetia Illucens) in Dairy Waste Management*. ProQuest
- American Public Health Association.** (1992). *Standard Methods for the Examination of Dairy Products. 16th Edn. APHA Inc. Washington DC.*
- Anderson, M., Reynolds, J., Rowe, J., Sverlow, K., Packham, A., Barr, B. and Conrad, P.** (1997). Evidence of vertical transmission of *Neospora* sp infection in dairy cattle. *Journal of the American Veterinary Medical Association* **210**, 1169-1172.
- Anderson, M., Palmer, C. W., Thurmond, M., Picanso, J., Blanchard, P., Breitmeyer, R., Layton, A., McAllister, M., Daft, B. and Kinde, H.** (1995). Evaluation of abortions in cattle attributable to neosporosis in selected dairy

- herds in California. *Journal of the American Veterinary Medical Association* **207**, 1206-1210
- Anderson, J. M. and Baird-Parker, A.** (1975). A rapid and direct plate method for enumerating *Escherichia coli* biotype I in food. *Journal of Applied Microbiology* **39**, 111-117.
- Ann, C., Walter, W., Mulbry, W.** (2002). Recovery of dairy manure nutrients by benthic freshwater algae. *Bioresource Technology* **84**, 81–91.
- Atlas, R.M. and Bartha, R.** (1998). *Microbial Ecology-Fundamentals and Applications*, Benjamin Cummings, Menlo Park.
- Barutzki, D. and Schaper, R.** (2003). Endoparasites in dogs and cats in Germany 1999–2002. *Parasitology Research*. **90**, 148-150.
- Baxter, D. and Leck, I.** (1984). The deleterious effects of dogs on human health: 2. Canine zoonoses. *Journal of Public Health* **6**, 185-197.
- Beuchat, L. R.** (1996). Pathogenic microorganisms associated with fresh produce. *Journal of Food Protection* **59**, 204-216.
- Beuchat, L. R.** (2006). Vectors and conditions for preharvest contamination of fruits and vegetables with pathogens capable of causing enteric diseases. *British Food Journal* **108**, 38-53.
- Bexfield, A., Nigam, Y., Thomas, S. and Ratcliffe, N. A.** (2004). Detection and partial characterisation of two antibacterial factors from the excretions/secretions of the medicinal maggot *Lucillia sericata* and their activity against methicillin-resistant *Staphylococcus aureus* (MRSA). *Microbes and Infection* **6**, 1297-1304.
- Bhattacharyya, P., Chakraborty, A., Bhattacharya, B., Chakrabarti, K.** (2003b). Evaluation of MSW compost as a component of integrated nutrient management in wetland rice. *Compost Science and Utilization*. **11**, 343–350.
- Biederbeck, V.O.** (1979). In *Soil Organic Matter*. Elsevier, New York.

- Blancou, J., Chomel, B. B., Belotto, A. and Meslin, F. X.** (2005). Emerging or re-emerging bacterial zoonoses: factors of emergence, surveillance and control. *Veterinary Research* **36**, 507-522.
- Bogdanov, S., Martin, P. and Lullmann, C.** (2002). Harmonised methods of the international honey commission. *Swiss Bee Research Centre, FAM, Liebefeld*.
- Brady, C. and Weil, R.** (1974). *The Nature and Properties of Soil*. Prentice Hall, London.
- Brinton, W.F and Storms, P.H.** (2004). Microbiological test qualities of composted manures and yard wastes. World Congress Organic Food, Michigan State University, March 29-31.
- Brown, K.A.** (1982). Sulphur in the environment a review. *Environmental Pollution* **3**, 47-80.
- Burns, G.R.** (1967). *Oxidation of Sulphur in Soils*. Sulphur Institute, Washington D.C. *Tech. Bull* 13, 1-41
- Butler, T.A., Sikora, L.J., Steinhilber, P.M. and Douglass, L.W.** (2001). Compost age and sample storage effects on maturity indicators of biosolids compost. *Journal of Environmental Quality* **30**, 2141–2148.
- Carmicheal, L.E. and Binn, L.N.** (1981). New enteric virus in the dog. *Advances in Veterinary Science and Comparative Medicine* **25**, 1-27
- Carvalho, R., Araújo, J., Braga, F., Araujo, J., Silva, A. and Tavela, A.** (2009). Predatory activity of nematophagous fungi on infective larvae of *Ancylostoma* sp.: evaluation in vitro and after passing through the gastrointestinal tract of dogs. *Journal of Helminthology* **83**, 231-236.
- Cekmecelioglu, D., Demirci, A., Graves, R. and Davitt, N.** (2005). Optimization of windrow food waste composting to inactivate pathogenic microorganisms. *Transactions of the ASAE* **48**, 2023.

- Chiu, S. Y., Kao, C. Y., Tsai, M. T., Ong, S. C., Chen, C. H. and Lin, C. S.** (2009). Lipid accumulation and CO₂ utilization of *Nannochloropsis oculata* in response to CO₂ aeration. *Bioresource Technology* **100**, 833-838.
- Chynoweth, D. P., Owens, J. M. and Legrand, R.** (2001). Renewable methane from anaerobic digestion of biomass. *Renewable Energy* **22**, 1-8.
- Clinical and Laboratory Standards Institute.** (2008). Performance standards for antimicrobial susceptibility testing; Eighteenth Informational Supplement, M100-S18. CLSI, Wayne, PA.
- Cociancich, S., Bulet, P., Hetru, C. and Hoffmann, J.** (1994). The inducible antibacterial peptides of insects. *Parasitology Today* **10**, 132-139.
- Colaizy, T. T., Kuforiji, T., Sklar, R. S. and Pillers, D. A. M.** (2003). PCR methods in clinical investigations of human ureaplasmas: a minireview. *Molecular Genetics and Metabolism* **80**, 389-397.
- Cooper, P.J.M.** (1972). Sequence of products formed during denitrification in some diverse western soils. *Soil science Society of America Proceedings* **27**, 659-662
- Craig Sheppard, D., Larry Newton, G., Thompson, S. A. and Savage, S.** (1994). A value added manure management system using the black Soldier Fly. *Bioresource Technology* **50**, 275-279.
- Craig Sheppard, D., Tomberlin, J. K., Joyce, J. A., Kiser, B. C. and Sumner, S. M.** (2002). Rearing methods for the black soldier fly (Diptera: Stratiomyidae). *Journal of Medical Entomology* **39**, 695-698.
- Crowley, L. D. and Houck, M. A.** (2002). The immune response of larvae and pupae of *Calliphora vicina* (Diptera: Calliphoridae), upon administered insult with *Escherichia coli*. *Journal of Medical Entomology* **39**, 931-934.
- Dawkins, G. J., M. L. Hislop and C. Bishop.** (1986). Transmission of bacteria soft rot of potatoes by slug. *Molluscan Studies*. **52**, 25-29.

- Davison, H., Otter, A. and Trees, A.** (1999). Estimation of vertical and horizontal transmission parameters of *Neospora caninum* infections in dairy cattle. *International Journal for Parasitology* **29**, 1683-1689.
- Demirbas, A.** (2008). *Biodiesel: a realistic fuel alternative for diesel engines*. Springer.
- Demirbas, A.** (2008). Biofuels sources, biofuel policy, biofuel economy and global biofuel projections. *Energy Conversion and Management* **49**, 2106-2116.
- Demirel, B., Scherer, P., Yenigun, O. and Onay, T. T.** (2010). Production of methane and hydrogen from biomass through conventional and high-rate anaerobic digestion processes. *Critical Reviews in Environmental Science and Technology* **40**, 116-146.
- De Montfort, G.** (2012). Waste Derived Biofuels: Are such fuels a suitable alternative to fossil fuels?.
- Despommier, D.** (2003). Toxocariasis: clinical aspects, epidemiology, medical ecology, and molecular aspects. *Clinical Microbiology Reviews* **16**, 265-272.
- Dox AW** (1910). The intracellular enzymes of *Penicillium* and *Aspergillus* with special reference to those of *P.camemberti*. *United States Department of Agriculture, Bureau of Animal Industry Bulletin*. No. 1; 120-170.
- Drasar, B.S. and P. A. Barrow.** (1985). Intestinal microbiology. In: Aspects of Microbiology (Schlessinger, D., Ed.), Washington, D. *American Society of Microbiology* **10**, 28– 38.
- Drasar, D.S. and M. J. Hill** (1974). *Human Intestinal Flora*. Academic Press, London, UK.
- Dubey, J and Williams, C.** (1980). Hammondia heydorni infection in sheep, goats, moose, dogs and coyotes. *Parasitology* **81**, 123-127.
- Dubey, J. P. and Lindsay, D.S.** (2006). Neosporosis, toxoplasmosis, and sarcocystosis in ruminants. *Veterinary Clinics Food and Animal Practise* **22**, 645-671.

- Dubey, J. P.** (2003). Review of *Neospora caninum* and neosporosis in animals. *The Korean journal of parasitology* 41, 1-16.
- Dusch, H. and Altwegg, M.** (1995). Evaluation of five new plating media for isolation of *Salmonella* species. *Journal of Clinical Microbiology* 33, 802-804.
- Elder, R. O., Keen, J. E., Siragusa, G.R., Barkocy-Gallagher, G. A., Koochmaraie, M., and Laegreid, W. W.** (2000). Correlation of enter hemorrhagic *Escherichia coli* O157 prevalence in feces, hides, and carcasses of beef cattle during processing. *Proceedings of the National Academy of Sciences of the United States of America* 97, 2999–3003.
- Elliot, L. P.** (1969). Certain bacteria, some of medical interest, associated with the slug *Limax maximus*. *Journal of Invertebrate Pathology* 15, 306–312.
- Emma, L. S., M. Macrae., I. D. Ogden., M. J. Wilson and N. J. C. Strachan.** (2006). Slugs: Potential Novel Vectors of *Escherichia coli* O157. *Applied and Environmental Microbiology* 72, 144–149.
- EN 14214** Automotive fuels – Fatty acid methyl esters (FAME) for diesel engines – Requirements and test methods. European Committee for Standardization (CEN). Brussels, 2003.
- Erickson, M. C., Islam, M., Sheppard, C., Liao, J. and Doyle, M. P.** (2004). Reduction of *Escherichia coli* O157: H7 and *Salmonella enterica* serovar Enteritidis in chicken manure by larvae of the Black Soldier Fly. *Journal of Food Protection* 67, 685-690.
- Falih, A. M.** (1995). *The Microbiology of Soil Receiving High Concentration of Sucrose from a Natural Source*. PhD Thesis, University of Sheffield
- Fangmeier, A., Hadwiger-Fangmeier, A., Van der Eerden, L. and Jager, H. J.** (1994). Effects of atmospheric ammonia on vegetation--A review. *Environmental pollution* 86, 43-82.

- Ferrer, E., Aho, M., Silvennoinen, J. and Nurminen, R.-V.** (2005). Fluidized bed combustion of refuse-derived fuel in presence of protective coal ash. *Fuel processing technology* 87, 33-44.
- Fredriksson-Ahomaa, M., Korte, T. and Korkeala, H.** (2001). Transmission of *Yersinia enterocolitica* 4/O:3 to pets via contaminated pork. *Letters in Applied Microbiology* 32, 375–378.
- Frenkel, J.K.** (1990). Transmission of toxoplasmosis and the role of immunity in limiting transmission and illness. *Journal of the American Veterinary Medical Association*. 196, 233-240.
- Frenkel, J.K., Parker, B.B.** (1996). An apparent role of dogs in the transmission of *Toxoplasma gondii*: the probable importance of xenosmophilia. *Annals of the New York Academy of Sciences*. 791, 402–407.
- Garber, K.** (1935). Über die Physiologie der Einwirkung von Ammoniakgasen auf die Pflanze. *Landw. Vers.-Stat* 123, 277-344.
- Garner, J.S.** (1996). Hospital Infection Control Practices Advisory Committee, U.S. Department of Health and Human Services, Centres for Disease Control and Prevention. Guideline for isolation precautions in hospitals. *Infection Control and Hospital Epidemiology* 17: 53-80
- Gillespie, S.** (1988). The epidemiology of *Toxocara canis*. *Parasitology Today* 4, 180-182.
- Godan, D.** (1983). *Pest Slugs and Snails: biology and control*. Springer-Verlag, Berlin, Germany. [translated by S. Gruber].

- Guittoneau, G.** (1927). Sur l'oxidation microbienne du soufre au course de l'ammonisation. *Comptes Rendus Hebdomadaires des Sciences de l'Academie des Sciences* **184**, 45-46.
- Hall, RE and Schulte, E.F.** (1979). *Using Dog and Cat Manure on Home Gardens*. University of Wisconsin, Madison, College of Agriculture and Life Sciences.
- Hansen, W. and Yourassowsky, E.** (1984). Detection of beta-glucuronidase in lactose-fermenting members of the family Enterobacteriaceae and its presence in bacterial urine cultures. *Journal of Clinical Microbiology* **20**, 1177-1179
- Hesse, P.R.** (1971). *A Textbook of Soil Chemical Analysis*, J. Murray, London.
- Hill, I.R. Denham, D.A. and Scheltz, C.L.** (1985). *Toxocara canis* larvae in the brain of a child. *Royal Society. Tropical Medicine and Hygiene*. **79**, 351-354.
- Himathongkham, S., and H. Riemann.** (1999). Destruction of *Salmonella typhimurium*, *Escherichia coli* O157:H7 and *Listeria monocytogenes* in chicken manure by drying and/or gassing with ammonia. *FEMS Microbiology Letters* **171**, 179-182.
- Hogan, J. M.** (1985). *The Behaviour of the Grey Field Slug (Deroceras reticulatum) with particular reference to control in winter wheat*. Ph.D. Thesis. Newcastle University, Newcastle upon-Tyne, UK.
- Hoque S, Heath SB and Killham K.** (1987). Evaluation of methods to assess adequacy of potential soil sulphur supply to crops. *Plant and Soil* **101**, 3-8
- Huberman, L., Gollop, N., Mumcuoglu, K., Block, C. and Galun, R.** (2007). Antibacterial properties of whole body extracts and haemolymph of *Lucilia sericata* maggots. *Journal of Wound Care* **16**, 123-127.
- Institut de la statistique Quebec** (2005a) : Estimation trimestrielle de la population au 1er avril 2005 et part par province. http://www.stat.gouv.qc.ca/donstat/econm_finncl_conjn_econm/TSCI [cited Aug. **30**, 2005].

- Islam, M., Doyle, M. P., Phatak, S. C., Millner, P. and Jiang, X.** (2005). Survival of *Escherichia coli* O157: H7 in soil and on carrots and onions grown in fields treated with contaminated manure composts or irrigation water. *Food Microbiology* **22**, 63-70.
- Jacobs-Reitsma, W. F., A. W. van de Giessen., N. M. Bolder, and R. W. A. W.b Mulder** (1995). Epidemiology of *Campylobacter* Spp. at two Dutch broiler farms. *Epidemiology and Infection*. **114**, 413-421.
- Jacquelyn, G.** (1999). *Microbiology-Principles and Explorations*. London, Wiley.
- Jaklič, D., Lapanje, A., Zupančič, K., Smrke, D. and Gunde-Cimerman, N.** (2008). Selective antimicrobial activity of maggots against pathogenic bacteria. *Journal of Medical Microbiology*. **57**, 617-625.
- Jeon, H., Park, S., Choi, J., Jeong, G., Lee, S. B., Choi, Y. and Lee, S. J.** (2011). The Intestinal Bacterial Community in the Food Waste-Reducing Larvae of *Hermetia illucens*. *Current Microbiology*. **62**, 1390-1399.
- Jiang, X., Morgan, J. and Doyle, M. P.** (2003). Fate of *Escherichia coli* O157: H7 during composting of bovine manure in a laboratory-scale bioreactor. *Journal of Food Protection* **174**, 25-30.
- Karadam, S. Y., Ertug, S., Ertabaklar, H. and Okyay, P.** (2008). The comparison of IgG antibodies specific to *Toxocara* Spp. among eosinophilic and non-eosinophilic groups. *Microbiologica-Quarterly Journal of Microbiological Sciences* **31**, 113-116.
- Kashmanian, R. M. and Rynk, R. F.** (1996). Agricultural composting in the United States: Trends and driving forces. *Journal of Soil and Water Conservation* **51**, 194.- 201

- Kebreab, E., Clark, K., Wagner-Riddle, C. and France, J.** (2006). Methane and nitrous oxide emissions from Canadian animal agriculture: A review. *Canadian Journal of Animal Science* **86**, 135-157.
- Killham, K.** (1994). *Soil Ecology*, Cambridge, Cambridge University Press.
- Kim R.E.** (2008). Dog Meat in Korea. A Socio- Legal Challenge” *Animal Law* **14**, 201-236.
- Kostov, O., Tzvetkov, Y., Kaloianova, N., Van Cleemput, O.** (1996). Production of tomato seedlings on composts of vine branches and grape prunings, hunks and seeds. *Compost Science and Utilization* **4**, 55–61.
- Kudva, I. T., Blanch, K. and Hovde, C. J.** (1998). Analysis of *Escherichia coli* O157: H7 survival in ovine or bovine manure and manure slurry. *Applied and Environmental Microbiology* **64**, 3166-3174.
- Kuenen, J.G. and Beudeker, R.F.** (1982). Soil fungistasis, relation to fungal spore nutrition. *Phytopathology* **298**, 473-497.
- LaGoy, P. K.** (1987). Estimated soil ingestion rates for use in risk assessment. *Risk Analysis* **7**, 355-359.
- Lam, H.-M., Coschigano, K., Oliveira, I., Melo-Oliveira, R. & Coruzzi, G.** (1996). The molecular-genetics of nitrogen assimilation into amino acids in higher plants. *Annual review of plant biology* **47**, 569-593.
- Lapage, S. P., Shelton J. E. and Mitchell T. G.** (1970) in *Methods in Microbiology*’ Eds. Norris J. R. and Ribbons D. W. Vol.3A. Academic Press. London. p. 116.
- Lee, B., Ellenbecker, M., Moure-Eraso, R.** (2004). Alternatives for treatment and disposal cost reduction of regulated medical wastes. *Waste Manage* **24**, 143-151.
- Leem, J. Y., Jeong, I. J., Park, K. T. and Park, H. Y.** (1999). Isolation of p-hydroxycinnamaldehyde as an antibacterial substance from the saw fly, *Acantholyda parki* . *FEBS letters* **442**, 53-56.

- Liao, W., Liu, Y., Wen, Z.Y., Frear, C. and Chen, S.L.** (2008). Kinetic modelling of enzymatic hydrolysis of cellulose in differently pre-treated fibers from dairy manure. *Biotechnology and Bioengineering* **101**, 441–451.
- Lincoln, C. K. and Gabridge, M. G.** (1998). Cell culture contamination: sources, consequences, prevention, and elimination. *Methods in Cell Biology* **57**, 49-65.
- Li, Q., Zheng, L., Cai, H., Garza, E., Yu, Z. and Zhou, S.** (2011a). From organic waste to biodiesel: Black soldier fly, *Hermetia illucens*, makes it feasible. *Fuel*. **90**, 1545-1548.
- Li, Q., Zheng, L., Hou, Y., Yang, S. and Yu, Z.** (2011b). Insect fat, a promising resource for biodiesel. *Journal of Petroleum and Environmental Biotechnology* **S 2**, 2-6.
- Li, Q., Zheng, L., Qiu, N., Cai, H., Tomberlin, J. K. and Yu, Z.** (2011c). Bioconversion of dairy manure by black soldier fly (Diptera: Stratiomyidae) for biodiesel and sugar production. *Waste Management* **31**, 1316-1320.
- Lindsay D.S., Dubai, J.P., Butler J.M., Blagburn, B.L.** (1997). Mechanical transmission of *Toxoplasma gondii* oocysts by dogs. *Veterinary Parasitology* **73**, 27–33.
- London, J. and Rittenberg, S.C.** (1967). *Thiobacillus permetabolis* nov. sp, a
- Lord, W. D., Lee Goff, M., Adkins, T. R. and Haskell, N.** (1994). The black soldier fly *Hermetia illucens* (Diptera: Stratiomyidae) as a potential measure of human postmortem interval: observations and case histories. *Journal of Forensic Sciences* **39**, 215-215.
- Lu, Y., Ludsin, S. A., Fanslow, D. L. and Pothoven, S. A.** (2008). Comparison of three microquantity techniques for measuring total lipids in fish. *Canadian Journal of Fisheries and Aquatic Sciences* **65**, 2233-2241.
- Lynch, J.M.** (1983). *Soil Biotechnology, Microbiological Factors in Crop Productivity*. Blackwell. Boston.

- Ma, F. and Hanna, M. A.** (1999). Biodiesel production: a review. *Bioresource technology* **70**, 1-15.
- Maier, R. M., Pepper, I. L. and Gerba, C. P.** (2009). *Environmental Microbiology*. London, Academic Press.
- Makarov, V. M. and Drozdovskii, V.** (1991). Reprocessing of tyres and rubber wastes: recycling from the rubber products industry. Ellis Horwood.
- Mallinson, E., Miller, R., de Rezende, C., Ferris, K., deGraft-Hanson, J. and Joseph, S.** (2000). Improved plating media for the detection of *Salmonella* species with typical and atypical hydrogen sulfide production. *Journal of Veterinary Diagnostic Investigation* **12**, 83.-87.
- Marketing, L.** (2002). *Canadians and Their Pets*. Montreal, Quebec, Canada: Leger Marketing Trends Report. Canadian Press/Leger Marketing, pgs 2-3.
- Mawdsley, J. L., Bardgett, R. D., Merry, R. J., Pain, B. F. and Theodorou, M. K.** (1995). Pathogens in livestock waste, their potential for movement through soil and environmental pollution. *Applied Soil Ecology* **2**, 1-15.
- McGarrity, G., Sarama, J and Vanaman, V.** (1985). Cell culture techniques. *ASM News* **51**, 170-183.
- McGarrity, G. J., Steiner, T and Vanaman, V.** (1983). Detection of mycoplasmal infection of cell cultures by DNA fluorochrome staining. *Methods in Mycoplasmaology* **2**, 155-208.
- McKinley, V., Vestal, J. and Eralp, A.** (1985). Microbial activity in composting. *Biocycle* **26**, 39-43.
- McLaughlin, J.** (2002). *Gardens As a Source of Infectious Disease: Reducing the Risk*. Miami Dade County Extension Service.

- Merlino, J.** (1997). Application of Chromagar Orientation in the identification and differentiation of Proteeae from other Enterobacteriaceae. *Australian Journal of Medical Science* **18**, 20-23.
- Merlino, J., Siarakas, S., Robertson, G. J., Funnell, G. R., Gottlieb, T. and Bradbury, R.** (1996). Evaluation of CHROMagar Orientation for differentiation and presumptive identification of gram-negative bacilli and *Enterococcus* species. *Journal of Clinical Microbiology*. **34**, 1788-1793.
- Miller, R. G. and Tate, C. R.** (1990). XLT4: A highly selective plating medium for the isolation of *Salmonella*. *Maryland Poultryman*, April p. 2-7.
- Miller, F.C.** (1996). Heat Evolution during Composting of Sewage Sludge. In: *The Science of Composting*, edited by de Bertoldi, M.D., Sequi, P., Lemmes, B., Papi, T. Glasgow, United-Kingdom: Blackie Academic and Professional, pgs 106-115.
- Muir, A., G. Forrest, noskralC ,J and slaehW, A** (2011). Detection of *Candida albicans* DNA from blood samples using a novel electrochemical assay. *Journal of Medical Microbiology* **60**, 467-471
- Mumcuoglu, K. Y., Miller, J., Mumcuoglu, M., Friger, M. and Tarshis, M.** (2001). Destruction of bacteria in the digestive tract of the maggot of *Lucilia sericata* (Diptera: Calliphoridae). *Journal of Medical Entomology* **38**, 161-166.
- Murry, A. C. and Hinckley, L. S.** (1992). Effect of the earthworm (*Eisenia foetida*) on *Salmonella enteritidis* in horse manure. *Bioresource Technology* **41**, 97-100.
- Nemiroff, L. and Patterson, J.** (2007). Design, Testing and Implementation of a Large-Scale Urban Dog Waste Composting Program. *Compost Science and Utilization* **15**, 237-242.
- Newton, L., Sheppard, C., Watson, D., Burtle, G. and Dove, R.** (2005). Using the black Soldier Fly, *Hermetia illucens*, as a value-added tool for the management

of swine manure. Animal and Poultry Waste Management Center, North Carolina State University, Raleigh, NC: 17.

Nor, Y. and Tabatabai, M. A. (1977). Extraction and colorimetric determination of thiosulphate and tetrathionate soils. *Soil Science* **122**, 175-177.

Otis, R. J., Boyle, W. C., Clements, E. V. and Schmidt, C. J. (1980). Design Manual; Onsite Wastewater Treatment and Disposal Systems. *Environmental Protection Agency Report EPA-625/1-80-012, October 1980. 412 p, 86 Fig, 82 Tab, 204 Ref. 1 Append.*

Overgaauw, P. A. M., Van Zutphen, L., Hoek, D., Yaya, F. O., Roelfsema, J., Pinelli, E., Van Knapen, F. and Kortbeek, L. M. (2009). Zoonotic parasites in fecal samples and fur from dogs and cats in The Netherlands. *Veterinary Parasitology* **163**, 115-122.

Paul, E.A and Clark, F.E. (1989). *Soil Microbiology and Biochemistry*. Academic Press, London.

Payne, W.J. (1981). *Dentitrification* . JohnWiley & Sons, Inc., New York, N.Y.

Peigne, J., Girardin, P. (2004). Environmental impacts of farm-scale composting practices. *Water, Air, and Soil Pollution* **153**, 45-68.

Pell, A. N. (1997). Manure and microbes: public and animal health problem? *Journal of Dairy Science* **80**, 2673-2681.

Phan, A. N. and Phan, T. M. (2008). Biodiesel production from waste cooking oils. *Fuel* **87**, 3490-3496.

PHLS *Standard methods for the Microbial Examination of Food and Dairy Samples* (1999) Aerobic plate count at 30°C - F10 and F11.

- Pryor Williams, A., Roberts, P., Avery, L. M., Killham, K. and Jones, D. L.** (2006). Earthworms as vectors of *Escherichia coli* O157: H7 in soil and vermicomposts. *FEMS Microbiology Ecology* **58**, 54-64.
- Richards B.N.** (1987). *The Microbiology of Terrestrial Ecosystem*. London, Longman.
- Rinaldi, L., Biggeri, A., Carbone, S., Musella, V., Catelan, D., Veneziano, V and Cringoli, G.** (2006). Canine faecal contamination and parasitic risk in the city of Naples (southern Italy). *BMC Veterinary Research* **2**, 29.
- Rippy, A, Gleason, M., Ash, A, Willingham, S.** (1997). *Composting Dog Waste in Interior Alaska, Fairbanks Soil and Water District, Study Details*, CWA Section 319 Progress Report, Project : 8-FY92-AK, 14 pgs
- Rottem, S. and Barile, M. F.** (1993). Beware of mycoplasmas. *Trends in Biotechnology* **11**, 143-151.
- Rubel, D. and Wisnivesky, C.** (2005). Magnitude and distribution of canine fecal contamination and helminth eggs in two areas of different urban structure, Greater Buenos Aires, Argentina. *Veterinary Parasitology*. **133**, 339-347.
- Rynk, R (ed.)** (1992). *On-Farm Composting Handbook*. Ithaca, New York: Natural Resource, Agriculture, and Engineering Service (NRAES), pps 12-20.
- Sahalan, A. Z., Omar, B., Mohamed, A. Y. and Jeffery, J.** (2006). Antibacterial activity of extracted Hemolymph from larvae and pupae of local fly species *Musca Domestica* and *Chrysomya Megacephala*. *Jurnal Sains Kesihatan Malaysia* **4**, 1-11.
- Samra, Z., Heifetz, M., Talmor, J., Bain, E. and Bahar, J.** (1998). Evaluation of use of a new chromogenic agar in detection of urinary tract pathogens. *Journal of Clinical Microbiology* **36**, 990-994.
- Santamaría, J. and Toranzos, G. A.** (2003). Enteric pathogens and soil: a short review. *International microbiology* **6**, 5-9.

- Schantz, P.** (1994). Of worms, dogs, and human hosts: Continuing challenges for veterinarians in prevention of human disease. *Journal of the American Veterinary Medical Association*. **204**, 1023-1028.
- Schook, L.B. and Berk, R.S.** (1978). Acid precipitation, Effects on fish. *Ambio* **5**, 228-230
- Sheppard, D. C. and Newton, G. L.** (2001). Insect digestion of manure. In: *Manure Management Strategies/technologies*. White paper prepared under the auspices of the National Center for Manure and Animal Waste Management, NCSU.
- Shulman, V. L.** (2004). *Tyre recycling*. iSmithers Rapra Publishing.
- Silvis, H.** (2006). Driving forces of European agriculture. *Wageningen, The Netherlands. LEI Agricultural Economics Research Institute*.
- Sims, J.R. and Jackson, G.D.** (1971) Rapid analysis of soil nitrate with chromotropic acid. *Soil Science Society American of Proceedings* **35**, 603-606.
- Sivasamy, R., Angayarkanni, J. and Palaniswamy, M.** (2011). A novel filarial topoisomerase II inhibitor produced by native isolate *Micrococcus luteus* B1252. *African Journal of Biotechnology*. **10**, 16069-16077.
- Small, P., Blankenhorn, D., Welty, D., Zinser, E. and Slonczewski, J. L.** (1994). Acid and base resistance in *Escherichia coli* and *Shigella flexneri*: role of rpoS and growth pH. *Journal of Bacteriology* **176**, 1729-1737.
- Simmons, S.** (1935). A bactericidal principle in excretions of surgical maggots which destroys important etiological agents of pyogenic infections. *Journal of Bacteriology* **30**: 253-267.
- Sims, J.R. and Jackson, G.D.** (1971). Rapid analysis of soil nitrite with chromic acid. *Soil Science Society of America Proceedings* **35**, 603-606.

- Smith, D.C., Hughes, J.C.** (2004). Changes in maturity indicators during the degradation of organic wastes subjected to simple composting procedures. *Biology and Fertility of Soils* **39**, 280–286.
- South, A.** (1992). *Terrestrial Slugs: biology, ecology and control*. London, Chapman Hall.
- Sproston, E. L., Macrae, M., Ogden, I. D., Wilson, M. J. and Strachan, N. J. C.** (2006). Slugs: Potential novel vectors of *Escherichia coli* O157. *Applied and Environmental Microbiology* **72**, 144-149.
- Starkey, R.L.** (1934). The production of polythionates from thiosulphate by microorganisms. *Journal of Bacteriology* **48**, 387-400.
- Stoffella, P. J. and Kahn, B. A. (eds.)** (2001). *Compost Utilization in Horticultural Cropping Systems*, Boca Raton, CRC Press.
- Strom, P. F.** (1985). Identification of thermophilic bacteria in solid-waste composting. *Applied and Environmental Microbiology*. **50**, 906-913.
- Tarsitano, E., Greco, G., Decaro, N., Nicassio, F., Lucente, M. S., Buonavoglia, C. and Tempesta, M.** (2010). Environmental monitoring and analysis of faecal contamination in an urban setting in the city of Bari (Apulia region, Italy): health and hygiene implications. *International Journal of Environmental Research and Public Health* **7**, 3972-3986.
- Tan, R.J.S., Lim, E.W. and Ishak, B.** (1977). Ecology of mycoplasmas in clinically healthy cats. *Australian Veterinary Journal* **53**, 515–519.
- Tan, R.J.S., and Miles, J.A.R.** (1974). Incidence and significance of mycoplasmas in sick cats. *Research in Veterinary Science* **16**, 27–34.
- Tate, C., Miller, R., Mallinson, E., Douglass, L. and Johnston, R.** (1990). The isolation of salmonellae from poultry environmental samples by several

enrichment procedures using plating media with and without novobiocin.
Poultry Science. **69**, 721-726.

Taylor, L. (2004). *Poop, Scoop and Compost: Proposal to Explore the Feasibility of Composting Dog Waste in Public Parks*. Vancouver, Canada: National Coalition for People and Dogs, Handout, pgs. 1-7.

Taylor-Robinson, D. (1996). Infections due to species of *Mycoplasma* and *Ureaplasma*: an update. *Clinical Infectious Diseases* **23**, 671-682.

Themelis, N. J. and Ulloa, P. A. (2007). Methane generation in landfills. *Renewable Energy* **32**, 1243-1257.

Thierry, F., G. Frédéric., N. Jacques and A. P. Burnens (2008). The Role of *Acinetobacter baumannii* as a nosocomial pathogen for dogs and cats in an intensive care unit. *Journal of Veterinary Internal Medicine* **2**, 177 – 183.

Tiedje, J.M. (1988). Ecology of denitrification and dissimilation nitrate reduction to ammonium. Pp. 179-244. In: A.J.B.Zehnder (ed.) *Biology of Anaerobic Microorganisms*. John Wiley, New York

Tisdale, S.L., Nelson, W. L., Beaton, J.D. (1985). *Soil Fertility and Fertiliser*, New York, Macmillan.

Thomas, S., Andrews, A., Hay, N. and Bourgoise, S. (1999). The anti-microbial activity of maggot secretions: results of a preliminary study. *Journal of Tissue Viability* **9**, 127.

Thomas, S., Jones, M., Shutler, S. and Jones, S. (1996). Using larvae in modern wound management. *Journal of Wound Care*. **5**, 60.

Thompson, D. E., Bundy, D. A. P., Cooper, E. S. and Schantz, P. M. (1986). Epidemiological characteristics of *Toxocara canis* zoonotic infection of children in a Caribbean community. *Bulletin of the World Health Organization* **64**, 283.

- Tomberlin, J. K., Sheppard, D. C. and Joyce, J. A.** (2002). Selected life-history traits of black soldier flies (Diptera: Stratiomyidae) reared on three artificial diets. *Annals of the Entomological Society of America* **95**, 379-386.
- Trudinger, P.A.** (1967). Metabolism of thiosulphate and tetrathionate by heterotrophic bacteria from soil, *Journal of Bacteriology* **93**, 550-559.
- Traub, R. J., Robertson, I. D., Irwin, P., Mencke, N. and Thompson, R. C. A.** (2002). The role of dogs in transmission of gastrointestinal parasites in a remote tea-growing community in northeastern India. *American Journal of Tropical Medicine and Hygiene* **67**, 539-545.
- Trautmann, N., Olynciw, E.** (2000). *Compost Microorganisms*. Cornell University Composting Page. <<http://www.cfe.cornell.edu/compost/microorg.html>>.
- Turnell, J., Faulkner, R. and Hinch, G.** (2007). Recent advances in Australian broiler litter utilisation. *World's Poultry Science Journal* **63**, 223-232.
- Van der Eerden, L.** (1982). Toxicity of ammonia to plants. *Agriculture and Environment* **7**. 223-235.
- Vaughn, J.** (2009). *Waste Management: A Reference Handbook*. Abc-clio. *pp* 311
- Vitolins, M.I. and Swaby, R.J.** (1969). Activity of sulphur-oxidizing microorganisms in some Australian soils. *Australian Journal of Soil Research* **7**, 171-183
- Wainwright, M.** (1978a). Distribution of sulphur oxidation products in soil and on *Acer pseudoplatanus* L. growing close to source of atmospheric pollution. *Environmental Pollution* **17**, 153-160.
- Wainwright, M.** (1981). Enzyme activity in intertidal sands and salt-marsh soils. *Plant and Soil* **59**, 357-363.
- Wainwright, M and Killham, K.** (1980). Sulphur oxidation by *Fusarium solani*. *Soil Biology and Biochemistry* **12**, 555-558.

- Wainwright, M. and Pugh, G.J.F.** (1973). The effect of three fungicides on nitrification and ammonification in soil. *Soil Biology and Biochemistry* **5**, 577-584.
- Walker, A. J., D. M. Glen and P. R. Shewry** (1999). Bacteria associated with the digestive system of the slug *Deroceras reticulatum* are not required for protein digestion. *Soil Biology and Biochemistry* **31**, 1387–1394.
- Wallace, G.D.** (1973). The role of the cat in the natural history of *Toxoplasma gondii*. *American Journal of Tropical Medicine and Hygiene* **22**, 313–22.
- WHO**, (1959). Joint expert committee on Zoonoses second report. *Tech. Rep. Ser.* No 169.
- Wang, R. F., Cao, W. W. and Cerniglia, C. E.** (1996). PCR detection and quantitation of predominant anaerobic bacteria in human and animal fecal samples. *Applied and Environmental Microbiology* **62**, 1242-1247.
- Weisburg, W. G., Barns, S. M., Pelletier, D. A. & Lane, D. J.** (1991). 16S ribosomal DNA amplification for phylogenetic study. *Journal of bacteriology* **173**, 697-703.
- Whitlock, J. E., Jones, D. T. and Harwood, V. J.** (2002). Identification of the sources of fecal coliforms in an urban watershed using antibiotic resistance analysis. *Water Research* **36**, 4273-428.
- Williams, D., Guy, C., McGarry, J., Guy, F., Tasker, L., Smith, R., MacEachern, K., Cripps, P., Kelly, D. and Trees, A.** (2000). Neospora caninum-associated abortion in cattle: the time of experimentally-induced parasitaemia during gestation determines foetal survival. *Parasitology* **121**, 347-358.
- Wilkinson, P. G.** (2011). *Characterisation of the Bacterial Flora Associated with the Grey Field Slug Deroceras reticulatum and Assessment of its Suitability as a Target for Biological Control*. PhD Thesis, University of Edinburgh

- Wilson, M. J., D. M. Glen and S. K. George** (1993). The rhabditid nematode *Phasmarhabditis hermaphrodita* as a potential biological control agent for slugs. *Biocontrol Science and Technology* **3**, 503–511.
- Wilson M J, Glen D M, George S K, Pearce J D and Wiltshire C W.** (1994) Biological control of slugs in winter wheat using the rhabditid nematode *Phasmarhabditis hermaphrodita*. *Annals of Applied Biology* **125**, 377-390.
- Wright, E. P.** (1982). The occurrence of *Campylobacter jejuni* in dog faeces from a public park. *Journal of Hygiene* **89**: 191-194.
- Yagi, S., Kitai, S. and Kimura, T.** (1971). Oxidation of element sulphur to thiosulphur by *Streptomyces*. *Archives of Microbiology* **22**, 157-159.
- Zhang, Y. a., Dube, M., McLean, D. and Kates, M.** (2003). Biodiesel production from waste cooking oil: 1. Process design and technological assessment. *Bioresource technology* **89**, 1-16.



Appendices



Appendix One: Preparation of chemical solutions and reagents

1.1 Solutions used in analysis of inorganic N-ions:

1.1.1 Indophenol blue method for the determination of ammonium-N (NH_4^{+-}N).

(a) Standard ammonium solution..

Standard ammonium solution was prepared by dissolving 0.4717 g ammonium sulphate in 1 litre distilled water for (100 pg N ml^{-1}). (Wainwright and Pugh, 1973)

(A) Ethylenediaminetetraacetic acid (EDTA)

EDTA (60g) was dissolved in (900ml) distilled water then diluted to 1 litre

(b) Phenol solution:

Phenol solution was prepared by dissolving phenol (62.5 g) in ethanol (25 ml) and adding acetone (12.5 ml) to give a total of 100 ml. The phenol solution was stored in the dark at 4°C .

(c) Phenolate reagent:

Phenolate reagent was prepared by mixing 20 ml of phenol solution with 20 ml caustic solution (27%NaOH w/v) and diluting to 100 ml. The reagent was prepared fresh daily.

1.1.2 Chromotropic acid method for nitrate-N determination.

(a) Standard nitrate solution:

Potassium nitrate (KNO_3), 0.229 g was dissolved in distilled water and made up to 1 litre volumetrically, for $100 \text{ NO}_3\text{-N ml}^{-1}$.

(b) Chromotropic acid reagent ($\text{C}_{10} \text{H}_6\text{O}_8\text{S}_2 \text{Na}_2$):

A 0.1% (v/v) stock solution of chromotropic acid in concentrated sulphuric acid (H_2SO_4) was prepared by dissolving 1.84 g chromotropic acid in 1 litre H_2SO_4 (1:1). This solution was stored in an amber bottle in refrigerator at 4°C for several months.

(c) Working chromotropic acid solution (CTA):

A working CTA-solution (0.01% v/v) was prepared by diluting 100 ml of stock solution to 990 ml with concentrated sulphuric acid (H₂SO₄) then adding 10 ml concentrated HCl. This solution was stored at 4°C for several weeks only

1.1.3 Analysis of inorganic S-ions.

(a) Standard sulphate-S solution:

Sodium sulphate (Na₂SO₄·10H₂O), 0.443 g was dissolved in 1 litre distilled water, which gives the concentration 100 µg SO₄²⁻ S ml⁻¹.

b) Gum acacia solution (0.250% w/v):

Gum acacia 0.25 g was dissolved in 100 ml distilled water.

1.3.4 Analysis of phosphate

(a) Chemical solution

1- Ascorbic acid solution (10%) was prepared by mixing 10 ml of ascorbic acid with 90 ml of distilled water

2- Ammonium molybdate (0.42% v/v) was prepared by dissolving 42 g of ammonium molybdate with 100 ml of in (1N) H₂SO₄

(b) Working solution

Working solution was prepared by mixing 1 volume of ascorbic acid solution (10%) and 6 volumes of ammonium molybdate (0.42%)

2. Elements standard curves

2.1 Standard ammonium solution (Wainwright and Pugh, 1973)

3.66 g of (NH₄)₂SO₄ were dissolved in 1 litre then diluted the solution 10 times (10 ml of ammonium solution with 90 ml distilled water) = 100 µg/NH₄⁺-N ml⁻¹. 2 ml of previous solution was added to 1 ml of EDTA (6% w/v) 7 ml of distilled water, 5 ml of phenolate reagent and 3 ml of sodium hypochlorite solution (10% v/v). The reaction mixture was mixed thoroughly and incubated at 25°C for 20 min in the dark. The volume

was made up to 50ml and mixed and the concentration of the indophenols-blue complex was measured at 630 nm.

2.2 Standard Nitrate solution (Sims and Jackson, 1971)

Weight 0.137g of NaNO_3 and were dissolved to 100ml of distilled water. Resulting solution is 1mg nitrate (NaNO_3) per ml. Solution was pipette to 6 tubes following volumes of the standard solution: (1 μl =1 μg nitrate) (0 μl , 10 μl , 20 μl , 40 μl , 80 μl , 100 μl). 3ml of filtrate was mixed to 7ml chromotropic acid (work solution) then incubated in water bath for 45 min in 40°C. The yellow colour formed was measured at 41 nm using spectrophotometer and the concentration of nitrate was determined by reference to a standard curve of nitrate concentration. (Fig 1 2)

2.3 Standard sulphate-s solution (Hesse, 1971)

1.47 g of (Na_2SO_4) and were dissolved in 1 litre of distilled water, which give the concentration **1000 $\mu\text{g}/\text{SO}_4^{-2}\text{-S ml}^{-1}$** then diluted the solution 10 times (10 ml of sodium sulphate solution with 90 ml of distal water) = **100 $\mu\text{g}/\text{SO}_4^{-2}\text{-S ml}^{-1}$** .

5ml of previous solution were added to 1g barium chloride BaCl_2 and 2ml of gum acacia (0.25% w/v) mixed well then the volume was made up to 25ml with distilled water. The white suspension resulting from precipitation of sulphate was measured at 470 nm. (Fig 13)

2.4 Standard phosphate solution (Hesse, 1971)

Weight 0.4393g of potassium dihydrogen (KH_2PO_4) into a litre volumetric flask. This solution was diluted with distilled water to produce 10 μg , 20 μg , 40 μg , 50 μg , 60 μg , 70 μg , 80 μg , 90 μg and 100 μg . The control was distilled water without KH_2PO_4 .

Mix 1 vol. of ascorbic acid mixed with 6 vol. of ammonium molybdate. The add 0.7 ml of working solution to 0.3 ml of sample and incubate at 37°C for 1 hour and read with blue colour at 820 nm. (Fig 1 4)




Fig1. 1 Ammonium standard curve




Fig1 2: Nitrate standard curve




Fig 1 3: Sulphate standard curve




Fig 1 4: Phosphate standard curve

Appendix Two: Identification of bacteria

1. Solid media

1.1 BRILLIANCEUTI CLARITY AGAR

Brillianceuti Clarity Agar (formerly Chromogenic UTI Clarity Agar) is for differentiation and presumptive identification of common urinary tract infection isolates.

Table 2 1 Composition of Brilliant clarity agar medium

| Typical Formula* | Gm/ litre |
|----------------------|--------------|
| Peptone..... | 9.00 |
| Chromogenic Mix..... | 17.00 |
| Tryptophan..... | 1.00 |
| Agar..... | 10.00 |
| pH 7.0 ± 0.2 @ 25°C | |

Interpretation

The expected colour reactions are laid out in Table 2 2

Table 2 2 Typical colour reactions on *Brilliance*UTI Clarity Agar


| Organism | β-galactosidase | β-glucosidase | TDA | Colony colour |
|---------------------------|-----------------|---------------|-----|--------------------|
| <i>E. coli</i> | + | | | Pink |
| enterococci | | + | | Blue / Turquoise |
| coliforms | + | + | | Dark Blue / Purple |
| <i>Proteus/Morganella</i> | | | + | Brown halo |
| pseudomonads | | | | Green/ Brown |
| staphylococci | | | | White / Cream |
| <i>S. saprophyticus</i> | | | | Pale Pink / White |
| streptococci | | | | White |

. Appearance

Dehydrated *Brilliance*UTI Clarity Agar is a free-flowing straw coloured powder. The prepared medium is a straw-coloured, transparent agar.

Table 2 1 Quality control

| Positive controls | Expected results |
|---|------------------------------------|
| <i>Escherichia coli</i> ATCC®25922* | Good growth; pink colonies |
| <i>Enterobacter aerogenes</i> ATCC®13048* | Good growth; purple colonies |
| <i>Enterococci</i> Fig 2 1: <i>E. coli</i> on Brilliant agar / Author's photo | |
| <i>Proteus mirabilis</i> NCTC10975 or <i>Proteus mirabilis</i> ATCC®29906* | Good growth; brown colony and halo |
| <i>Staphylococcus aureus</i> ATCC®25923* | Good growth; typical appearance |
| Negative control | |
| Uninoculated medium | |



E. coli on Brillianceuti Clarity Agar by (Authors)

1.2 CHROMagar Orientation

The principle of this medium is the use of Chromogenic substrates revealing metabolic enzymes. Dehydrated powder was provided by the CHROMagar Company, Paris, France.

The medium is composed of 16g each of peptone, meat, and yeast extracts and 15g of agar per Litre and a special chromogenic mixture. The powder was introduced into an Automatic pippeter, and sterilization process was performed at 120°C for 15 min. Samples were Streaked onto plate and incubate at 37°C for 18-24 hours.

Table 2 4: Composition of Chromagar Orientation medium

| Typical Formula..... | gm/litre |
|--------------------------------|---------------|
| Agar..... | 15.0 g |
| Peptone and yeast extract..... | 17.0 |
| Chromogenic mix..... | 1.0 g |
| Total..... | 33 g/L |

pH 7.0 ± 0.2 @ 25°C




FIG. 1. Representative chromogenic reactions of selected organisms on CHROMagar. 1, *E. coli*; 2, *P. vulgaris*; 3, *P. mirabilis*; 4, *E. cloacae*; 5, *C. diversus*; 6, *Klebsiella pneumoniae*; 7, *Salmonella typhimurium*; 8, *Aeromonas hydrophila*; 9, *Shigella sonnei*; 10, *P. aeruginosa*; 11, *Y. enterocolitica*; 12, *M.morganii*; 13, *Plesiomonas shigelloides*; 14, *Serratia marcescens*; 15, *A. calcoaceticus*; 16, *Stenotrophomonas maltophilia*; 17, *Enterococcus faecalis*.

(Merlino *et al.*, 1996)

Fig, 2 2: The colour of colonies of CHROMagar Orientation media reference

1.3 CHROMagar MRSA

BBL CHROMagar MRSA is a selective and differential medium, which incorporates cefoxitin, for the detection of MRSA from specimens.

For pre-weighed dose of medium of CHROMagaTMRSA powder, add dry powder to the corresponding volume of purified water. Alternatively, suspend the powder slowly in water by rotating for swelling of the agar. Heat and bring to boiling (100°C) while swirling or stirring regularly. If using an autoclave, do so without pressure. Do not heat to more than 100°C. the mixture may also be brought to a boil in a microwave oven: after initial boiling, remove from oven, stir gently, then return to oven for short repeated

Table 2 5: Composition of Chromagar MRSA

| Typical Formula..... | | gm/litre |
|----------------------|----------|---------------|
| Chromopeptone..... | | 40.0 g |
| Sodium..... | Chloride | 25.0 g |
| Chromogenic mix..... | | 0.5 g |
| Inhibitory..... | Agents | 0.07 g |
| Cefoxitin..... | | 6.0mg |
| Agar..... | | 14.0 g |
| pH 7.0 ± 0.2 @ 25°C | | |

Table 2 6: Quality control

| Positive control: | Expected result |
|---|--|
| <i>Staphylococcus aureus</i> ATCC® 29213 | Inhibition (partial to complete); N/A |
| * <i>Staphylococcus aureus</i> ATCC®25923 | Inhibition (partial to complete); N/A |
| * <i>Staphylococcus aureus</i> ATCC®43300 | Growth; Mauve |
| <i>Staphylococcus aureus</i> ATCC ®3359 | Growth; Mauve |
| <i>Enterococcus faecalis</i> ATCC ®29212 | Growth; blue |

(Garner, 1996 ; Clinical and Laboratory Standards Institute. 2008)

1.4 Czapek Dox agar

Code: CM0097 / **Oxoid**

Semi-synthetic solid medium, containing sucrose as C-source and nitrate as the sole source of nitrogen, useful for the general cultivation of fungi, yeasts and soil bacteria.

Recommended by Czapek (1902-1903) and Dox (1910). Table 2 1 1

Table 2.7: Composition of Czapek Dox agar medium

| Typical Formula | gm/litre |
|---|---------------|
| Na NO..... | 2.00 g |
| KCl..... | 0.50 g |
| Magnesium glycerophosphate..... | 0.50 g |
| Fe SO ₄ (7H ₂ O)..... | 0.01 g |
| K ₂ SO ₄ | 0.35 g |
| Sucrose..... | 30.0 g |
| Agar (Oxoid No. 3)..... | 12.0 g |

pH 6.8 ± 0.2 @ 25°C

Directions

A proprietary formulation (Oxoid) of Czapek Dox Agar was used for fungal growth. It was prepared by suspending 45.4 g of the powder in a litre of distilled water. The medium Was dissolved, and the pH adjusted to 6.8, and sterilised by autoclaving at 121° C for 15 min.

Appearance

Dehydrated medium: White coloured, free-flowing powder

Prepared medium: Off-white coloured gel

Table 2 8: Cultural characteristics after 24-48 hours at 35°C.

| Organisms (ATCC) | Growth |
|---|--------|
| <i>Aspergillus braseliensis</i> (16404) | +++ |
| <i>Saccharomyces cerevisiae</i> (9763) | +++ |
| <i>Candida albicans</i> (10231) | ++ |
| <i>Bacillus subtilis</i> (6633) | ++ |
| <i>Staphylococcus aureus</i> (25923) | - |

1.5 HiCrome (TM) E. coli Agar A

Code: 70722/ Fluka/ Sigma

HiCrome *E.coli* Agar B is recommended for the detection and enumeration of Escherichia coli in foods without further confirmation on membrane filter or by indole reagent.

Table 2 9: Composition of HiCrome (TM) E. coli Agar A

| Formula | gm/litre |
|---------|----------|
| | |

| | | |
|---------------------|------------|----------------|
| Casein | enzymic | 14.0 g |
| hydrolysate..... | | |
| Peptone | | 5.0 g |
| especial..... | | |
| Bile | salts | 1.50 g |
| mixture..... | | |
| Disodium | | 1.0 g |
| hydrogen..... | | |
| Sodium | dihydrogen | 0.6 g |
| phosphate..... | | |
| Sodium | | 2.4 g |
| chloride..... | | |
| Glucuronide..... | | 0.075 g |
| Agar | | 12.0 g |
| pH 7.2 ± 0.2 @ 25°C | | |

Directions :

Suspend 36,6g in 1 litre distilled water. Sterilize by autoclaving at 121°C for 15 minutes. Cool to 50°C and pour into sterile petri plates

Table 2 10: Culture characterisation after 18-24 hours at 44°C

| Positive control: | Expected results |
|---|------------------------------|
| <i>Escherichia coli</i> ATCC® 25922 * | Luxuriant; blue |
| <i>Klebsiella pneumoniae</i> ATCC® 13883 | luxuriant colourless, mucoid |
| <i>Salmonella enteritidis</i> ATCC® 13076 | luxuriant colourless |
| <i>Staphylococcus aureus</i> ATCC® 25923 | inhibited |

Anderson and Baird-Parker., 1975; Hansen and Yourassawsky



E. coli

Fig 2 3: *E. coli* on HiCrome agar / Author's photo

1.6 Nutrient Agar (Oxoid)

Code: CM0003 (Powder)

A general purpose medium which may be enriched with up to 10% blood or other biological fluid.

Table 2 11: Composition of Nutrient Agar

| Typical Formula..... | gm/litre |
|---------------------------------|---------------|
| 'Lab-Lemco' powder (Oxoid)..... | 1.0 g |
| Yeast extract..... | 2.0 e |
| Peptone..... | 5.0 g |
| Sodium chloride..... | 5.0 g |
| Agar..... | 15.0 g |

pH 7.4 ± 0.2 @ 25°C

Directions

The medium was prepared by suspending 28g in litre of distilled water, boiled dissolve completely. Sterilised by autoclaving at 121°C for 15 minutes.

Table 2 2 12: Quality control

| Positive controls: | Expected results |
|--|-----------------------------------|
| <i>Staphylococcus aureus</i> ATCC® 25923 * | Good growth; straw/white colonies |
| <i>Escherichia coli</i> ATCC® 25922 * | Good growth; straw colonies |
| Negative control: | |
| Uninoculated medium | No change |

(Lapage *et al.* 1970)

1.7 Plate Count Agar

The medium was prepared by suspending 17.5g in litre of distilled water, boiled to dissolve completely, pH 7.0, sterilised by autoclaving at 121°C for 15 minutes' Silica gel medium (Parkinson et al.,1989)

Table 2 13: Composition Plate Count Agar (Oxoid)

| Typical Formula..... | gm/litre |
|----------------------|--------------|
| Tryptone..... | 1.0 g |
| Yeast extract..... | 2.5 e |
| Glucose..... | 1.0 g |
| Agar..... | 9.0 g |

pH 7.0 ± 0.2 @ 25°C

Directions

Add 17.5g to 1 litre of distilled water. Dissolve by bringing to the boil with frequent stirring, mix and distribute into final containers. Sterilise by autoclaving at 121°C for 15 minutes.

Appearance

Dehydrated medium: Straw coloured, free-flowing powder
Prepared medium: Straw coloured gel

Table 2 14: Quality control

| Positive control: | Expected results |
|---------------------------------------|--------------------------------------|
| <i>Escherichia coli</i> ATCC® 25922 * | Good growth; straw coloured colonies |
| Negative control: | |
| Uninoculated plate | No change |

(PHLS, 1999)

1.8 POTATO DEXTROSE AGAR (EP/USP/JP/BP)

Code: CM0139 (Oxoid)

For the detection and enumeration of yeasts and moulds in butter and other dairy and food products. Also for the preparation of *Aspergillus niger* for the Harmonised Microbial Limit Tests from EP/USP/JP (enumeration test)

Table 2 15: Composition of PAD medium

| Formula | gm/litre |
|------------|-------------|
| Potato | |
| tract..... | 4.0* |
| Glucose | 20.0 |
| | |
| Agar | 15.0 |
| | |

pH 5.6 ± 0.2 @ 25°C

Directions

Suspend 39g in 1 litre of water (purified as requested). Bring to the boil to dissolve completely. Sterilise by autoclaving at 121°C for 15 minutes. Mix well before pouring.

Appearance

Dehydrated medium: Off-white, free-flowing powder

Prepared medium: Light straw coloured gel

Table 2 16: Quality Control

| Positive control: | Expected result |
|---|-----------------------------------|
| <i>Aspergillus fumigatus</i> ATCC® 9197 * | White mycelium, blue green spores |
| Negative controls: | |
| Uninoculated medium | No change |
| At pH 3.5 <i>Bacillus subtilis</i> ATCC® 6633 * | No growth |

American Public Health Association. (1992)

1.10 XLT-4 AGAR

Code: CM1061 from / **Oxoid**

A highly selective medium for isolation and identification of Salmonellae from clinical, environmental and food samples.

Table 17; Composition of XLT-4 Agar

| Typical Formula* | gm/litre |
|------------------------------|-------------|
| Proteose Peptone | 1.6 |
| Yeast extract..... | 3.0 |
| Lysine..... | 5.0 |
| Xylose..... | 3.75 |
| Lactose..... | 7.5 |
| Sucrose..... | 7.5 |
| Ferric ammonium citrate..... | 0.8 |
| Sodium thiosulphate..... | 6.8 |
| Sodium chloride..... | 5.0 |
| Phenol Red..... | 0.08 |
| Agar..... | 18.0 |

pH 7.4 ± 0.2 @ 25°C

Directions

Suspend 59g of XLT-4 Agar Base in 1 litre of distilled water, add 4.6ml of XLT-4 Selective Supplement and bring the medium to the boil.

Do not overheat, do not autoclave.

Cool to approximately 50°C and pour into sterile Petri dishes. It is not advised to hold the medium at 50°C for longer than 1 hour as this may cause the medium to precipitate.

Description

XLT-4 (Xylose Lactose Tergitol™ 4) Agar is a highly selective plating medium used for isolation and identification of salmonellae from clinical, environmental and food samples according to Miller

b) XLT-4 SELECTIVE SUPPLEMENT

Code: SR0237/ Oxoid

Table 2 18: Composition of XLT-4 selective supplement

| Supplement available in 100 ml (SR0237C) | per litre |
|--|-----------|
| Tergitol™ | 4.6ml |
| 4..... | |

Appearance:

Dehydrated medium: straw coloured, free-flowing powder
 Prepared medium: clear red gel

Table 9: Quality control

| Positive control: | Expected results |
|--|--|
| <i>Salmonella enteritidis</i> ATCC®13076 * | Good growth: black or, red with black centre |
| Negative controls: | |
| <i>Escherichia coli</i> ATCC®25922 * | Reduced growth, yellow |
| <i>Enterococcus faecalis</i> ATCC®29212 * | |

(Miller and Tate, 1990; Dusch and Altwegg, 1995)



Salmonella enterica

Fig 2 4: *S. enterica* on XLT-4 agar / Author's photo

gb|GQ418110.1| Uncultured Salmonella sp. clone F4jun.23 16S ribosomal RNA gene, partial sequence
 Length=1463

Score = 481 bits (260), Expect = 1e-132
 Identities = 261/262 (99%), Gaps = 0/262 (0%)
 Strand=Plus/Minus

```


Query 1      TTCTTTTGAACCCACTCCCATGGTGTGACGGGCGGTGTGTACAAGGCCCGGGAACGTAT 60
              |||
3bjct 1405   TTCTTTTGAACCCACTCCCATGGTGTGACGGGCGGTGTGTACAAGGCCCGGGAACGTAT 1346

Query 61     TCACCGTGGCATTCTGATCCACGATTACTAGCGATTCCGACTTCATGGAGTCGAGTTGCA 120
              |||
3bjct 1345   TCACCGTGGCATTCTGATCCACGATTACTAGCGATTCCGACTTCATGGAGTCGAGTTGCA 1286

Query 121    GACTCCAATCCGGACTACGACATACTTTATGAGGTCGCGTTGTCTCGCGAGGTCGCTTC 180
              |||
3bjct 1285   GACTCCAATCCGGACTACGACATACTTTATGAGGTCGCGTTGTCTCGCGAGGTCGCTTC 1226

Query 181    TCTTTGTATATGCCATTGTAGCACGTGTGTAGCCCTGGTCGTAAGGNCCATGATGACTTG 240
              |||
3bjct 1225   TCTTTGTATATGCCATTGTAGCACGTGTGTAGCCCTGGTCGTAAGGNCCATGATGACTTG 1166

Query 241    ACGTCATCCCCACCTTCCTCCA 262
              |||
3bjct 1165   ACGTCATCCCCACCTTCCTCCA 1144
  
```



> [gb|JF718785.1](#) Trichoderma asperellum strain C3 internal transcribed spacer 1, partial sequence; 5.8S ribosomal RNA gene, complete sequence; and internal transcribed spacer 2, partial sequence
 Length=541

Score = 435 bits (482), Expect = 5e-119
 Identities = 243/244 (99%), Gaps = 0/244 (0%)
 Strand=Plus/Plus

```

Query 1   ACTCGTTCTGTAGTCCCCTCGGGACGTATTTCTTACAGCTCTGAGCAAAAATTCAAAT 60
          |||||
Sbjct 135  ACTCTTTCTGTAGTCCCCTCGGGACGTATTTCTTACAGCTCTGAGCAAAAATTCAAAT 194

Query 61   GAATCAAAACTTTCAACAACGGATCTCTTGTTCTGGCATCGATGAAGAACGCAGCGAAA 120
          |||||
Sbjct 195  GAATCAAAACTTTCAACAACGGATCTCTTGTTCTGGCATCGATGAAGAACGCAGCGAAA 254

Query 121  TCGATAAGTAATGTGAATTGCAGAATTCAGTGAATCATCGAATCTTTGAACGCACATTG 180
          |||||
Sbjct 255  TCGATAAGTAATGTGAATTGCAGAATTCAGTGAATCATCGAATCTTTGAACGCACATTG 314

Query 181  CGCCCGCCAGTATTCTGGCGGGCATGCCTGTCCGAGCGTCATTTCAACCCTCGAACCCT 240
          |||||
Sbjct 315  CGCCCGCCAGTATTCTGGCGGGCATGCCTGTCCGAGCGTCATTTCAACCCTCGAACCCT 374

Query 241  CCGG 244
          ||||
Sbjct 375  CCGG 378
  
```





Fig 2 5: g DNA of isolated bacterial species; (Lane 1) *Salmonella* and (Lane 2) *E. coli*, (lane L); hyper ladder




Fig 2 6: PCR- 16Sr RNA, amplification products of *Salmonella enteric* species analyzed by electrophoresis in agarose gel lanes represent; (lane L); hyper ladder;(lane lane 1); *S. enterica* R1, (lane 2); *S. enterica* R2 and (lane 3) positive control.



Fig 2 7: PCR- 16Sr RNA, amplification products of *Salmonella enteric* species analyzed by electrophoresis in agarose gel lanes represent; (lane L); hyper ladder;(lane lane 1); *S. enterica* R1, (lane 2); *S. enterica* R2 negative result 1.




Fig 2 8: PCR- 16Sr RNA, amplification products of *E. coli* analyzed by electrophoresis in agarose gel lanes represent; (lane L); hyper ladder;(lane lane 1); *E.coli* .

Appendix Three: ANOVA tables

Three Way Analysis of Variance

Saturday, March 17, 2012, 14:06:28

Data source: E. coli ANOVA of BSFL in BSFL AONVA E. coli

Balanced Design (No Interactions)

Dependent Variable: Y

Normality Test: Failed (P < 0.050)

Equal Variance Test: Passed (P = 1.000)

| Source of Variation | DF | SS | MS | F | P |
|---------------------|----|------------|------------|----------|--------|
| T | 1 | 1.974E+015 | 1.974E+015 | 431.898 | <0.001 |
| R | 4 | 1.914E+016 | 4.784E+015 | 1046.919 | <0.001 |
| C | 2 | 2.713E+013 | 1.357E+013 | 2.969 | 0.109 |
| Residual | 8 | 3.656E+013 | 4.570E+012 | | |
| Total | 29 | 2.724E+016 | 9.394E+014 | | |

The difference in the mean values among the different levels of T are greater than would be expected by chance after allowing for the effects of differences in R and C. There is a statistically significant difference (P = <0.001). To isolate which group(s) differ from the others use a multiple comparison procedure.

The difference in the mean values among the different levels of R are greater than would be expected by chance after allowing for the effects of differences in T and C. There is a statistically significant difference (P = <0.001). To isolate which group(s) differ from the others use a multiple comparison procedure.

The difference in the mean values among the different levels of C are not great enough to exclude the possibility that the difference is just due to random sampling variability after allowing for the effects of differences in T and R. There is not a statistically significant difference (P = 0.109).

All Pairwise Multiple Comparison Procedures (Holm-Sidak method):

Overall significance level = 0.05

Comparisons for factor: **T**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|--------|--------------|----------------|--------------|
| 2.000 vs. 1.000 | 16221816.000 | 20.782 | <0.001 | 0.050 | Yes |

Comparisons for factor: **R**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|--------|--------------|----------------|--------------|
| 1.000 vs. 5.000 | 63319106.667 | 51.305 | <0.001 | 0.005 | Yes |

| | | | | | |
|-----------------|--------------|--------|--------|-------|-----|
| 1.000 vs. 4.000 | 63293650.000 | 51.284 | <0.001 | 0.006 | Yes |
| 1.000 vs. 3.000 | 59012683.333 | 47.815 | <0.001 | 0.006 | Yes |
| 2.000 vs. 5.000 | 36459106.667 | 29.541 | <0.001 | 0.007 | Yes |
| 2.000 vs. 4.000 | 36433650.000 | 29.521 | <0.001 | 0.009 | Yes |
| 2.000 vs. 3.000 | 32152683.333 | 26.052 | <0.001 | 0.010 | Yes |
| 1.000 vs. 2.000 | 26860000.000 | 21.763 | <0.001 | 0.013 | Yes |
| 3.000 vs. 5.000 | 4306423.333 | 3.489 | 0.008 | 0.017 | Yes |
| 3.000 vs. 4.000 | 4280966.667 | 3.469 | 0.008 | 0.025 | Yes |
| 4.000 vs. 5.000 | 25456.667 | 0.0206 | 0.984 | 0.050 | No |

Comparisons for factor: **C**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-------------------|----------------------|----------|---------------------|-----------------------|---------------------|
| 1.000 vs. 3.000 | 2284562.000 | 2.390 | 0.044 | 0.017 | No |
| 2.000 vs. 3.000 | 1536809.000 | 1.608 | 0.147 | 0.025 | No |
| 1.000 vs. 2.000 | 747753.000 | 0.782 | 0.457 | 0.050 | No |

Power of performed test with alpha = 0.0500: for T : 1.000

Power of performed test with alpha = 0.0500: for R : 1.000

Power of performed test with alpha = 0.0500: for C : 0.288

Least square means for T :

Group Mean

1.000 12725337.333

2.000 28947153.333

Std Err of LS Mean = 551942.360

Least square means for R :

Group Mean

1.000 63333333.333

2.000 36473333.333

3.000 4320650.000

4.000 39683.333

5.000 14226.667

Std Err of LS Mean = 872697.497

Least square means for C :

Group Mean

1.000 21847017.000

2.000 21099264.000

3.000 19562455.000

Std Err of LS Mean = 675988.575

Data source: ANOVA OF BSFL Salmonella in BSFL ANOVA Salmonella

Balanced Design (No Interactions)

Dependent Variable: Y

Normality Test: Failed (P < 0.050)**Equal Variance Test:** Passed (P = 1.000)

| Source of Variation | DF | SS | MS | F | P |
|---------------------|----|------------------|-----------------|----------|--------|
| T | 1 | 2.293E+013 | 2.293E+013 | 955.133 | <0.001 |
| R | 4 | 3.360E+014 | 8.400E+013 | 3499.420 | <0.001 |
| C | 2 | 62712566000.000 | 31356283000.0 | | |
| Residual | 8 | 192037221999.938 | 24004652749.992 | | |
| Total | 29 | 4.441E+014 | 1.531E+013 | | |

The difference in the mean values among the different levels of T are greater than would be expected by chance after allowing for the effects of differences in R and C. There is a statistically significant difference (P = <0.001). To isolate which group(s) differ from the others use a multiple comparison procedure.

The difference in the mean values among the different levels of R are greater than would be expected by chance after allowing for the effects of differences in T and C. There is a statistically significant difference (P = <0.001). To isolate which group(s) differ from the others use a multiple comparison procedure.

The difference in the mean values among the different levels of C are not great enough to exclude the possibility that the difference is just due to random sampling variability after allowing for the effects of differences in T and R. There is not a statistically significant difference (P = 0.323).

All Pairwise Multiple Comparison Procedures (Holm-Sidak method):

Overall significance level = 0.05

Comparisons for factor: **T**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|--------|--------------|----------------|--------------|
| 2.000 vs. 1.000 | 1748433.333 | 30.905 | <0.001 | 0.050 | Yes |

Comparisons for factor: **R**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|--------|--------------|----------------|--------------|
| 1.000 vs. 5.000 | 8397250.000 | 93.875 | <0.001 | 0.005 | Yes |
| 1.000 vs. 4.000 | 8379783.333 | 93.680 | <0.001 | 0.006 | Yes |

| | | | | | |
|-----------------|-------------|--------|--------|-------|-----|
| 1.000 vs. 3.000 | 8256250.000 | 92.299 | <0.001 | 0.006 | Yes |
| 2.000 vs. 5.000 | 4309083.333 | 48.172 | <0.001 | 0.007 | Yes |
| 2.000 vs. 4.000 | 4291616.667 | 47.977 | <0.001 | 0.009 | Yes |
| 2.000 vs. 3.000 | 4168083.333 | 46.596 | <0.001 | 0.010 | Yes |
| 1.000 vs. 2.000 | 4088166.667 | 45.703 | <0.001 | 0.013 | Yes |
| 3.000 vs. 5.000 | 141000.000 | 1.576 | 0.154 | 0.017 | No |
| 3.000 vs. 4.000 | 123533.333 | 1.381 | 0.205 | 0.025 | No |
| 4.000 vs. 5.000 | 17466.667 | 0.195 | 0.850 | 0.050 | No |

Comparisons for factor: C

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|-------|--------------|----------------|--------------|
| 1.000 vs. 3.000 | 111920.000 | 1.615 | 0.145 | 0.017 | No |
| 1.000 vs. 2.000 | 59470.000 | 0.858 | 0.416 | 0.025 | No |
| 2.000 vs. 3.000 | 52450.000 | 0.757 | 0.471 | 0.050 | No |

Power of performed test with alpha = 0.0500: for T : 1.000
Power of performed test with alpha = 0.0500: for R : 1.000
Power of performed test with alpha = 0.0500: for C : 0.0826

Least square means for T :

Group Mean
1.000 1701493.333
2.000 3449926.667
Std Err of LS Mean = 40003.877

Least square means for R :

Group Mean
1.000 8400000.000
2.000 4311833.333
3.000 143750.000
4.000 20216.667
5.000 2750.000
Std Err of LS Mean = 63251.683

Least square means for C :

Group Mean
1.000 2632840.000
2.000 2573370.000
3.000 2520920.000
Std Err of LS Mean = 48994.543

Data source: BSFL AONVA Plate count media in Plate count media ANOVA

Balanced Design (No Interactions)

Dependent Variable: Y

Normality Test: Failed (P < 0.050)

Equal Variance Test: Passed (P = 1.000)

| Source of Variation | DF | SS | MS | F | P |
|---------------------|----|------------|------------|---------|--------|
| T | 1 | 1.581E+017 | 1.581E+017 | 102.181 | <0.001 |
| R | 4 | 7.626E+017 | 1.906E+017 | 123.201 | <0.001 |
| C | 2 | 7.304E+015 | 3.652E+015 | 2.360 | 0.156 |
| Residual | 8 | 1.238E+016 | 1.547E+015 | | |
| Total | 29 | 1.560E+018 | 5.378E+016 | | |

The difference in the mean values among the different levels of T are greater than would be expected by chance after allowing for the effects of differences in R and C. There is a statistically significant difference (P = <0.001). To isolate which group(s) differ from the others use a multiple comparison procedure.

The difference in the mean values among the different levels of R are greater than would be expected by chance after allowing for the effects of differences in T and C. There is a statistically significant difference (P = <0.001). To isolate which group(s) differ from the others use a multiple comparison procedure.

The difference in the mean values among the different levels of C are not great enough to exclude the possibility that the difference is just due to random sampling variability after allowing for the effects of differences in T and R. There is not a statistically significant difference (P = 0.156).

All Pairwise Multiple Comparison Procedures (Holm-Sidak method):
Overall significance level = 0.05

Comparisons for factor: **T**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|--------|--------------|----------------|--------------|
| 2.000 vs. 1.000 | 145196940.000 | 10.108 | <0.001 | 0.050 | Yes |

Comparisons for factor: **R**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|--------|--------------|----------------|--------------|
| 2.000 vs. 5.000 | 363737183.333 | 16.016 | <0.001 | 0.005 | Yes |
| 2.000 vs. 4.000 | 361496500.000 | 15.917 | <0.001 | 0.006 | Yes |

| | | | | | |
|-----------------|---------------|--------|--------|-------|-----|
| 2.000 vs. 3.000 | 359237333.333 | 15.818 | <0.001 | 0.006 | Yes |
| 1.000 vs. 5.000 | 283270516.667 | 12.473 | <0.001 | 0.007 | Yes |
| 1.000 vs. 4.000 | 281029833.333 | 12.374 | <0.001 | 0.009 | Yes |
| 1.000 vs. 3.000 | 278770666.667 | 12.275 | <0.001 | 0.010 | Yes |
| 2.000 vs. 1.000 | 80466666.667 | 3.543 | 0.008 | 0.013 | Yes |
| 3.000 vs. 5.000 | 4499850.000 | 0.198 | 0.848 | 0.017 | No |
| 3.000 vs. 4.000 | 2259166.667 | 0.0995 | 0.923 | 0.025 | No |
| 4.000 vs. 5.000 | 2240683.333 | 0.0987 | 0.924 | 0.050 | No |

Comparisons for factor: C

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|-------|--------------|----------------|--------------|
| 1.000 vs. 2.000 | 37619750.000 | 2.138 | 0.065 | 0.017 | No |
| 1.000 vs. 3.000 | 24658710.000 | 1.402 | 0.199 | 0.025 | No |
| 3.000 vs. 2.000 | 12961040.000 | 0.737 | 0.482 | 0.050 | No |

Power of performed test with alpha = 0.0500: for T : 1.000

Power of performed test with alpha = 0.0500: for R : 1.000

Power of performed test with alpha = 0.0500: for C : 0.209

Least square means for T :

Group Mean

1.000 58213993.333

2.000 203410933.333

Std Err of LS Mean = 10156812.630

Least square means for R :

Group Mean

1.000 283333333.333

2.000 363800000.000

3.000 4562666.667

4.000 2303500.000

5.000 62816.667

Std Err of LS Mean = 16059330.839

Least square means for C :

Group Mean

1.000 151571950.000

2.000 113952200.000

3.000 126913240.000

Std Err of LS Mean = 12439504.178

Data source: E. coli ANOVA in FBL E. coli ANOVA

Balanced Design (No Interactions)

Dependent Variable: Y

Normality Test: Failed (P < 0.050)**Equal Variance Test:** Passed (P = 1.000)

| Source of Variation | DF | SS | MS | F | P |
|---------------------|----|------------|------------|---------|--------|
| R | 1 | 8.554E+015 | 8.554E+015 | 455.393 | <0.001 |
| T | 4 | 3.470E+016 | 8.676E+015 | 461.905 | <0.001 |
| C | 2 | 4.231E+013 | 2.115E+013 | 1.126 | 0.371 |
| Residual | 8 | 1.503E+014 | 1.878E+013 | | |
| Total | 29 | 7.776E+016 | 2.681E+015 | | |

The difference in the mean values among the different levels of R are greater than would be expected by chance after allowing for the effects of differences in T and C. There is a statistically significant difference (P = <0.001). To isolate which group(s) differ from the others use a multiple comparison procedure.

The difference in the mean values among the different levels of T are greater than would be expected by chance after allowing for the effects of differences in R and C. There is a statistically significant difference (P = <0.001). To isolate which group(s) differ from the others use a multiple comparison procedure.

The difference in the mean values among the different levels of C are not great enough to exclude the possibility that the difference is just due to random sampling variability after allowing for the effects of differences in R and T. There is not a statistically significant difference (P = 0.371).

All Pairwise Multiple Comparison Procedures (Holm-Sidak method):

Overall significance level = 0.05

Comparisons for factor: **R**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|--------|--------------|----------------|--------------|
| 2.000 vs. 1.000 | 33771066.667 | 21.340 | <0.001 | 0.050 | Yes |

Comparisons for factor: **T**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|--------|--------------|----------------|--------------|
| 2.000 vs. 5.000 | 88992000.000 | 35.566 | <0.001 | 0.005 | Yes |
| 2.000 vs. 4.000 | 88841000.000 | 35.505 | <0.001 | 0.006 | Yes |

| | | | | | |
|-----------------|--------------|--------|--------|-------|-----|
| 2.000 vs. 3.000 | 84650000.000 | 33.830 | <0.001 | 0.006 | Yes |
| 2.000 vs. 1.000 | 71783333.333 | 28.688 | <0.001 | 0.007 | Yes |
| 1.000 vs. 5.000 | 17208666.667 | 6.877 | <0.001 | 0.009 | Yes |
| 1.000 vs. 4.000 | 17057666.667 | 6.817 | <0.001 | 0.010 | Yes |
| 1.000 vs. 3.000 | 12866666.667 | 5.142 | <0.001 | 0.013 | Yes |
| 3.000 vs. 5.000 | 4342000.000 | 1.735 | 0.121 | 0.017 | No |
| 3.000 vs. 4.000 | 4191000.000 | 1.675 | 0.132 | 0.025 | No |
| 4.000 vs. 5.000 | 151000.000 | 0.0603 | 0.953 | 0.050 | No |

Comparisons for factor: C

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|-------|--------------|----------------|--------------|
| 1.000 vs. 3.000 | 2812600.000 | 1.451 | 0.185 | 0.017 | No |
| 1.000 vs. 2.000 | 2049200.000 | 1.057 | 0.321 | 0.025 | No |
| 2.000 vs. 3.000 | 763400.000 | 0.394 | 0.704 | 0.050 | No |

Power of performed test with alpha = 0.0500: for R : 1.000
Power of performed test with alpha = 0.0500: for T : 1.000
Power of performed test with alpha = 0.0500: for C : 0.0630

Least square means for R :

Group Mean
1.000 5377866.667
2.000 39148933.333
Std Err of LS Mean = 1119016.803

Least square means for T :

Group Mean
1.000 17333333.333
2.000 89116666.667
3.000 4466666.667
4.000 275666.667
5.000 124666.667
Std Err of LS Mean = 1769320.918

Least square means for C :

Group Mean
1.000 23884000.000
2.000 21834800.000
3.000 21071400.000
Std Err of LS Mean = 1370510.090

Three Way Analysis of Variance

Tuesday, January 17, 2012, 21:49:21

Data source: FBL Salmonella ANOVA in FBL Salmonella ANOVA

Balanced Design (No Interactions)

Dependent Variable: Y

Normality Test: Failed (P < 0.050)

Equal Variance Test: Passed (P = 1.000)

| Source of Variation | DF | SS | MS | F | P |
|---------------------|----|------------|------------|---------|--------|
| R | 1 | 2.473E+017 | 2.473E+017 | 237.919 | <0.001 |
| T | 4 | 7.879E+017 | 1.970E+017 | 189.502 | <0.001 |
| C | 2 | 2.672E+015 | 1.336E+015 | 1.285 | 0.328 |
| Residual | 8 | 8.315E+015 | 1.039E+015 | | |
| Total | 29 | 1.872E+018 | 6.454E+016 | | |

The difference in the mean values among the different levels of R are greater than would be expected by chance after allowing for the effects of differences in T and C. There is a statistically significant difference (P = <0.001). To isolate which group(s) differ from the others use a multiple comparison procedure.

The difference in the mean values among the different levels of T are greater than would be expected by chance after allowing for the effects of differences in R and C. There is a statistically significant difference (P = <0.001). To isolate which group(s) differ from the others use a multiple comparison procedure.

The difference in the mean values among the different levels of C are not great enough to exclude the possibility that the difference is just due to random sampling variability after allowing for the effects of differences in R and T. There is not a statistically significant difference (P = 0.328).

All Pairwise Multiple Comparison Procedures (Holm-Sidak method):
Overall significance level = 0.05

Comparisons for factor: **R**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|--------|--------------|----------------|--------------|
| 2.000 vs. 1.000 | 181583786.667 | 15.425 | <0.001 | 0.050 | Yes |

Comparisons for factor: **T**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|--------|--------------|----------------|--------------|
| 2.000 vs. 5.000 | 420681950.000 | 22.601 | <0.001 | 0.005 | Yes |
| 2.000 vs. 4.000 | 420623483.333 | 22.598 | <0.001 | 0.006 | Yes |

| | | | | | |
|-----------------|---------------|---------|--------|-------|-----|
| 2.000 vs. 1.000 | 386316666.667 | 20.754 | <0.001 | 0.006 | Yes |
| 2.000 vs. 3.000 | 385480000.000 | 20.709 | <0.001 | 0.007 | Yes |
| 3.000 vs. 5.000 | 35201950.000 | 1.891 | 0.095 | 0.009 | No |
| 3.000 vs. 4.000 | 35143483.333 | 1.888 | 0.096 | 0.010 | No |
| 1.000 vs. 5.000 | 34365283.333 | 1.846 | 0.102 | 0.013 | No |
| 1.000 vs. 4.000 | 34306816.667 | 1.843 | 0.103 | 0.017 | No |
| 3.000 vs. 1.000 | 836666.667 | 0.0449 | 0.965 | 0.025 | No |
| 4.000 vs. 5.000 | 58466.667 | 0.00314 | 0.998 | 0.050 | No |

Comparisons for factor: C

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|--------|--------------|----------------|--------------|
| 1.000 vs. 3.000 | 20346370.000 | 1.411 | 0.196 | 0.017 | No |
| 2.000 vs. 3.000 | 19675210.000 | 1.365 | 0.210 | 0.025 | No |
| 1.000 vs. 2.000 | 671160.000 | 0.0465 | 0.964 | 0.050 | No |

Power of performed test with alpha = 0.0500: for R : 1.000
Power of performed test with alpha = 0.0500: for T : 1.000
Power of performed test with alpha = 0.0500: for C : 0.0802

Least square means for R :

Group Mean
1.000 7304353.333
2.000 188888140.000
Std Err of LS Mean = 8324291.397

Least square means for T :

Group Mean
1.000 34400000.000
2.000 420716666.667
3.000 35236666.667
4.000 93183.333
5.000 34716.667
Std Err of LS Mean = 13161860.360

Least square means for C :

Group Mean
1.000 105102090.000
2.000 104430930.000
3.000 84755720.000
Std Err of LS Mean = 10195133.196

Three Way Analysis of Variance

Saturday, March 17, 2012, 20:02:03

Data source: FBL Plate count media ANOVA in FBL Plate count media ANOVA

Balanced Design (No Interactions)

Dependent Variable: Y

Normality Test: Failed (P < 0.050)

Equal Variance Test: Passed (P = 1.000)

| Source of Variation | DF | SS | MS | F | P |
|---------------------|----|------------|------------|----------|--------|
| R | 1 | 2.338E+017 | 2.338E+017 | 1114.570 | <0.001 |
| T | 4 | 8.676E+017 | 2.169E+017 | 1034.149 | <0.001 |
| C | 2 | 1.363E+014 | 6.814E+013 | 0.325 | 0.732 |
| Residual | 8 | 1.678E+015 | 2.097E+014 | | |
| Total | 29 | 1.815E+018 | 6.259E+016 | | |

The difference in the mean values among the different levels of R are greater than would be expected by chance after allowing for the effects of differences in T and C. There is a statistically significant difference (P = <0.001). To isolate which group(s) differ from the others use a multiple comparison procedure.

The difference in the mean values among the different levels of T are greater than would be expected by chance after allowing for the effects of differences in R and C. There is a statistically significant difference (P = <0.001). To isolate which group(s) differ from the others use a multiple comparison procedure.

The difference in the mean values among the different levels of C are not great enough to exclude the possibility that the difference is just due to random sampling variability after allowing for the effects of differences in R and T. There is not a statistically significant difference (P = 0.732).

All Pairwise Multiple Comparison Procedures (Holm-Sidak method):
Overall significance level = 0.05

Comparisons for factor: **R**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|--------|--------------|----------------|--------------|
| 2.000 vs. 1.000 | 176550300.000 | 33.385 | <0.001 | 0.050 | Yes |

Comparisons for factor: **T**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|--------|--------------|----------------|--------------|
| 2.000 vs. 5.000 | 452886583.333 | 54.163 | <0.001 | 0.005 | Yes |
| 2.000 vs. 4.000 | 452667666.667 | 54.137 | <0.001 | 0.006 | Yes |

| | | | | | |
|-----------------|---------------|--------|--------|-------|-----|
| 2.000 vs. 3.000 | 403100000.000 | 48.209 | <0.001 | 0.006 | Yes |
| 2.000 vs. 1.000 | 309666666.667 | 37.035 | <0.001 | 0.007 | Yes |
| 1.000 vs. 5.000 | 143219916.667 | 17.128 | <0.001 | 0.009 | Yes |
| 1.000 vs. 4.000 | 143001000.000 | 17.102 | <0.001 | 0.010 | Yes |
| 1.000 vs. 3.000 | 93433333.333 | 11.174 | <0.001 | 0.013 | Yes |
| 3.000 vs. 5.000 | 49786583.333 | 5.954 | <0.001 | 0.017 | Yes |
| 3.000 vs. 4.000 | 49567666.667 | 5.928 | <0.001 | 0.025 | Yes |
| 4.000 vs. 5.000 | 218916.667 | 0.0262 | 0.980 | 0.050 | No |

Comparisons for factor: C

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|-------|--------------|----------------|--------------|
| 3.000 vs. 1.000 | 5214920.000 | 0.805 | 0.444 | 0.017 | No |
| 2.000 vs. 1.000 | 2822550.000 | 0.436 | 0.675 | 0.025 | No |
| 3.000 vs. 2.000 | 2392370.000 | 0.369 | 0.721 | 0.050 | No |

Power of performed test with alpha = 0.0500: for R : 1.000
Power of performed test with alpha = 0.0500: for T : 1.000
Power of performed test with alpha = 0.0500: for C : 0.0500

Least square means for R :

Group Mean
1.000 41060666.667
2.000 217610966.667
Std Err of LS Mean = 3739382.136

Least square means for T :

Group Mean
1.000 143333333.333
2.000 453000000.000
3.000 49900000.000
4.000 332333.333
5.000 113416.667
Std Err of LS Mean = 5912482.295

Least square means for C :

Group Mean
1.000 126656660.000
2.000 129479210.000
3.000 131871580.000
Std Err of LS Mean = 4579789.093

Larvae Guts

Three Way Analysis of Variance

Monday, March 19, 2012, 14:24:45

Data source: ANOVA BSFL gut *E. coli* in ANOVA BSFL gut *E. coli*

Balanced Design (No Interactions)

Dependent Variable: Col 4

Normality Test: Failed (P < 0.050)

Equal Variance Test: Passed (P = 1.000)

| Source of Variation | DF | SS | MS | F | P |
|---------------------|----|-----------------|----------------|---------|--------|
| T | 1 | 3491104687.500 | 3491104687.500 | 141.130 | <0.001 |
| C | 4 | 5039706996.667 | 1259926749.167 | 50.933 | <0.001 |
| R | 2 | 91331151.667 | 45665575.833 | 1.846 | 0.219 |
| Residual | 8 | 197894770.000 | 24736846.250 | | |
| Total | 29 | 14123564784.167 | 487019475.316 | | |

The difference in the mean values among the different levels of T are greater than would be expected by chance after allowing for the effects of differences in C and R. There is a statistically significant difference (P = <0.001). To isolate which group(s) differ from the others use a multiple comparison procedure.

The difference in the mean values among the different levels of C are greater than would be expected by chance after allowing for the effects of differences in T and R. There is a statistically significant difference (P = <0.001). To isolate which group(s) differ from the others use a multiple comparison procedure.

The difference in the mean values among the different levels of R are not great enough to exclude the possibility that the difference is just due to random sampling variability after allowing for the effects of differences in T and C. There is not a statistically significant difference (P = 0.219).

All Pairwise Multiple Comparison Procedures (Holm-Sidak method):

Overall significance level = 0.05

Comparisons for factor: **T**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|--------|--------------|----------------|--------------|
| 2.000 vs. 1.000 | 21575.000 | 11.880 | <0.001 | 0.050 | Yes |

Comparisons for factor: **C**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|--------|--------------|----------------|--------------|
| 2.000 vs. 1.000 | 34163.333 | 11.897 | <0.001 | 0.005 | Yes |
| 2.000 vs. 5.000 | 34017.500 | 11.847 | <0.001 | 0.006 | Yes |

| | | | | | |
|-----------------|-----------|--------|--------|-------|-----|
| 2.000 vs. 4.000 | 30211.667 | 10.521 | <0.001 | 0.006 | Yes |
| 2.000 vs. 3.000 | 19218.333 | 6.693 | <0.001 | 0.007 | Yes |
| 3.000 vs. 1.000 | 14945.000 | 5.205 | <0.001 | 0.009 | Yes |
| 3.000 vs. 5.000 | 14799.167 | 5.154 | <0.001 | 0.010 | Yes |
| 3.000 vs. 4.000 | 10993.333 | 3.828 | 0.005 | 0.013 | Yes |
| 4.000 vs. 1.000 | 3951.667 | 1.376 | 0.206 | 0.017 | No |
| 4.000 vs. 5.000 | 3805.833 | 1.325 | 0.222 | 0.025 | No |
| 5.000 vs. 1.000 | 145.833 | 0.0508 | 0.961 | 0.050 | No |

Comparisons for factor: **R**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-------------------|----------------------|----------|---------------------|-----------------------|---------------------|
| 1.000 vs. 3.000 | 4167.500 | 1.874 | 0.098 | 0.017 | No |
| 2.000 vs. 3.000 | 2904.500 | 1.306 | 0.228 | 0.025 | No |
| 1.000 vs. 2.000 | 1263.000 | 0.568 | 0.586 | 0.050 | No |

Power of performed test with alpha = 0.0500: for T : 1.000
 Power of performed test with alpha = 0.0500: for C : 1.000
 Power of performed test with alpha = 0.0500: for R : 0.145

Least square means for T :

Group Mean
 1.000 120.333
 2.000 21695.333
 Std Err of LS Mean = 1284.182

Least square means for C :

Group Mean
 1.000 266.667
 2.000 34430.000
 3.000 15211.667
 4.000 4218.333
 5.000 412.500
 Std Err of LS Mean = 2030.470

Least square means for R :

Group Mean
 1.000 12718.000
 2.000 11455.000
 3.000 8550.500
 Std Err of LS Mean = 1572.795

Three Way Analysis of Variance

Monday, March 19, 2012, 14:32:22

Data source: ANOVA BSFL gut Salmonella in ANOVA BSFL gut Salmonella

Balanced Design (No Interactions)

Dependent Variable: Col 4

Normality Test: Passed (P = 0.097)

Equal Variance Test: Passed (P = 1.000)

| Source of Variation | DF | SS | MS | F | P |
|---------------------|----|------------------|-----------------|--------|-------|
| T | 1 | 21075980853.333 | 21075980853.333 | 12.305 | 0.008 |
| C | 4 | 35539650118.133 | 8884912529.533 | 5.188 | 0.023 |
| R | 2 | 2254822678.400 | 1127411339.200 | 0.658 | 0.544 |
| Residual | 8 | 13702023586.667 | 1712752948.333 | | |
| Total | 29 | 124078968596.800 | 4278585124.028 | | |

The difference in the mean values among the different levels of T are greater than would be expected by chance after allowing for the effects of differences in C and R. There is a statistically significant difference (P = 0.008). To isolate which group(s) differ from the others use a multiple comparison procedure.

The difference in the mean values among the different levels of C are greater than would be expected by chance after allowing for the effects of differences in T and R. There is a statistically significant difference (P = 0.023). To isolate which group(s) differ from the others use a multiple comparison procedure.

The difference in the mean values among the different levels of R are not great enough to exclude the possibility that the difference is just due to random sampling variability after allowing for the effects of differences in T and C. There is not a statistically significant difference (P = 0.544).

All Pairwise Multiple Comparison Procedures (Holm-Sidak method):

Overall significance level = 0.05

Comparisons for factor: **T**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|-------|--------------|----------------|--------------|
| 2.000 vs. 1.000 | 53010.667 | 3.508 | 0.008 | 0.050 | Yes |

Comparisons for factor: **C**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|-------|--------------|----------------|--------------|
| 3.000 vs. 1.000 | 88390.667 | 3.699 | 0.006 | 0.005 | No |
| 3.000 vs. 5.000 | 87605.000 | 3.666 | 0.006 | 0.006 | No |

| | | | | | |
|-----------------|-----------|--------|-------|-------|----|
| 3.000 vs. 4.000 | 85338.333 | 3.572 | 0.007 | 0.006 | No |
| 3.000 vs. 2.000 | 48490.000 | 2.029 | 0.077 | 0.007 | No |
| 2.000 vs. 1.000 | 39900.667 | 1.670 | 0.133 | 0.009 | No |
| 2.000 vs. 5.000 | 39115.000 | 1.637 | 0.140 | 0.010 | No |
| 2.000 vs. 4.000 | 36848.333 | 1.542 | 0.162 | 0.013 | No |
| 4.000 vs. 1.000 | 3052.333 | 0.128 | 0.902 | 0.017 | No |
| 4.000 vs. 5.000 | 2266.667 | 0.0949 | 0.927 | 0.025 | No |
| 5.000 vs. 1.000 | 785.667 | 0.0329 | 0.975 | 0.050 | No |

Comparisons for factor: **R**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-------------------|----------------------|----------|---------------------|-----------------------|---------------------|
| 3.000 vs. 1.000 | 18648.400 | 1.008 | 0.343 | 0.017 | No |
| 3.000 vs. 2.000 | 18122.000 | 0.979 | 0.356 | 0.025 | No |
| 2.000 vs. 1.000 | 526.400 | 0.0284 | 0.978 | 0.050 | No |

Power of performed test with alpha = 0.0500: for T : 0.847

Power of performed test with alpha = 0.0500: for C : 0.689

Power of performed test with alpha = 0.0500: for R : 0.0500

Least square means for T :

Group Mean

1.000 39.867

2.000 53050.533

Std Err of LS Mean = 10685.669

Least square means for C :

Group Mean

1.000 119.333

2.000 40020.000

3.000 88510.000

4.000 3171.667

5.000 905.000

Std Err of LS Mean = 16895.527

Least square means for R :

Group Mean

1.000 20153.600

2.000 20680.000

3.000 38802.000

Std Err of LS Mean = 13087.219

Three Way Analysis of Variance

Monday, March 19, 2012, 16:14:50

Data source: ANOVA BSFL gut Plate count medium in ANOVA BSFL gut Plate count medium

Balanced Design (No Interactions)

Dependent Variable: Col 4

Normality Test: Failed (P < 0.050)

Equal Variance Test: Passed (P = 1.000)

| Source of Variation | DF | SS | MS | F | P |
|---------------------|----|------------------|------------------|---------|--------|
| Col 1 | 1 | 795360395700.300 | 795360395700.300 | 179.832 | <0.001 |
| Col 2 | 4 | 1.220E+012 30 | 5092657541.967 | 68.982 | <0.001 |
| Col 3 | 2 | 15524723293.267 | 7762361646.633 | 1.755 | 0.233 |
| Residual | 8 | 35382363173.067 | 4422795396.633 | | |
| Total | 23 | 3.327E+012 | 14719680659.151 | | |

The difference in the mean values among the different levels of Col 1 are greater than would be expected by chance after allowing for the effects of differences in Col 2 and Col 3. There is a statistically significant difference (P = <0.001). To isolate which group(s) differ from the others use a multiple comparison procedure.

The difference in the mean values among the different levels of Col 2 are greater than would be expected by chance after allowing for the effects of differences in Col 1 and Col 3. There is a statistically significant difference (P = <0.001). To isolate which group(s) differ from the others use a multiple comparison procedure.

The difference in the mean values among the different levels of Col 3 are not great enough to exclude the possibility that the difference is just due to random sampling variability after allowing for the effects of differences in Col 1 and Col 2. There is not a statistically significant difference (P = 0.233).

All Pairwise Multiple Comparison Procedures (Holm-Sidak method):

Overall significance level = 0.05

Comparisons for factor: **Col 1**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|--------|--------------|----------------|--------------|
| 2.000 vs. 1.000 | 325650.200 | 13.410 | <0.001 | 0.050 | Yes |

Comparisons for factor: **Col 2**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|--------|--------------|----------------|--------------|
| 2.000 vs. 5.000 | 483278.833 | 12.587 | <0.001 | 0.005 | Yes |

| | | | | | |
|-----------------|------------|--------|--------|-------|-----|
| 2.000 vs. 1.000 | 479883.333 | 12.498 | <0.001 | 0.006 | Yes |
| 2.000 vs. 4.000 | 474980.000 | 12.371 | <0.001 | 0.006 | Yes |
| 3.000 vs. 5.000 | 320612.167 | 8.350 | <0.001 | 0.007 | Yes |
| 3.000 vs. 1.000 | 317216.667 | 8.262 | <0.001 | 0.009 | Yes |
| 3.000 vs. 4.000 | 312313.333 | 8.134 | <0.001 | 0.010 | Yes |
| 2.000 vs. 3.000 | 162666.667 | 4.237 | 0.003 | 0.013 | Yes |
| 4.000 vs. 5.000 | 8298.833 | 0.216 | 0.834 | 0.017 | No |
| 4.000 vs. 1.000 | 4903.333 | 0.128 | 0.902 | 0.025 | No |
| 1.000 vs. 5.000 | 3395.500 | 0.0884 | 0.932 | 0.050 | No |

Comparisons for factor: **Col 3**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-------------------|----------------------|----------|---------------------|-----------------------|---------------------|
| 1.000 vs. 3.000 | 55721.700 | 1.874 | 0.098 | 0.017 | No |
| 2.000 vs. 3.000 | 28027.000 | 0.942 | 0.374 | 0.025 | No |
| 1.000 vs. 2.000 | 27694.700 | 0.931 | 0.379 | 0.050 | No |

Power of performed test with alpha = 0.0500: for Col 1 : 1.000

Power of performed test with alpha = 0.0500: for Col 2 : 1.000

Power of performed test with alpha = 0.0500: for Col 3 : 0.134

Least square means for Col 1 :

Group Mean

1.000 2563.133

2.000 328213.333

Std Err of LS Mean = 17171.285

Least square means for Col 2 :

Group Mean

1.000 5666.667

2.000 485550.000

3.000 322883.333

4.000 10570.000

5.000 2271.167

Std Err of LS Mean = 27150.185

Least square means for Col 3 :

Group Mean

1.000 193193.700

2.000 165499.000

3.000 137472.000

Std Err of LS Mean = 21030.443

Three Way Analysis of Variance

Monday, March 19, 2012, 15:50:56

Data source: ANOVA FBL gut *E. coli* in ANOVA FBL gut *E. coli*

Balanced Design (No Interactions)

Dependent Variable: Y

Normality Test: Failed ($P < 0.050$)

Equal Variance Test: Passed ($P = 1.000$)

| Source of Variation | DF | SS | MS | F | P |
|---------------------|----|------------------|------------------|---------|--------|
| T | 1 | 289342177348.033 | 289342177348.033 | 549.679 | <0.001 |
| C | 4 | 865296991978.467 | 216324247994.617 | 410.963 | <0.001 |
| R | 2 | 2437231576.467 | 1218615788.233 | 2.315 | 0.161 |
| Residual | 8 | 4211070551.867 | 526383818.983 | | |
| Total | 29 | 2.057E+012 | 70946674171.551 | | |

The difference in the mean values among the different levels of T are greater than would be expected by chance after allowing for the effects of differences in C and R. There is a statistically significant difference ($P = <0.001$). To isolate which group(s) differ from the others use a multiple comparison procedure.

The difference in the mean values among the different levels of C are greater than would be expected by chance after allowing for the effects of differences in T and R. There is a statistically significant difference ($P = <0.001$). To isolate which group(s) differ from the others use a multiple comparison procedure.

The difference in the mean values among the different levels of R are not great enough to exclude the possibility that the difference is just due to random sampling variability after allowing for the effects of differences in T and C. There is not a statistically significant difference ($P = 0.161$).

All Pairwise Multiple Comparison Procedures (Holm-Sidak method):

Overall significance level = 0.05

Comparisons for factor: **T**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|--------|--------------|----------------|--------------|
| 2.000 vs. 1.000 | 196415.267 | 23.445 | <0.001 | 0.050 | Yes |

Comparisons for factor: **C**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|--------|--------------|----------------|--------------|
| 3.000 vs. 5.000 | 441169.000 | 33.305 | <0.001 | 0.005 | Yes |

| | | | | | |
|-----------------|------------|--------|--------|-------|-----|
| 3.000 vs. 4.000 | 430199.333 | 32.477 | <0.001 | 0.006 | Yes |
| 3.000 vs. 1.000 | 418376.167 | 31.585 | <0.001 | 0.006 | Yes |
| 3.000 vs. 2.000 | 404107.833 | 30.508 | <0.001 | 0.007 | Yes |
| 2.000 vs. 5.000 | 37061.167 | 2.798 | 0.023 | 0.009 | No |
| 2.000 vs. 4.000 | 26091.500 | 1.970 | 0.084 | 0.010 | No |
| 1.000 vs. 5.000 | 22792.833 | 1.721 | 0.124 | 0.013 | No |
| 2.000 vs. 1.000 | 14268.333 | 1.077 | 0.313 | 0.017 | No |
| 1.000 vs. 4.000 | 11823.167 | 0.893 | 0.398 | 0.025 | No |
| 4.000 vs. 5.000 | 10969.667 | 0.828 | 0.432 | 0.050 | No |

Comparisons for factor: **R**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-------------------|----------------------|----------|---------------------|-----------------------|---------------------|
| 2.000 vs. 1.000 | 21282.200 | 2.074 | 0.072 | 0.017 | No |
| 3.000 vs. 1.000 | 15728.900 | 1.533 | 0.164 | 0.025 | No |
| 2.000 vs. 3.000 | 5553.300 | 0.541 | 0.603 | 0.050 | No |

Power of performed test with alpha = 0.0500: for T : 1.000

Power of performed test with alpha = 0.0500: for C : 1.000

Power of performed test with alpha = 0.0500: for R : 0.204

Least square means for T :

Group Mean

1.000 4731.400

2.000 201146.667

Std Err of LS Mean = 5923.872

Least square means for C :

Group Mean

1.000 23333.333

2.000 37601.667

3.000 441709.500

4.000 11510.167

5.000 540.500

Std Err of LS Mean = 9366.463

Least square means for R :

Group Mean

1.000 90602.000

2.000 111884.200

3.000 106330.900

Std Err of LS Mean = 7255.231

Three Way Analysis of Variance

Monday, March 19, 2012, 15:59:35

Data source: ANOVA FBL gut *Salmonella* in ANOVA FBL gut *Salmonella*

Balanced Design (No Interactions)

Dependent Variable: Y

Normality Test: Failed (P < 0.050)

Equal Variance Test: Passed (P = 1.000)

| Source of Variation | DF | SS | MS | F | P |
|---------------------|----|-----------------|----------------|--------|--------|
| T | 1 | 5083997648.133 | 5083997648.133 | 97.628 | <0.001 |
| C | 4 | 3124422187.200 | 781105546.800 | 15.000 | <0.001 |
| R | 2 | 179414888.467 | 89707444.233 | 1.723 | 0.239 |
| Residual | 8 | 416599594.200 | 52074949.275 | | |
| Total | 29 | 13693679987.867 | 472195861.651 | | |

The difference in the mean values among the different levels of T are greater than would be expected by chance after allowing for the effects of differences in C and R. There is a statistically significant difference (P = <0.001). To isolate which group(s) differ from the others use a multiple comparison procedure.

The difference in the mean values among the different levels of C are greater than would be expected by chance after allowing for the effects of differences in T and R. There is a statistically significant difference (P = <0.001). To isolate which group(s) differ from the others use a multiple comparison procedure.

The difference in the mean values among the different levels of R are not great enough to exclude the possibility that the difference is just due to random sampling variability after allowing for the effects of differences in T and C. There is not a statistically significant difference (P = 0.239).

All Pairwise Multiple Comparison Procedures (Holm-Sidak method):

Overall significance level = 0.05

Comparisons for factor: **T**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|-------|--------------|----------------|--------------|
| 2.000 vs. 1.000 | 26035.867 | 9.881 | <0.001 | 0.050 | Yes |

Comparisons for factor: **C**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|-------|--------------|----------------|--------------|
| 3.000 vs. 4.000 | 30753.333 | 7.381 | <0.001 | 0.005 | Yes |

| | | | | | |
|-----------------|-----------|-------|--------|-------|-----|
| 1.000 vs. 4.000 | 21574.000 | 5.178 | <0.001 | 0.006 | Yes |
| 3.000 vs. 5.000 | 18341.000 | 4.402 | 0.002 | 0.006 | Yes |
| 2.000 vs. 4.000 | 18207.333 | 4.370 | 0.002 | 0.007 | Yes |
| 3.000 vs. 2.000 | 12546.000 | 3.011 | 0.017 | 0.009 | No |
| 5.000 vs. 4.000 | 12412.333 | 2.979 | 0.018 | 0.010 | No |
| 3.000 vs. 1.000 | 9179.333 | 2.203 | 0.059 | 0.013 | No |
| 1.000 vs. 5.000 | 9161.667 | 2.199 | 0.059 | 0.017 | No |
| 2.000 vs. 5.000 | 5795.000 | 1.391 | 0.202 | 0.025 | No |
| 1.000 vs. 2.000 | 3366.667 | 0.808 | 0.442 | 0.050 | No |

Comparisons for factor: **R**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-------------------|----------------------|----------|---------------------|-----------------------|---------------------|
| 1.000 vs. 2.000 | 5439.400 | 1.685 | 0.130 | 0.017 | No |
| 1.000 vs. 3.000 | 4892.700 | 1.516 | 0.168 | 0.025 | No |
| 3.000 vs. 2.000 | 546.700 | 0.169 | 0.870 | 0.050 | No |

Power of performed test with alpha = 0.0500: for T : 1.000

Power of performed test with alpha = 0.0500: for C : 0.997

Power of performed test with alpha = 0.0500: for R : 0.130

Least square means for T :

Group Mean

1.000 4664.133

2.000 30700.000

Std Err of LS Mean = 1863.240

Least square means for C :

Group Mean

1.000 22666.667

2.000 19300.000

3.000 31846.000

4.000 1092.667

5.000 13505.000

Std Err of LS Mean = 2946.041

Least square means for R :

Group Mean

1.000 21126.100

2.000 15686.700

3.000 16233.400

Std Err of LS Mean = 2281.994

Three Way Analysis of Variance

Monday, March 19, 2012, 14:42:25

Data source: ANOVA FBL gut Plate countmedium in ANOVA FBL gut Plate count medium

Balanced Design (No Interactions)

Dependent Variable: Y

Normality Test: Failed (P < 0.050)

Equal Variance Test: Passed (P = 1.000)

| Source of Variation | DF | SS | MS | F | P |
|---------------------|----|------------------|-----------------|--------------|--------|
| T | 1 | 143431898781.633 | 143431898 | 781.6106.291 | <0.001 |
| C | 4 | 1.541E+012 | 385284530525.8 | 285.517 | <0.001 |
| R | 2 | 2612610782. | 6001306305391.3 | 0.968 | 0.420 |
| Residual | 8 | 10795425067.067 | 1349428133.383 | | |
| Total | 29 | 1.930E+0126 | 6549719835.197 | | |

The difference in the mean values among the different levels of T are greater than would be expected by chance after allowing for the effects of differences in C and R. There is a statistically significant difference (P = <0.001). To isolate which group(s) differ from the others use a multiple comparison procedure.

The difference in the mean values among the different levels of C are greater than would be expected by chance after allowing for the effects of differences in T and R. There is a statistically significant difference (P = <0.001). To isolate which group(s) differ from the others use a multiple comparison procedure.

The difference in the mean values among the different levels of R are not great enough to exclude the possibility that the difference is just due to random sampling variability after allowing for the effects of differences in T and C. There is not a statistically significant difference (P = 0.420).

All Pairwise Multiple Comparison Procedures (Holm-Sidak method):

Overall significance level = 0.05

Comparisons for factor: **T**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|--------|--------------|----------------|--------------|
| 2.000 vs. 1.000 | 138290.467 | 10.310 | <0.001 | 0.050 | Yes |

Comparisons for factor: **C**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-----------------|---------------|--------|--------------|----------------|--------------|
| 1.000 vs. 4.000 | 604935.000 | 28.523 | <0.001 | 0.005 | Yes |

| | | | | | |
|-----------------|------------|--------|--------|-------|-----|
| 1.000 vs. 5.000 | 598791.167 | 28.233 | <0.001 | 0.006 | Yes |
| 1.000 vs. 3.000 | 522166.667 | 24.620 | <0.001 | 0.006 | Yes |
| 1.000 vs. 2.000 | 357500.000 | 16.856 | <0.001 | 0.007 | Yes |
| 2.000 vs. 4.000 | 247435.000 | 11.667 | <0.001 | 0.009 | Yes |
| 2.000 vs. 5.000 | 241291.167 | 11.377 | <0.001 | 0.010 | Yes |
| 2.000 vs. 3.000 | 164666.667 | 7.764 | <0.001 | 0.013 | Yes |
| 3.000 vs. 4.000 | 82768.333 | 3.903 | 0.005 | 0.017 | Yes |
| 3.000 vs. 5.000 | 76624.500 | 3.613 | 0.007 | 0.025 | Yes |
| 5.000 vs. 4.000 | 6143.833 | 0.290 | 0.779 | 0.050 | No |

Comparisons for factor: **R**

| Comparison | Diff of Means | t | Unadjusted P | Critical Level | Significant? |
|-------------------|----------------------|----------|---------------------|-----------------------|---------------------|
| 1.000 vs. 3.000 | 22800.700 | 1.388 | 0.203 | 0.017 | No |
| 2.000 vs. 3.000 | 12810.200 | 0.780 | 0.458 | 0.025 | No |
| 1.000 vs. 2.000 | 9990.500 | 0.608 | 0.560 | 0.050 | No |

Power of performed test with alpha = 0.0500: for T : 1.000
Power of performed test with alpha = 0.0500: for C : 1.000
Power of performed test with alpha = 0.0500: for R : 0.0500

Least square means for T :

Group Mean
1.000 130842.867
2.000 269133.333
Std Err of LS Mean = 9484.823

Least square means for C :

Group Mean
1.000 616666.667
2.000 259166.667
3.000 94500.000
4.000 11731.667
5.000 17875.500
Std Err of LS Mean = 14996.823

Least square means for R :

Group Mean
1.000 210918.500
2.000 200928.000
3.000 188117.800
Std Err of LS Mean = 11616.489

Appendix Four: Turbo mass spectrometer GC/MS




Fig 4 1: Turbo mass spectrometer detector used in GC/MS; a) Outer view b) Inside view




Fig 4 2: Capillary column (Zebran ZB-S, 30 m length x 0.25 mm Diameter x 0.25 μ m phase thickness)



Interactions between invertebrates and pathogenic and non-pathogenic bacteria

S. Jaber and M. Wainwright

Department of Molecular Biology and Biotechnology, Firth Court, Western Bank, University of Sheffield, S10 2TN, England
mbp008sma@sheffield.ac.uk



Introduction

The risks of canine waste accumulation in urban environment and agro-ecosystem is an ever-growing issue, pathogenic bacteria such as *Escherichia coli* and *Salmonella* Spp. have been isolated from animals faeces including canine faeces. Invertebrates such as slugs and snails play an important role in the food chain and can act as agricultural pests. (Emma et al, 2006) observed that the Great Gray Slug, *Limax maximus* and the Yellow Slug, *Limax flavus* can carry *E. coli* O157 from animals manure on their bodies surface and internally as intermediate hosts vectors. Recycling is a sustainable strategy for disposing of animals waste, and composting can be an important part of a recycling programme. The microorganisms involved oxidize carbon as an energy source for growth and ingest nitrogen for protein synthesis (Taylor 2004). Consequently, the right carbon to nitrogen ratio in composting systems is required for efficient decomposition of wet dog manure contains 0.7% nitrogen (N), and 0.25% phosphate (P2O4) (compared to wet cattle manure, dog waste which contains 40% more nitrogen, the same amount of phosphate (Hall and Schulte 1979). Insects such as Black soldier fly larvae have been used in bioremediation to breakdown and recycle animals faeces which lead to a reduction in bacterial number in animal's faeces which may be used for biodiesel production and animal food-stock. This study aims to investigate the problems resulting from dog faeces accumulation in the environment and to find scientific solutions for utilizing and converting dogs waste to a safer material which should be more compatible with environmental aspects.

Results

1. Microorganisms isolated from canine faeces, slugs, snails and lettuce

The results (Figure 1.A) show seven differently coloured bacterial colonies which were isolated from dog faeces grown on CHROMagar media. Six of them were transmitted to the inside (gut) of slugs fed on dog faeces (Figure 1.B); as well, *Staphylococcus lentus* grow in the slime on the surface of slugs. However, in the contaminated lettuce samples, four colonies, different from *Enterococci*, were isolated. The colonies were initially identified using colour reference as: *S. lentus*, *Deftia* sp., *E. coli* and *S. enterica* in addition to *Enterococci* spp which were isolated everywhere (Table. 1). Further identification of bacterial species using 16S rRNA procedure has been investigated. The isolations JS1, JS2, JS3, JS4 and JS5 (Table. 2) were subjected to the extraction of genomic DNA (Figure 3.4) which were isolated from; outside control slugs (Metallic blue), inside slugs fed on faeces (Beige), outside slugs fed on faeces (Beige), outside slugs fed on faeces (Cream) and contaminated salad(Beige) respectively. In addition to three isolations MR1, MR2 and MR3 (Table. 2) which were isolated from the Inside of slugs




Figure 1(A)Bacteria isolated from dog faeces grown on Chromagar medium. 1. *S. lentus*; 2. *Enterococcus* spp; 3. *Proteus* spp; 4. *S. enterica*; 5. *E. coli*; 6. *Deftia* sp; 7. *Acinetobacter* sp. B) Bacteria grown on Petri dish isolated from slugs gut following 1. *S. lentus*; 2. *S. enterica*; 3. *Enterococcus* spp; 4. *E. coli*; 5. *Proteus* sp; 6. *Deftia* sp; C) *E. coli* and D) *S. enterica* are bacteria grown on selective media isolated from public park and E) *Aspergillus niger* and A. *flavus*

Table 1. Different colours of bacteria species grown on CHROMagar medium

| Bacterial | Colony color on CHROMagar medium | Presence of bacterial species in different samples | Slugs | Slugs | Slugs | Slugs | Slugs | Slugs | Slugs | Slugs | Slugs | Slugs |
|-----------------------|----------------------------------|--|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| | | | | | | | | | | | | |
| Acinetobacter sp | Blue transparent | - | - | - | - | - | - | - | - | - | - | - |
| Acinetobacter | Metallic Blue | - | - | - | - | - | - | - | - | - | - | - |
| Staphylococcus aureus | Metallic Blue | - | - | - | - | - | - | - | - | - | - | - |
| Acinetobacter | Beige | - | - | - | - | - | - | - | - | - | - | - |
| Enterococcus sp | Blue | - | - | - | - | - | - | - | - | - | - | - |
| Deftia sp | Red | - | - | - | - | - | - | - | - | - | - | - |
| Proteus sp | Red with black halo | - | - | - | - | - | - | - | - | - | - | - |
| E. coli | Pink and | - | - | - | - | - | - | - | - | - | - | - |
| Salmonella sp | Orange | - | - | - | - | - | - | - | - | - | - | - |

2. Identification of bacteria and fungi transmitted from canine faeces to slugs and snails to lettuce



Figure 2. PCR amplification products of four bacterial species analyzed by electrophoresis in agarose gel lanes represent: (lane 1); hyper ladder;(lane JS1) outside control slugs (MB); (lane JS 2) inside slugs fed on faeces (B); (lane JS3) outside slugs fed on faeces (C); (lane JS 4) outside slugs fed on faeces (C); (lane 5) Contaminated salad



Figure 3. PCR amplification products of bacterial species detecting bacteria test analyzed by electrophoresis in agarose gel lanes represent: (lane 1); hyper ladder;(lane MR1) inside slugs fed on faeces (MB); (lane MR 2) inside slugs fed on faeces (C); (lane MR3) Contaminated salad




Figure 4. Polymerase chain reaction (PCR) detection of Mycoplasma species in Dog faeces and lettuce samples. EZ-PCR, detecting Mycoplasma test analyzed by electrophoresis in 2% agarose gel; the lanes represent: (lane L); hyper ladder; (lane KG and K6) dog faeces samples.




Figure 5. PCR-16S rRNA amplification products of fungal species detecting fungi test analyzed by electrophoresis in 2% agarose gel; the lanes represent: (lane L); hyper ladder; (lane K6 and K5) dog faeces samples.

Table 2. Bacterial identification using gDNA-16S rRNA

| S.No. | Strains | Species | % |
|-------|--------------------------------|--------------------------------|-----|
| JS1 | Outside control slug (MR) | <i>Acinetobacter baumannii</i> | 99% |
| JS2 | Inside slug fed on faeces (B) | <i>Acinetobacter</i> sp. | 98% |
| JS3 | Outside slug fed on faeces (B) | <i>Enterococcus</i> sp. | 99% |
| JS4 | Outside slug fed on faeces (C) | <i>Acinetobacter</i> sp. | 98% |
| JS5 | Contaminated salad (B) | <i>Deftia</i> sp. | 99% |
| MR1 | Inside slug fed on faeces (MB) | <i>Staphylococcus aureus</i> | 99% |
| MR2 | Inside slug fed on faeces (C) | <i>Acinetobacter</i> sp. | 99% |
| MR3 | Contaminated salad (FB) | <i>E. coli</i> | 99% |

Table 3. The presence of Mycoplasma in various samples

| Sample | Symbol | Reaction |
|---------------------------|--------|----------|
| Earth worm (Whole body) | RM1 | Positive |
| Dog faeces | SM1 | Positive |
| Contaminated lettuce | SM2 | Positive |
| Control lettuce | SM3 | Negative |
| Outside control slug | AZ1 | Negative |
| Outside contaminated slug | AZ2 | Positive |
| Outside contaminated slug | AZ3 | Negative |
| Outside contaminated slug | AZ4 | Negative |
| Inside control slug | AZ5 | Positive |
| Inside contaminated slug | AZ6 | Positive |
| Inside control slug | AZ7 | Negative |
| Inside contaminated slug | AZ8 | Positive |

3. Summary of the canine waste accumulation problems

Dog faeces can carry species of bacteria, mycoplasma and pathogenic fungi

Insert BSF and FB larvae as a solution suggested for bioremediation of dog faeces

contaminated slugs and snails can carry species of bacteria and mycoplasma

Species of bacteria and mycoplasma can be transmitted to lettuce through

4. The effect of BSF and FB larvae activity on on E. coli and Salmonella enterica populations




Figure 6. Influence of BSFL on bacterial activity on dog faeces.




Figure 7. Influence of FBFL on bacterial activity on dog faeces.



Figure 8. The influence of BSFL gut on bacterial in dog faeces



Figure 9. The influence of FBFL gut on bacterial in dog faeces




Figure 10. The final compound of untreated dog faeces after 28 days



Figure 11. The final compound of treated dog faeces after 28 days




Figure 12. Illustrates: a) the lettuce seeds planted in different concentration of treated dog faeces with BSFL No. 1, control dog faeces 2, 3 and 4 are 75, 50 and 25% of dog faeces concentration and 5 is a control compost. From 6 to 10 are the lettuce seeds planted at the same concentration of treated dog faeces with FBFL No. 1, control dog faeces 2, 3 and 4 were 75, 50 and 25% of dog faeces and 5 is a control compost. From 6 to 10 are the lettuce seeds planted at the same concentration

5. SEM and TEM studies on the morphological effects of Hemolymph and Whole body secretions of BSF and FB larvae on MRSA, E. coli, S. enterica




Figure 13. The effect of hemolymph and whole body secretions of BSF and FB larvae on the morphological of MRSA (a-e), *E. coli* (f-i), internal morphology different bacterial species (k-o). Z and y the effect the secretion of larvae on *E. coli* and MRSA growth




Figure 14. The amount of lipids in dry weight of 1g of dog faeces and BSF and FB larvae




Figure 15. Use the GC/MS to identify the extracted fatty acids from dog's faeces, BSF and FB Larvae

15. Use the GC/MS to identify the extracted fatty acids from dog's faeces, BSF and FB Larvae

7. Dogs waste from biomass to biofuel

The results of this study show that seven species of bacteria were found in dog faeces. Six out of these seven species were confirmed for being carried by slugs and snails, when obligatory fed on dog's faeces. Slugs transmitted four bacterial species i.e.: *E. coli* sp., *Salmonella enterica*, *Staphylococcus lentus* and *Deftia* sp to fresh lettuce. Our study suggests a range of different solutions to convert the dog faeces from a poisonous substance to a less offensive material using insect larvae. Bioremediation of dog faeces was successful in converting dog waste to modified compost for agricultural purpose. Lipids of poeul use as as befouts were also extracted from dog faeces treated with larvae.

8. Acknowledgments

I thank my supervisor Prof. M. Wainwright for the valuable guidance and advice. I also thank my advisors Dr. D. J. Gilmour and Dr L. Partridge for their assistance. I am grateful to Mr S. J. Thorpe from chemistry dept for his help in GC/MS, and Mr C.J. Hill in SEM/TEM.

9. Conclusion

The results of this study show that seven species of bacteria were found in dog faeces. Six out of these seven species were confirmed for being carried by slugs and snails, when obligatory fed on dog's faeces. Slugs transmitted four bacterial species i.e.: *E. coli* sp., *Salmonella enterica*, *Staphylococcus lentus* and *Deftia* sp to fresh lettuce. Our study suggests a range of different solutions to convert the dog faeces from a poisonous substance to a less offensive material using insect larvae. Bioremediation of dog faeces was successful in converting dog waste to modified compost for agricultural purpose. Lipids of poeul use as as befouts were also extracted from dog faeces treated with larvae.

10. References

Emma, L.S., M. Meece, I.D. Cooper, M. J. Wilson and N. J. C. Strachan (2006). Slugs: Potential Novel Vectors of *Escherichia coli* O157. *Applied and Environmental Microbiology*, 72: 144-149

Hall, RE., Schulte, E.F. (1979). Using dog and cat manure onhome gardens. University of Wisconsin, Madison, College of Agriculture and Life Sciences.

Taylor, L. (2004). Poop, Scoop and Compost: Proposal to Explore the Feasibility of Composting Dog Waste in Public Parks. Vancouver, Canada: National Coalition for People and Dogs, Handout, pgs. 1-7

Poster Presentation Relating to Some of the Work Described in this Thesis