

References

- Ahmed, Z., Schuller, A.C., Suhling, K., Tregidgo, C., Ladbury, J.E., 2008. Extracellular point mutations in FGFR2 elicit unexpected changes in intracellular signalling. *The Biochemical journal* 413, 37-49.
- Ahrens, K., Schlosser, G., 2005. Tissues and signals involved in the induction of placodal Six1 expression in *Xenopus laevis*. *Developmental biology* 288, 40-59.
- Aikawa, T., Segre, G.V., Lee, K., 2001. Fibroblast growth factor inhibits chondrocytic growth through induction of p21 and subsequent inactivation of cyclin E-Cdk2. *The Journal of biological chemistry* 276, 29347-29352.
- Alvarez, Y., Alonso, M.T., Vendrell, V., Zelarayan, L.C., Chamero, P., Theil, T., Bosl, M.R., Kato, S., Maconochie, M., Riethmacher, D., Schimmang, T., 2003. Requirements for FGF3 and FGF10 during inner ear formation. *Development* 130, 6329-6338.
- Avery, S., Zafarana, G., Gokhale, P.J., Andrews, P.W., 2010. The role of SMAD4 in human embryonic stem cell self-renewal and stem cell fate. *Stem cells* 28, 863-873.
- Avila, M.A., Varela-Nieto, I., Romero, G., Mato, J.M., Giraldez, F., Van De Water, T.R., Represa, J., 1993. Brain-derived neurotrophic factor and neurotrophin-3 support the survival and neuritogenesis response of developing cochleovestibular ganglion neurons. *Developmental biology* 159, 266-275.
- Bailey, A.P., Bhattacharyya, S., Bronner-Fraser, M., Streit, A., 2006. Lens specification is the ground state of all sensory placodes, from which FGF promotes olfactory identity. *Developmental cell* 11, 505-517.
- Barald, K.F., Kelley, M.W., 2004. From placode to polarization: new tunes in inner ear development. *Development* 131, 4119-4130.
- Bermingham, N.A., Hassan, B.A., Price, S.D., Vollrath, M.A., Ben-Arie, N., Eatock, R.A., Bellen, H.J., Lysakowski, A., Zoghbi, H.Y., 1999. Math1: an essential gene for the generation of inner ear hair cells. *Science* 284, 1837-1841.
- Besseyrias, V., Fiorini, E., Strobl, L.J., Zimmer-Strobl, U., Dumortier, A., Koch, U., Arcangeli, M.L., Ezine, S., Macdonald, H.R., Radtke, F., 2007. Hierarchy of Notch-Delta interactions

promoting T cell lineage commitment and maturation. *The Journal of experimental medicine* 204, 331-343.

Bhattacharyya, S., Bailey, A.P., Bronner-Fraser, M., Streit, A., 2004. Segregation of lens and olfactory precursors from a common territory: cell sorting and reciprocity of *Dlx5* and *Pax6* expression. *Developmental biology* 271, 403-414.

Bichler, E., Spöndlin, H., Rauegger, H., 1983. Degeneration of cochlear neurons after amikacin intoxication in the rat. *Archives of oto-rhino-laryngology* 237, 201-208.

Blauwkamp, T.A., Nigam, S., Ardehali, R., Weissman, I.L., Nusse, R., 2012. Endogenous Wnt signalling in human embryonic stem cells generates an equilibrium of distinct lineage-specified progenitors. *Nature communications* 3, 1070.

Boddy, S.L., Chen, W., Romero-Guevara, R., Kottam, L., Bellantuono, I., Rivolta, M.N., 2012. Inner ear progenitor cells can be generated in vitro from human bone marrow mesenchymal stem cells. *Regenerative medicine* 7, 757-767.

Bouchard, M., de Caprona, D., Busslinger, M., Xu, P., Fritzsche, B., 2010. *Pax2* and *Pax8* cooperate in mouse inner ear morphogenesis and innervation. *BMC developmental biology* 10, 89.

Bouchard, M., Souabni, A., Busslinger, M., 2004. Tissue-specific expression of cre recombinase from the *Pax8* locus. *Genesis* 38, 105-109.

Brons, I.G., Smithers, L.E., Trotter, M.W., Rugg-Gunn, P., Sun, B., Chuva de Sousa Lopes, S.M., Howlett, S.K., Clarkson, A., Ahrlund-Richter, L., Pedersen, R.A., Vallier, L., 2007. Derivation of pluripotent epiblast stem cells from mammalian embryos. *Nature* 448, 191-195.

Brugmann, S.A., Pandur, P.D., Kenyon, K.L., Pignoni, F., Moody, S.A., 2004. *Six1* promotes a placodal fate within the lateral neurogenic ectoderm by functioning as both a transcriptional activator and repressor. *Development* 131, 5871-5881.

Brummelkamp, T.R., Bernards, R., Agami, R., 2002. Stable suppression of tumorigenicity by virus-mediated RNA interference. *Cancer cell* 2, 243-247.

Burton, Q., Cole, L.K., Mulheisen, M., Chang, W., Wu, D.K., 2004. The role of *Pax2* in mouse inner ear development. *Developmental biology* 272, 161-175.

Caplen, N.J., Parrish, S., Imani, F., Fire, A., Morgan, R.A., 2001. Specific inhibition of gene expression by small double-stranded RNAs in invertebrate and vertebrate systems. *Proceedings of the National Academy of Sciences of the United States of America* 98, 9742-9747.

Cappellen, D., De Oliveira, C., Ricol, D., de Medina, S., Bourdin, J., Sastre-Garau, X., Chopin, D., Thiery, J.P., Radvanyi, F., 1999. Frequent activating mutations of FGFR3 in human bladder and cervix carcinomas. *Nature genetics* 23, 18-20.

Carstens, R.P., Eaton, J.V., Krigman, H.R., Walther, P.J., Garcia-Blanco, M.A., 1997. Alternative splicing of fibroblast growth factor receptor 2 (FGF-R2) in human prostate cancer. *Oncogene* 15, 3059-3065.

Celli, G., LaRochelle, W.J., Mackem, S., Sharp, R., Merlino, G., 1998. Soluble dominant-negative receptor uncovers essential roles for fibroblast growth factors in multi-organ induction and patterning. *The EMBO journal* 17, 1642-1655.

Chambers, S.M., Fasano, C.A., Papapetrou, E.P., Tomishima, M., Sadelain, M., Studer, L., 2009. Highly efficient neural conversion of human ES and iPS cells by dual inhibition of SMAD signaling. *Nature biotechnology* 27, 275-280.

Chellaiah, A., Yuan, W., Chellaiah, M., Ornitz, D.M., 1999. Mapping ligand binding domains in chimeric fibroblast growth factor receptor molecules. Multiple regions determine ligand binding specificity. *The Journal of biological chemistry* 274, 34785-34794.

Chellaiah, A.T., McEwen, D.G., Werner, S., Xu, J., Ornitz, D.M., 1994. Fibroblast growth factor receptor (FGFR) 3. Alternative splicing in immunoglobulin-like domain III creates a receptor highly specific for acidic FGF/FGF-1. *The Journal of biological chemistry* 269, 11620-11627.

Chen, C.W., Liu, C.S., Chiu, I.M., Shen, S.C., Pan, H.C., Lee, K.H., Lin, S.Z., Su, H.L., 2010. The signals of FGFs on the neurogenesis of embryonic stem cells. *Journal of biomedical science* 17, 33.

Chen, G., Gulbranson, D.R., Yu, P., Hou, Z., Thomson, J.A., 2012. Thermal stability of fibroblast growth factor protein is a determinant factor in regulating self-renewal, differentiation, and reprogramming in human pluripotent stem cells. *Stem cells* 30, 623-630.

Chen, W., Jongkamonwiwat, N., Abbas, L., Eshtan, S.J., Johnson, S.L., Kuhn, S., Milo, M., Thurlow, J.K., Andrews, P.W., Marcotti, W., Moore, H.D., Rivolta, M.N., 2012. Restoration of auditory evoked responses by human ES-cell-derived otic progenitors. *Nature* 490, 278-282.

Chesi, M., Brents, L.A., Ely, S.A., Bais, C., Robbiani, D.F., Mesri, E.A., Kuehl, W.M., Bergsagel, P.L., 2001. Activated fibroblast growth factor receptor 3 is an oncogene that contributes to tumor progression in multiple myeloma. *Blood* 97, 729-736.

Chi, N., Epstein, J.A., 2002. Getting your Pax straight: Pax proteins in development and disease. *Trends in genetics : TIG* 18, 41-47.

Chikar, J.A., Colesa, D.J., Swiderski, D.L., Di Polo, A., Raphael, Y., Pflugst, B.E., 2008. Over-expression of BDNF by adenovirus with concurrent electrical stimulation improves cochlear implant thresholds and survival of auditory neurons. *Hearing research* 245, 24-34.

Cohen, M.A., Itsykson, P., Reubinoff, B.E., 2010. The role of FGF-signaling in early neural specification of human embryonic stem cells. *Developmental biology* 340, 450-458.

Colvin, J.S., Bohne, B.A., Harding, G.W., McEwen, D.G., Ornitz, D.M., 1996. Skeletal overgrowth and deafness in mice lacking fibroblast growth factor receptor 3. *Nature genetics* 12, 390-397.

Corrales, C.E., Pan, L., Li, H., Liberman, M.C., Heller, S., Edge, A.S., 2006. Engraftment and differentiation of embryonic stem cell-derived neural progenitor cells in the cochlear nerve trunk: growth of processes into the organ of Corti. *Journal of neurobiology* 66, 1489-1500.

Coumoul, X., Shukla, V., Li, C., Wang, R.H., Deng, C.X., 2005. Conditional knockdown of Fgfr2 in mice using Cre-LoxP induced RNA interference. *Nucleic acids research* 33, e102.

Cunha, G.R., Hayward, S.W., Wang, Y.Z., Ricke, W.A., 2003. Role of the stromal microenvironment in carcinogenesis of the prostate. *International journal of cancer. Journal international du cancer* 107, 1-10.

Darr, H., Mayshar, Y., Benvenisty, N., 2006. Overexpression of NANOG in human ES cells enables feeder-free growth while inducing primitive ectoderm features. *Development* 133, 1193-1201.

De Moerlooze, L., Spencer-Dene, B., Revest, J.M., Hajihosseini, M., Rosewell, I., Dickson, C., 2000. An important role for the IIIb isoform of fibroblast growth factor receptor 2 (FGFR2) in mesenchymal-epithelial signalling during mouse organogenesis. *Development* 127, 483-492.

Delorme, B., Nivet, E., Gaillard, J., Haupl, T., Ringe, J., Deveze, A., Magnan, J., Sohier, J., Khrestchatsky, M., Roman, F.S., Charbord, P., Sensebe, L., Layrolle, P., Feron, F., 2010. The human nose harbors a niche of olfactory ectomesenchymal stem cells displaying neurogenic and osteogenic properties. *Stem cells and development* 19, 853-866.

Ding, V.M., Ling, L., Natarajan, S., Yap, M.G., Cool, S.M., Choo, A.B., 2010. FGF-2 modulates Wnt signaling in undifferentiated hESC and iPS cells through activated PI3-K/GSK3beta signaling. *Journal of cellular physiology* 225, 417-428.

Dodson, H.C., Mohuiddin, A., 2000. Response of spiral ganglion neurones to cochlear hair cell destruction in the guinea pig. *Journal of neurocytology* 29, 525-537.

Doetschman, T.C., Eistetter, H., Katz, M., Schmidt, W., Kemler, R., 1985. The in vitro development of blastocyst-derived embryonic stem cell lines: formation of visceral yolk sac, blood islands and myocardium. *Journal of embryology and experimental morphology* 87, 27-45.

Doyle, K.L., Kazda, A., Hort, Y., McKay, S.M., Oleskevich, S., 2007. Differentiation of adult mouse olfactory precursor cells into hair cells in vitro. *Stem cells* 25, 621-627.

Draper, J.S., Moore, H.D., Ruban, L.N., Gokhale, P.J., Andrews, P.W., 2004. Culture and characterization of human embryonic stem cells. *Stem cells and development* 13, 325-336.

Duran Alonso, M.B., Feijoo-Redondo, A., Conde de Felipe, M., Carnicero, E., Garcia, A.S., Garcia-Sancho, J., Rivolta, M.N., Giraldez, F., Schimmang, T., 2012. Generation of inner ear sensory cells from bone marrow-derived human mesenchymal stem cells. *Regenerative medicine* 7, 769-783.

Dvorak, P., Dvorakova, D., Koskova, S., Vodinska, M., Najvirtova, M., Krekac, D., Hampl, A., 2005. Expression and potential role of fibroblast growth factor 2 and its receptors in human embryonic stem cells. *Stem cells* 23, 1200-1211.

Dvorak, P., Hampl, A., 2005. Basic fibroblast growth factor and its receptors in human embryonic stem cells. *Folia histochemica et cytobiologica / Polish Academy of Sciences, Polish Histochemical and Cytochemical Society* 43, 203-208.

Elbashir, S.M., Harborth, J., Lendeckel, W., Yalcin, A., Weber, K., Tuschl, T., 2001. Duplexes of 21-nucleotide RNAs mediate RNA interference in cultured mammalian cells. *Nature* 411, 494-498.

Elbashir, S.M., Martinez, J., Patkaniowska, A., Lendeckel, W., Tuschl, T., 2001. Functional anatomy of siRNAs for mediating efficient RNAi in *Drosophila melanogaster* embryo lysate. *The EMBO journal* 20, 6877-6888.

Ernfors, P., Duan, M.L., ElShamy, W.M., Canlon, B., 1996. Protection of auditory neurons from aminoglycoside toxicity by neurotrophin-3. *Nature medicine* 2, 463-467.

Ernfors, P., Van De Water, T., Loring, J., Jaenisch, R., 1995. Complementary roles of BDNF and NT-3 in vestibular and auditory development. *Neuron* 14, 1153-1164.

Eswarakumar, V.P., Lax, I., Schlessinger, J., 2005. Cellular signaling by fibroblast growth factor receptors. *Cytokine & growth factor reviews* 16, 139-149.

Fekete, D.M., Wu, D.K., 2002. Revisiting cell fate specification in the inner ear. *Current opinion in neurobiology* 12, 35-42.

Feng, S., Wang, F., Matsubara, A., Kan, M., McKeegan, W.L., 1997. Fibroblast growth factor receptor 2 limits and receptor 1 accelerates tumorigenicity of prostate epithelial cells. *Cancer research* 57, 5369-5378.

Fire, A., Xu, S., Montgomery, M.K., Kostas, S.A., Driver, S.E., Mello, C.C., 1998. Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* 391, 806-811.

Freter, S., Muta, Y., Mak, S.S., Rinkwitz, S., Ladher, R.K., 2008. Progressive restriction of otic fate: the role of FGF and Wnt in resolving inner ear potential. *Development* 135, 3415-3424.

Fritzsche, B., Silos-Santiago, I., Bianchi, L.M., Farinas, I., 1997. The role of neurotrophic factors in regulating the development of inner ear innervation. *Trends in neurosciences* 20, 159-164.

Fritzsche, B., Tessarollo, L., Coppola, E., Reichardt, L.F., 2004. Neurotrophins in the ear: their roles in sensory neuron survival and fiber guidance. *Progress in brain research* 146, 265-278.

Gadue, P., Huber, T.L., Paddison, P.J., Keller, G.M., 2006. Wnt and TGF-beta signaling are required for the induction of an in vitro model of primitive streak formation using embryonic stem cells. *Proceedings of the National Academy of Sciences of the United States of America* 103, 16806-16811.

Garcia, C.M., Huang, J., Madakashira, B.P., Liu, Y., Rajagopal, R., Dattilo, L., Robinson, M.L., Beebe, D.C., 2011. The function of FGF signaling in the lens placode. *Developmental biology* 351, 176-185.

Garcia, M.A., Gil, J., Ventoso, I., Guerra, S., Domingo, E., Rivas, C., Esteban, M., 2006. Impact of protein kinase PKR in cell biology: from antiviral to antiproliferative action. *Microbiology and molecular biology reviews* : MMBR 70, 1032-1060.

Ginis, I., Luo, Y., Miura, T., Thies, S., Brandenberger, R., Gerecht-Nir, S., Amit, M., Hoke, A., Carpenter, M.K., Itskovitz-Eldor, J., Rao, M.S., 2004. Differences between human and mouse embryonic stem cells. *Developmental biology* 269, 360-380.

Glueckert, R., Bitsche, M., Miller, J.M., Zhu, Y., Prieskorn, D.M., Altschuler, R.A., Schrott-Fischer, A., 2008. Deafferentation-associated changes in afferent and efferent processes in the guinea pig cochlea and afferent regeneration with chronic intrascalar brain-derived neurotrophic factor and acidic fibroblast growth factor. *The Journal of comparative neurology* 507, 1602-1621.

Gospodarowicz, D., 1974. Localisation of a fibroblast growth factor and its effect alone and with hydrocortisone on 3T3 cell growth. *Nature* 249, 123-127.

Gossen, M., Bujard, H., 1992. Tight control of gene expression in mammalian cells by tetracycline-responsive promoters. *Proceedings of the National Academy of Sciences of the United States of America* 89, 5547-5551.

Greber, B., Coulon, P., Zhang, M., Moritz, S., Frank, S., Muller-Molina, A.J., Arauzo-Bravo, M.J., Han, D.W., Pape, H.C., Scholer, H.R., 2011. FGF signalling inhibits neural induction in human embryonic stem cells. *The EMBO journal* 30, 4874-4884.

Greber, B., Lehrach, H., Adjaye, J., 2007. Fibroblast growth factor 2 modulates transforming growth factor beta signaling in mouse embryonic fibroblasts and human ESCs (hESCs) to support hESC self-renewal. *Stem cells* 25, 455-464.

Greber, B., Wu, G., Bernemann, C., Joo, J.Y., Han, D.W., Ko, K., Tapia, N., Sabour, D., Sternecker, J., Tesar, P., Scholer, H.R., 2010. Conserved and divergent roles of FGF signaling in mouse epiblast stem cells and human embryonic stem cells. *Cell stem cell* 6, 215-226.

Grose, R., Dickson, C., 2005. Fibroblast growth factor signaling in tumorigenesis. *Cytokine & growth factor reviews* 16, 179-186.

Groves, A.K., Bronner-Fraser, M., 2000. Competence, specification and commitment in otic placode induction. *Development* 127, 3489-3499.

Halder, G., Callaerts, P., Gehring, W.J., 1995. Induction of ectopic eyes by targeted expression of the *eyeless* gene in *Drosophila*. *Science* 267, 1788-1792.

Hammond, S.M., Caudy, A.A., Hannon, G.J., 2001. Post-transcriptional gene silencing by double-stranded RNA. *Nature reviews. Genetics* 2, 110-119.

Hans, S., Christison, J., Liu, D., Westerfield, M., 2007. Fgf-dependent otic induction requires competence provided by *Foxi1* and *Dlx3b*. *BMC developmental biology* 7, 5.

Harper, L.V., Hilton, A.C., Jones, A.F., 2003. RT-PCR for the pseudogene-free amplification of the glyceraldehyde-3-phosphate dehydrogenase gene (*gapd*). *Molecular and cellular probes* 17, 261-265.

Hart, K.C., Robertson, S.C., Kanemitsu, M.Y., Meyer, A.N., Tynan, J.A., Donoghue, D.J., 2000. Transformation and Stat activation by derivatives of FGFR1, FGFR3, and FGFR4. *Oncogene* 19, 3309-3320.

Hatini, V., Ye, X., Balas, G., Lai, E., 1999. Dynamics of placodal lineage development revealed by targeted transgene expression. *Developmental dynamics : an official publication of the American Association of Anatomists* 215, 332-343.

Hayashi, T., Ray, C.A., Younkins, C., Bermingham-McDonogh, O., 2010. Expression patterns of FGF receptors in the developing mammalian cochlea. *Developmental dynamics : an official publication of the American Association of Anatomists* 239, 1019-1026.

Heller, N., Brandli, A.W., 1997. *Xenopus Pax-2* displays multiple splice forms during embryogenesis and pronephric kidney development. *Mechanisms of development* 69, 83-104.

Heller, N., Brandli, A.W., 1999. *Xenopus Pax-2/5/8* orthologues: novel insights into Pax gene evolution and identification of Pax-8 as the earliest marker for otic and pronephric cell lineages. *Developmental genetics* 24, 208-219.

Hildebrand, M.S., Dahl, H.H., Hardman, J., Coleman, B., Shepherd, R.K., de Silva, M.G., 2005. Survival of partially differentiated mouse embryonic stem cells in the scala media of the

guinea pig cochlea. *Journal of the Association for Research in Otolaryngology* : JARO 6, 341-354.

Hu, B.Y., Weick, J.P., Yu, J., Ma, L.X., Zhang, X.Q., Thomson, J.A., Zhang, S.C., 2010. Neural differentiation of human induced pluripotent stem cells follows developmental principles but with variable potency. *Proceedings of the National Academy of Sciences of the United States of America* 107, 4335-4340.

International Stem Cell Initiative, C., Akopian, V., Andrews, P.W., Beil, S., Benvenisty, N., Brehm, J., Christie, M., Ford, A., Fox, V., Gokhale, P.J., Healy, L., Holm, F., Hovatta, O., Knowles, B.B., Ludwig, T.E., McKay, R.D., Miyazaki, T., Nakatsuji, N., Oh, S.K., Pera, M.F., Rossant, J., Stacey, G.N., Suemori, H., 2010. Comparison of defined culture systems for feeder cell free propagation of human embryonic stem cells. *In vitro cellular & developmental biology. Animal* 46, 247-258.

Iyengar, L., Wang, Q., Rasko, J.E., McAvoy, J.W., Lovicu, F.J., 2007. Duration of ERK1/2 phosphorylation induced by FGF or ocular media determines lens cell fate. *Differentiation; research in biological diversity* 75, 662-668.

Izumikawa, M., Minoda, R., Kawamoto, K., Abrashkin, K.A., Swiderski, D.L., Dolan, D.F., Brough, D.E., Raphael, Y., 2005. Auditory hair cell replacement and hearing improvement by Atoh1 gene therapy in deaf mammals. *Nature medicine* 11, 271-276.

Jabs, E.W., Li, X., Scott, A.F., Meyers, G., Chen, W., Eccles, M., Mao, J.I., Charnas, L.R., Jackson, C.E., Jaye, M., 1994. Jackson-Weiss and Crouzon syndromes are allelic with mutations in fibroblast growth factor receptor 2. *Nature genetics* 8, 275-279.

Jacobson, A.G., 1963. The Determination and Positioning of the Nose, Lens and Ear. Iii. Effects of Reversing the Antero-Posterior Axis of Epidermis, Neural Plate and Neural Fold. *The Journal of experimental zoology* 154, 293-303.

Jacobson, A.G., 1966. Inductive processes in embryonic development. *Science* 152, 25-34.

Jeon, S.J., Oshima, K., Heller, S., Edge, A.S., 2007. Bone marrow mesenchymal stem cells are progenitors in vitro for inner ear hair cells. *Molecular and cellular neurosciences* 34, 59-68.

Johnson, D.E., Lu, J., Chen, H., Werner, S., Williams, L.T., 1991. The human fibroblast growth factor receptor genes: a common structural arrangement underlies the mechanisms for generating receptor forms that differ in their third immunoglobulin domain. *Molecular and cellular biology* 11, 4627-4634.

Johnston, C.L., Cox, H.C., Gomm, J.J., Coombes, R.C., 1995. Fibroblast growth factor receptors (FGFRs) localize in different cellular compartments. A splice variant of FGFR-3 localizes to the nucleus. *The Journal of biological chemistry* 270, 30643-30650.

Jouanneau, J., Moens, G., Bourgeois, Y., Poupon, M.F., Thiery, J.P., 1994. A minority of carcinoma cells producing acidic fibroblast growth factor induces a community effect for tumor progression. *Proceedings of the National Academy of Sciences of the United States of America* 91, 286-290.

Jouanneau, J., Plouet, J., Moens, G., Thiery, J.P., 1997. FGF-2 and FGF-1 expressed in rat bladder carcinoma cells have similar angiogenic potential but different tumorigenic properties in vivo. *Oncogene* 14, 671-676.

Kandel, J., Bossy-Wetzler, E., Radvanyi, F., Klagsbrun, M., Folkman, J., Hanahan, D., 1991. Neovascularization is associated with a switch to the export of bFGF in the multistep development of fibrosarcoma. *Cell* 66, 1095-1104.

Kawamoto, K., Ishimoto, S., Minoda, R., Brough, D.E., Raphael, Y., 2003. Math1 gene transfer generates new cochlear hair cells in mature guinea pigs in vivo. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 23, 4395-4400.

Kelley, M.W., 2006. Hair cell development: commitment through differentiation. *Brain research* 1091, 172-185.

Kiefer, P., Dickson, C., 1995. Nucleolar association of fibroblast growth factor 3 via specific sequence motifs has inhibitory effects on cell growth. *Molecular and cellular biology* 15, 4364-4374.

King, F.W., Ritner, C., Liszewski, W., Kwan, H.C., Pedersen, A., Leavitt, A.D., Bernstein, H.S., 2009. Subpopulations of human embryonic stem cells with distinct tissue-specific fates can be selected from pluripotent cultures. *Stem cells and development* 18, 1441-1450.

Kondo, T., Johnson, S.A., Yoder, M.C., Romand, R., Hashino, E., 2005. Sonic hedgehog and retinoic acid synergistically promote sensory fate specification from bone marrow-derived pluripotent stem cells. *Proceedings of the National Academy of Sciences of the United States of America* 102, 4789-4794.

Kunath, T., Saba-El-Leil, M.K., Almousaillekh, M., Wray, J., Meloche, S., Smith, A., 2007. FGF stimulation of the Erk1/2 signalling cascade triggers transition of pluripotent embryonic stem cells from self-renewal to lineage commitment. *Development* 134, 2895-2902.

Kunii, K., Davis, L., Gorenstein, J., Hatch, H., Yashiro, M., Di Bacco, A., Elbi, C., Lutterbach, B., 2008. FGFR2-amplified gastric cancer cell lines require FGFR2 and Erbb3 signaling for growth and survival. *Cancer research* 68, 2340-2348.

Kwabi-Addo, B., Ozen, M., Ittmann, M., 2004. The role of fibroblast growth factors and their receptors in prostate cancer. *Endocrine-related cancer* 11, 709-724.

Ladher, R.K., 2000. Identification of Synergistic Signals Initiating Inner Ear Development. *Science* 290, 1965-1967.

Ladher, R.K., Wright, T.J., Moon, A.M., Mansour, S.L., Schoenwolf, G.C., 2005. FGF8 initiates inner ear induction in chick and mouse. *Genes & development* 19, 603-613.

Lanford, P.J., Lan, Y., Jiang, R., Lindsell, C., Weinmaster, G., Gridley, T., Kelley, M.W., 1999. Notch signalling pathway mediates hair cell development in mammalian cochlea. *Nature genetics* 21, 289-292.

Lawoko-Kerali, G., Rivolta, M.N., Lawlor, P., Cacciabue-Rivolta, D.I., Langton-Hewer, C., van Doorninck, J.H., Holley, M.C., 2004. GATA3 and NeuroD distinguish auditory and vestibular neurons during development of the mammalian inner ear. *Mechanisms of development* 121, 287-299.

Leavitt, A.D., Hamlett, I., 2011. Homologous recombination in human embryonic stem cells: a tool for advancing cell therapy and understanding and treating human disease. *Clinical and translational science* 4, 298-305.

Legeai-Mallet, L., Benoist-Lasselain, C., Munnich, A., Bonaventure, J., 2004. Overexpression of FGFR3, Stat1, Stat5 and p21Cip1 correlates with phenotypic severity and defective chondrocyte differentiation in FGFR3-related chondrodysplasias. *Bone* 34, 26-36.

Levenstein, M.E., Ludwig, T.E., Xu, R.H., Llanas, R.A., VanDenHeuvel-Kramer, K., Manning, D., Thomson, J.A., 2006. Basic fibroblast growth factor support of human embryonic stem cell self-renewal. *Stem cells* 24, 568-574.

Li, C., Yang, Y., Lu, X., Sun, Y., Gu, J., Feng, Y., Jin, Y., 2010. Efficient derivation of Chinese human embryonic stem cell lines from frozen embryos. *In vitro cellular & developmental biology. Animal* 46, 186-191.

Li, H., Liu, H., Heller, S., 2003. Pluripotent stem cells from the adult mouse inner ear. *Nature medicine* 9, 1293-1299.

Li, H., Roblin, G., Liu, H., Heller, S., 2003. Generation of hair cells by stepwise differentiation of embryonic stem cells. *Proceedings of the National Academy of Sciences of the United States of America* 100, 13495-13500.

Li, H.S., Yang, J.M., Jacobson, R.D., Pasko, D., Sundin, O., 1994. Pax-6 is first expressed in a region of ectoderm anterior to the early neural plate: implications for stepwise determination of the lens. *Developmental biology* 162, 181-194.

Li, L.C., Okino, S.T., Zhao, H., Pookot, D., Place, R.F., Urakami, S., Enokida, H., Dahiya, R., 2006. Small dsRNAs induce transcriptional activation in human cells. *Proceedings of the National Academy of Sciences of the United States of America* 103, 17337-17342.

Li, Z., Zhu, Y.X., Plowright, E.E., Bergsagel, P.L., Chesi, M., Patterson, B., Hawley, T.S., Hawley, R.G., Stewart, A.K., 2001. The myeloma-associated oncogene fibroblast growth factor receptor 3 is transforming in hematopoietic cells. *Blood* 97, 2413-2419.

- Liew, C.G., Draper, J.S., Walsh, J., Moore, H., Andrews, P.W., 2007. Transient and stable transgene expression in human embryonic stem cells. *Stem cells* 25, 1521-1528.
- Litsiou, A., Hanson, S., Streit, A., 2005. A balance of FGF, BMP and WNT signalling positions the future placode territory in the head. *Development* 132, 4051-4062.
- Livak, K.J., Schmittgen, T.D., 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2⁻($\Delta\Delta C_T$) Method. *Methods* 25, 402-408.
- Lombardo, A., Isaacs, H.V., Slack, J.M., 1998. Expression and functions of FGF-3 in *Xenopus* development. *The International journal of developmental biology* 42, 1101-1107.
- Lovicu, F.J., McAvoy, J.W., 2001. FGF-induced lens cell proliferation and differentiation is dependent on MAPK (ERK1/2) signalling. *Development* 128, 5075-5084.
- Ludwig, T.E., Bergendahl, V., Levenstein, M.E., Yu, J., Probasco, M.D., Thomson, J.A., 2006. Feeder-independent culture of human embryonic stem cells. *Nature methods* 3, 637-646.
- Ludwig, T.E., Levenstein, M.E., Jones, J.M., Berggren, W.T., Mitchen, E.R., Frane, J.L., Crandall, L.J., Daigh, C.A., Conard, K.R., Piekarczyk, M.S., Llanas, R.A., Thomson, J.A., 2006. Derivation of human embryonic stem cells in defined conditions. *Nature biotechnology* 24, 185-187.
- Luo, T., Matsuo-Takasaki, M., Lim, J.H., Sargent, T.D., 2001. Differential regulation of *Dlx* gene expression by a BMP morphogenetic gradient. *The International journal of developmental biology* 45, 681-684.
- MacArthur, C.A., Shankar, D.B., Shackelford, G.M., 1995. *Fgf-8*, activated by proviral insertion, cooperates with the *Wnt-1* transgene in murine mammary tumorigenesis. *Journal of virology* 69, 2501-2507.
- Maher, P.A., 1996. Nuclear Translocation of fibroblast growth factor (FGF) receptors in response to FGF-2. *The Journal of cell biology* 134, 529-536.
- Malgrange, B., Lefebvre, P.P., Martin, D., Staecker, H., Van de Water, T.R., Moonen, G., 1996. NT-3 has a tropic effect on process outgrowth by postnatal auditory neurones in vitro. *Neuroreport* 7, 2495-2499.
- Mansukhani, A., Ambrosetti, D., Holmes, G., Cornivelli, L., Basilico, C., 2005. *Sox2* induction by FGF and FGFR2 activating mutations inhibits Wnt signaling and osteoblast differentiation. *The Journal of cell biology* 168, 1065-1076.
- Marks, H., Kalkan, T., Menafra, R., Denissov, S., Jones, K., Hofemeister, H., Nichols, J., Kranz, A., Stewart, A.F., Smith, A., Stunnenberg, H.G., 2012. The transcriptional and epigenomic foundations of ground state pluripotency. *Cell* 149, 590-604.

Maroon, H., Walshe, J., Mahmood, R., Kiefer, P., Dickson, C., Mason, I., 2002. Fgf3 and Fgf8 are required together for formation of the otic placode and vesicle. *Development* 129, 2099-2108.

Martin, G.R., 1981. Isolation of a pluripotent cell line from early mouse embryos cultured in medium conditioned by teratocarcinoma stem cells. *Proceedings of the National Academy of Sciences of the United States of America* 78, 7634-7638.

Martin, G.R., Evans, M.J., 1975. Differentiation of clonal lines of teratocarcinoma cells: formation of embryoid bodies in vitro. *Proceedings of the National Academy of Sciences of the United States of America* 72, 1441-1445.

Martin, K., Groves, A.K., 2006. Competence of cranial ectoderm to respond to Fgf signaling suggests a two-step model of otic placode induction. *Development* 133, 877-887.

Matsubara, A., Kan, M., Feng, S., McKeehan, W.L., 1998. Inhibition of growth of malignant rat prostate tumor cells by restoration of fibroblast growth factor receptor 2. *Cancer research* 58, 1509-1514.

McLarren, K.W., Litsiou, A., Streit, A., 2003. DLX5 positions the neural crest and preplacode region at the border of the neural plate. *Developmental biology* 259, 34-47.

Mehta, A., Mathew, S., Viswanathan, C., Sen Majumdar, A., 2010. Intrinsic properties and external factors determine the differentiation bias of human embryonic stem cell lines. *Cell biology international* 34, 1021-1031.

Meyers, E.N., Lewandoski, M., Martin, G.R., 1998. An Fgf8 mutant allelic series generated by Cre- and Flp-mediated recombination. *Nature genetics* 18, 136-141.

Miki, T., Bottaro, D.P., Fleming, T.P., Smith, C.L., Burgess, W.H., Chan, A.M., Aaronson, S.A., 1992. Determination of ligand-binding specificity by alternative splicing: two distinct growth factor receptors encoded by a single gene. *Proceedings of the National Academy of Sciences of the United States of America* 89, 246-250.

Miller, D.L., Ortega, S., Bashayan, O., Basch, R., Basilico, C., 2000. Compensation by fibroblast growth factor 1 (FGF1) does not account for the mild phenotypic defects observed in FGF2 null mice. *Molecular and cellular biology* 20, 2260-2268.

Minagawa, A., Iitaka, M., Suzuki, M., Yasuda, S., Kameyama, K., Shimada, S., Kitahama, S., Wada, S., Katayama, S., 2002. A case of primary mucoepidermoid carcinoma of the thyroid: molecular evidence of its origin. *Clinical endocrinology* 57, 551-556.

Miralles, F., Czernichow, P., Ozaki, K., Itoh, N., Scharfmann, R., 1999. Signaling through fibroblast growth factor receptor 2b plays a key role in the development of the exocrine pancreas. *Proceedings of the National Academy of Sciences of the United States of America* 96, 6267-6272.

- Miyazaki, J., Takaki, S., Araki, K., Tashiro, F., Tominaga, A., Takatsu, K., Yamamura, K., 1989. Expression vector system based on the chicken beta-actin promoter directs efficient production of interleukin-5. *Gene* 79, 269-277.
- Mohammadi, M., 1997. Structures of the Tyrosine Kinase Domain of Fibroblast Growth Factor Receptor in Complex with Inhibitors. *Science* 276, 955-960.
- Morisada, N., Rendtorff, N.D., Nozu, K., Morishita, T., Miyakawa, T., Matsumoto, T., Hisano, S., Iijima, K., Tranebjaerg, L., Shirahata, A., Matsuo, M., Kusahara, K., 2010. Branchio-otorenal syndrome caused by partial EYA1 deletion due to LINE-1 insertion. *Pediatric nephrology* 25, 1343-1348.
- Mueller, K.L., Jacques, B.E., Kelley, M.W., 2002. Fibroblast growth factor signaling regulates pillar cell development in the organ of corti. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 22, 9368-9377.
- Munoz-Sanjuan, I., Brivanlou, A.H., 2002. Neural induction, the default model and embryonic stem cells. *Nature reviews. Neuroscience* 3, 271-280.
- Murata, J., Ikeda, K., Okano, H., 2012. Notch signaling and the developing inner ear. *Advances in experimental medicine and biology* 727, 161-173.
- Na, J., Furue, M.K., Andrews, P.W., 2010. Inhibition of ERK1/2 prevents neural and mesendodermal differentiation and promotes human embryonic stem cell self-renewal. *Stem cell research* 5, 157-169.
- Nadol, J.B., Jr., 1997. Patterns of neural degeneration in the human cochlea and auditory nerve: implications for cochlear implantation. *Otolaryngology--head and neck surgery : official journal of American Academy of Otolaryngology-Head and Neck Surgery* 117, 220-228.
- Nadol, J.B., Jr., Eddington, D.K., 2006. Histopathology of the inner ear relevant to cochlear implantation. *Advances in oto-rhino-laryngology* 64, 31-49.
- Naimi, B., Latil, A., Fournier, G., Mangin, P., Cussenot, O., Berthon, P., 2002. Down-regulation of (IIIb) and (IIIc) isoforms of fibroblast growth factor receptor 2 (FGFR2) is associated with malignant progression in human prostate. *The Prostate* 52, 245-252.
- Naito, Y., Nakamura, T., Nakagawa, T., Iguchi, F., Endo, T., Fujino, K., Kim, T.S., Hiratsuka, Y., Tamura, T., Kanemaru, S., Shimizu, Y., Ito, J., 2004. Transplantation of bone marrow stromal cells into the cochlea of chinchillas. *Neuroreport* 15, 1-4.
- Naski, M.C., Colvin, J.S., Coffin, J.D., Ornitz, D.M., 1998. Repression of hedgehog signaling and BMP4 expression in growth plate cartilage by fibroblast growth factor receptor 3. *Development* 125, 4977-4988.

Nat, R., Nilbratt, M., Narkilahti, S., Winblad, B., Hovatta, O., Nordberg, A., 2007. Neurogenic neuroepithelial and radial glial cells generated from six human embryonic stem cell lines in serum-free suspension and adherent cultures. *Glia* 55, 385-399.

Nechiporuk, A., Linbo, T., Raible, D.W., 2005. Endoderm-derived Fgf3 is necessary and sufficient for inducing neurogenesis in the epibranchial placodes in zebrafish. *Development* 132, 3717-3730.

Nelson, A.D., Svendsen, C.N., 2006. Low concentrations of extracellular FGF-2 are sufficient but not essential for neurogenesis from human neural progenitor cells. *Molecular and cellular neurosciences* 33, 29-35.

Ng, K.T., Man, K., Sun, C.K., Lee, T.K., Poon, R.T., Lo, C.M., Fan, S.T., 2006. Clinicopathological significance of homeoprotein Six1 in hepatocellular carcinoma. *British journal of cancer* 95, 1050-1055.

Niwa, H., Miyazaki, J., Smith, A.G., 2000. Quantitative expression of Oct-3/4 defines differentiation, dedifferentiation or self-renewal of ES cells. *Nature genetics* 24, 372-376.

Niwa, H., Yamamura, K., Miyazaki, J., 1991. Efficient selection for high-expression transfectants with a novel eukaryotic vector. *Gene* 108, 193-199.

Nornes, H.O., Dressler, G.R., Knapik, E.W., Deutsch, U., Gruss, P., 1990. Spatially and temporally restricted expression of Pax2 during murine neurogenesis. *Development* 109, 797-809.

Ohyama, T., Groves, A.K., 2004. Expression of mouse Foxi class genes in early craniofacial development. *Developmental dynamics : an official publication of the American Association of Anatomists* 231, 640-646.

Ohyama, T., Groves, A.K., 2004. Generation of Pax2-Cre mice by modification of a Pax2 bacterial artificial chromosome. *Genesis* 38, 195-199.

Ohyama, T., Groves, A.K., Martin, K., 2007. The first steps towards hearing: mechanisms of otic placode induction. *The International journal of developmental biology* 51, 463-472.

Ornitz, D.M., 2005. FGF signaling in the developing endochondral skeleton. *Cytokine & growth factor reviews* 16, 205-213.

Ornitz, D.M., Xu, J., Colvin, J.S., McEwen, D.G., MacArthur, C.A., Coulier, F., Gao, G., Goldfarb, M., 1996. Receptor specificity of the fibroblast growth factor family. *The Journal of biological chemistry* 271, 15292-15297.

Ornitz, D.M., Yayon, A., Flanagan, J.G., Svahn, C.M., Levi, E., Leder, P., 1992. Heparin is required for cell-free binding of basic fibroblast growth factor to a soluble receptor and for mitogenesis in whole cells. *Molecular and cellular biology* 12, 240-247.

Orr-Urtreger, A., Bedford, M.T., Burakova, T., Arman, E., Zimmer, Y., Yayon, A., Givol, D., Lonai, P., 1993. Developmental localization of the splicing alternatives of fibroblast growth factor receptor-2 (FGFR2). *Developmental biology* 158, 475-486.

Osafune, K., Caron, L., Borowiak, M., Martinez, R.J., Fitz-Gerald, C.S., Sato, Y., Cowan, C.A., Chien, K.R., Melton, D.A., 2008. Marked differences in differentiation propensity among human embryonic stem cell lines. *Nature biotechnology* 26, 313-315.

Oshima, K., Shin, K., Diensthuber, M., Peng, A.W., Ricci, A.J., Heller, S., 2010. Mechanosensitive hair cell-like cells from embryonic and induced pluripotent stem cells. *Cell* 141, 704-716.

Ouji, Y., Ishizaka, S., Nakamura-Uchiyama, F., Yoshikawa, M., 2012. In vitro differentiation of mouse embryonic stem cells into inner ear hair cell-like cells using stromal cell conditioned medium. *Cell death & disease* 3, e314.

Paddison, P.J., Caudy, A.A., Bernstein, E., Hannon, G.J., Conklin, D.S., 2002. Short hairpin RNAs (shRNAs) induce sequence-specific silencing in mammalian cells. *Genes & development* 16, 948-958.

Pandit, S.R., Sullivan, J.M., Egger, V., Borecki, A.A., Oleskevich, S., 2011. Functional effects of adult human olfactory stem cells on early-onset sensorineural hearing loss. *Stem cells* 29, 670-677.

Pauley, S., Lai, E., Fritzschn, B., 2006. Foxg1 is required for morphogenesis and histogenesis of the mammalian inner ear. *Developmental dynamics : an official publication of the American Association of Anatomists* 235, 2470-2482.

Pera, E.M., Wessely, O., Li, S.Y., De Robertis, E.M., 2001. Neural and head induction by insulin-like growth factor signals. *Developmental cell* 1, 655-665.

Peters, K.G., Werner, S., Chen, G., Williams, L.T., 1992. Two FGF receptor genes are differentially expressed in epithelial and mesenchymal tissues during limb formation and organogenesis in the mouse. *Development* 114, 233-243.

Pfeffer, P.L., Gerster, T., Lun, K., Brand, M., Busslinger, M., 1998. Characterization of three novel members of the zebrafish Pax2/5/8 family: dependency of Pax5 and Pax8 expression on the Pax2.1 (noi) function. *Development* 125, 3063-3074.

Phillips, B.T., Bolding, K., Riley, B.B., 2001. Zebrafish fgf3 and fgf8 encode redundant functions required for otic placode induction. *Developmental biology* 235, 351-365.

Pirvola, U., Spencer-Dene, B., Xing-Qun, L., Kettunen, P., Thesleff, I., Fritzschn, B., Dickson, C., Ylikoski, J., 2000. FGF/FGFR-2(IIIb) signaling is essential for inner ear morphogenesis. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 20, 6125-6134.

- Pirvola, U., Ylikoski, J., Trokovic, R., Hebert, J.M., McConnell, S.K., Partanen, J., 2002. FGFR1 is required for the development of the auditory sensory epithelium. *Neuron* 35, 671-680.
- Portnoy, V., Huang, V., Place, R.F., Li, L.C., 2011. Small RNA and transcriptional upregulation. *Wiley interdisciplinary reviews. RNA* 2, 748-760.
- Presta, M., Belleri, M., Vecchi, A., Hesselgesser, J., Mantovani, A., Horuk, R., 1998. Noncompetitive, chemokine-mediated inhibition of basic fibroblast growth factor-induced endothelial cell proliferation. *The Journal of biological chemistry* 273, 7911-7919.
- Puschel, A.W., Westerfield, M., Dressler, G.R., 1992. Comparative analysis of Pax-2 protein distributions during neurulation in mice and zebrafish. *Mechanisms of development* 38, 197-208.
- Qian, X., Davis, A.A., Goderie, S.K., Temple, S., 1997. FGF2 concentration regulates the generation of neurons and glia from multipotent cortical stem cells. *Neuron* 18, 81-93.
- Quint, E., Zerucha, T., Ekker, M., 2000. Differential expression of orthologous Dlx genes in zebrafish and mice: implications for the evolution of the Dlx homeobox gene family. *The Journal of experimental zoology* 288, 235-241.
- Reardon, W., Winter, R.M., Rutland, P., Pulleyn, L.J., Jones, B.M., Malcolm, S., 1994. Mutations in the fibroblast growth factor receptor 2 gene cause Crouzon syndrome. *Nature genetics* 8, 98-103.
- Reiter, A., Sohal, J., Kulkarni, S., Chase, A., Macdonald, D.H., Aguiar, R.C., Goncalves, C., Hernandez, J.M., Jennings, B.A., Goldman, J.M., Cross, N.C., 1998. Consistent fusion of ZNF198 to the fibroblast growth factor receptor-1 in the t(8;13)(p11;q12) myeloproliferative syndrome. *Blood* 92, 1735-1742.
- Revoltella, R.P., Papini, S., Rosellini, A., Michelini, M., Franceschini, V., Ciorba, A., Bertolaso, L., Magosso, S., Hatzopoulos, S., Lorito, G., Giordano, P., Simoni, E., Ognio, E., Cilli, M., Saccardi, R., Urbani, S., Jeffery, R., Poulson, R., Martini, A., 2008. Cochlear repair by transplantation of human cord blood CD133+ cells to nod-scid mice made deaf with kanamycin and noise. *Cell transplantation* 17, 665-678.
- Ricol, D., Cappellen, D., El Marjou, A., Gil-Diez-de-Medina, S., Girault, J.M., Yoshida, T., Ferry, G., Tucker, G., Poupon, M.F., Chopin, D., Thiery, J.P., Radvanyi, F., 1999. Tumour suppressive properties of fibroblast growth factor receptor 2-IIIb in human bladder cancer. *Oncogene* 18, 7234-7243.
- Robledo, R.F., Lufkin, T., 2006. Dlx5 and Dlx6 homeobox genes are required for specification of the mammalian vestibular apparatus. *Genesis* 44, 425-437.

Roghani, M., Moscatelli, D., 1992. Basic fibroblast growth factor is internalized through both receptor-mediated and heparan sulfate-mediated mechanisms. *The Journal of biological chemistry* 267, 22156-22162.

Ronaghi, M., Nasr, M., Heller, S., 2012. Concise review: Inner ear stem cells--an oxymoron, but why? *Stem cells* 30, 69-74.

Roumiantsev, S., Krause, D.S., Neumann, C.A., Dimitri, C.A., Asiedu, F., Cross, N.C., Van Etten, R.A., 2004. Distinct stem cell myeloproliferative/T lymphoma syndromes induced by ZNF198-FGFR1 and BCR-FGFR1 fusion genes from 8p11 translocations. *Cancer cell* 5, 287-298.

Rousseau, F., Bonaventure, J., Legeai-Mallet, L., Pelet, A., Rozet, J.M., Maroteaux, P., Le Merrer, M., Munnich, A., 1994. Mutations in the gene encoding fibroblast growth factor receptor-3 in achondroplasia. *Nature* 371, 252-254.

Sahni, M., Ambrosetti, D.C., Mansukhani, A., Gertner, R., Levy, D., Basilico, C., 1999. FGF signaling inhibits chondrocyte proliferation and regulates bone development through the STAT-1 pathway. *Genes & development* 13, 1361-1366.

Sakamoto, T., Nakagawa, T., Endo, T., Kim, T.S., Iguchi, F., Naito, Y., Sasai, Y., Ito, J., 2004. Fates of mouse embryonic stem cells transplanted into the inner ears of adult mice and embryonic chickens. *Acta oto-laryngologica. Supplementum*, 48-52.

Sato, N., Meijer, L., Skaltsounis, L., Greengard, P., Brivanlou, A.H., 2004. Maintenance of pluripotency in human and mouse embryonic stem cells through activation of Wnt signaling by a pharmacological GSK-3-specific inhibitor. *Nature medicine* 10, 55-63.

Sato, N., Sanjuan, I.M., Heke, M., Uchida, M., Naef, F., Brivanlou, A.H., 2003. Molecular signature of human embryonic stem cells and its comparison with the mouse. *Developmental biology* 260, 404-413.

Schimmang, T., 2007. Expression and functions of FGF ligands during early otic development. *The International journal of developmental biology* 51, 473-481.

Schmahl, J., Kim, Y., Colvin, J.S., Ornitz, D.M., Capel, B., 2004. Fgf9 induces proliferation and nuclear localization of FGFR2 in Sertoli precursors during male sex determination. *Development* 131, 3627-3636.

Schuldiner, M., Yanuka, O., Itskovitz-Eldor, J., Melton, D.A., Benvenisty, N., 2000. Effects of eight growth factors on the differentiation of cells derived from human embryonic stem cells. *Proceedings of the National Academy of Sciences of the United States of America* 97, 11307-11312.

Schule, B., Li, H.H., Fisch-Kohl, C., Purmann, C., Francke, U., 2007. DLX5 and DLX6 expression is biallelic and not modulated by MeCP2 deficiency. *American journal of human genetics* 81, 492-506.

Schwartz, M.S., Otto, S.R., Shannon, R.V., Hitselberger, W.E., Brackmann, D.E., 2008. Auditory brainstem implants. *Neurotherapeutics : the journal of the American Society for Experimental NeuroTherapeutics* 5, 128-136.

Shackleford, G.M., MacArthur, C.A., Kwan, H.C., Varmus, H.E., 1993. Mouse mammary tumor virus infection accelerates mammary carcinogenesis in Wnt-1 transgenic mice by insertional activation of int-2/Fgf-3 and hst/Fgf-4. *Proceedings of the National Academy of Sciences of the United States of America* 90, 740-744.

Sharp, P.A., 2001. RNA interference--2001. *Genes & development* 15, 485-490.

Shi, E., Kan, M., Xu, J., Wang, F., Hou, J., McKeehan, W.L., 1993. Control of fibroblast growth factor receptor kinase signal transduction by heterodimerization of combinatorial splice variants. *Molecular and cellular biology* 13, 3907-3918.

Shi, F., Corrales, C.E., Liberman, M.C., Edge, A.S., 2007. BMP4 induction of sensory neurons from human embryonic stem cells and reinnervation of sensory epithelium. *The European journal of neuroscience* 26, 3016-3023.

Shiang, R., Thompson, L.M., Zhu, Y.Z., Church, D.M., Fielder, T.J., Bocian, M., Winokur, S.T., Wasmuth, J.J., 1994. Mutations in the transmembrane domain of FGFR3 cause the most common genetic form of dwarfism, achondroplasia. *Cell* 78, 335-342.

Shibata, S.B., Cortez, S.R., Beyer, L.A., Wiler, J.A., Di Polo, A., Pflingst, B.E., Raphael, Y., 2010. Transgenic BDNF induces nerve fiber regrowth into the auditory epithelium in deaf cochleae. *Experimental neurology* 223, 464-472.

Sperger, J.M., Chen, X., Draper, J.S., Antosiewicz, J.E., Chon, C.H., Jones, S.B., Brooks, J.D., Andrews, P.W., Brown, P.O., Thomson, J.A., 2003. Gene expression patterns in human embryonic stem cells and human pluripotent germ cell tumors. *Proceedings of the National Academy of Sciences of the United States of America* 100, 13350-13355.

Spoendlin, H., 1975. Retrograde degeneration of the cochlear nerve. *Acta oto-laryngologica* 79, 266-275.

Staecker, H., Kopke, R., Malgrange, B., Lefebvre, P., Van de Water, T.R., 1996. NT-3 and/or BDNF therapy prevents loss of auditory neurons following loss of hair cells. *Neuroreport* 7, 889-894.

Stavridis, M.P., Lunn, J.S., Collins, B.J., Storey, K.G., 2007. A discrete period of FGF-induced Erk1/2 signalling is required for vertebrate neural specification. *Development* 134, 2889-2894.

Steele, I.A., Edmondson, R.J., Bulmer, J.N., Bolger, B.S., Leung, H.Y., Davies, B.R., 2001. Induction of FGF receptor 2-IIIb expression and response to its ligands in epithelial ovarian cancer. *Oncogene* 20, 5878-5887.

Streit, A., 2002. Extensive cell movements accompany formation of the otic placode. *Developmental biology* 249, 237-254.

Streit, A., 2004. Early development of the cranial sensory nervous system: from a common field to individual placodes. *Developmental biology* 276, 1-15.

Streit, A., 2007. The preplacodal region: an ectodermal domain with multipotential progenitors that contribute to sense organs and cranial sensory ganglia. *The International journal of developmental biology* 51, 447-461.

Streit, A., Stern, C.D., 1999. Establishment and maintenance of the border of the neural plate in the chick: involvement of FGF and BMP activity. *Mechanisms of development* 82, 51-66.

Tavassoli, K., Ruger, W., Horst, J., 1997. Alternative splicing in PAX2 generates a new reading frame and an extended conserved coding region at the carboxy terminus. *Human genetics* 101, 371-375.

Tesar, P.J., Chenoweth, J.G., Brook, F.A., Davies, T.J., Evans, E.P., Mack, D.L., Gardner, R.L., McKay, R.D., 2007. New cell lines from mouse epiblast share defining features with human embryonic stem cells. *Nature* 448, 196-199.

Theodorou, V., Boer, M., Weigelt, B., Jonkers, J., van der Valk, M., Hilkens, J., 2004. Fgf10 is an oncogene activated by MMTV insertional mutagenesis in mouse mammary tumors and overexpressed in a subset of human breast carcinomas. *Oncogene* 23, 6047-6055.

Thomson, J.A., Itskovitz-Eldor, J., Shapiro, S.S., Waknitz, M.A., Swiergiel, J.J., Marshall, V.S., Jones, J.M., 1998. Embryonic stem cell lines derived from human blastocysts. *Science* 282, 1145-1147.

Torres, M., Gomez-Pardo, E., Gruss, P., 1996. Pax2 contributes to inner ear patterning and optic nerve trajectory. *Development* 122, 3381-3391.

Tropepe, V., Hitoshi, S., Sirard, C., Mak, T.W., Rossant, J., van der Kooy, D., 2001. Direct neural fate specification from embryonic stem cells: a primitive mammalian neural stem cell stage acquired through a default mechanism. *Neuron* 30, 65-78.

Turner, C.W., Gantz, B.J., Karsten, S., Fowler, J., Reiss, L.A., 2010. Impact of hair cell preservation in cochlear implantation: combined electric and acoustic hearing. *Otology & neurotology : official publication of the American Otological Society, American Neurotology Society [and] European Academy of Otology and Neurotology* 31, 1227-1232.

- Tuschl, T., Zamore, P.D., Lehmann, R., Bartel, D.P., Sharp, P.A., 1999. Targeted mRNA degradation by double-stranded RNA in vitro. *Genes & development* 13, 3191-3197.
- Vallier, L., Alexander, M., Pedersen, R.A., 2005. Activin/Nodal and FGF pathways cooperate to maintain pluripotency of human embryonic stem cells. *Journal of cell science* 118, 4495-4509.
- Vallier, L., Touboul, T., Chng, Z., Brimpari, M., Hannan, N., Millan, E., Smithers, L.E., Trotter, M., Rugg-Gunn, P., Weber, A., Pedersen, R.A., 2009. Early cell fate decisions of human embryonic stem cells and mouse epiblast stem cells are controlled by the same signalling pathways. *PLoS one* 4, e6082.
- Wang, B., Gao, Y., Xiao, Z., Chen, B., Han, J., Zhang, J., Wang, X., Dai, J., 2009. Erk1/2 promotes proliferation and inhibits neuronal differentiation of neural stem cells. *Neuroscience letters* 461, 252-257.
- Warchol, M.E., Lambert, P.R., Goldstein, B.J., Forge, A., Corwin, J.T., 1993. Regenerative proliferation in inner ear sensory epithelia from adult guinea pigs and humans. *Science* 259, 1619-1622.
- Watanabe, K., Ueno, M., Kamiya, D., Nishiyama, A., Matsumura, M., Wataya, T., Takahashi, J.B., Nishikawa, S., Muguruma, K., Sasai, Y., 2007. A ROCK inhibitor permits survival of dissociated human embryonic stem cells. *Nature biotechnology* 25, 681-686.
- Webster, M.K., Donoghue, D.J., 1997. Enhanced signaling and morphological transformation by a membrane-localized derivative of the fibroblast growth factor receptor 3 kinase domain. *Molecular and cellular biology* 17, 5739-5747.
- Weinstein, M., Xu, X., Ohyama, K., Deng, C.X., 1998. FGFR-3 and FGFR-4 function cooperatively to direct alveogenesis in the murine lung. *Development* 125, 3615-3623.
- Wernig, M., Lengner, C.J., Hanna, J., Lodato, M.A., Steine, E., Foreman, R., Staerk, J., Markoulaki, S., Jaenisch, R., 2008. A drug-inducible transgenic system for direct reprogramming of multiple somatic cell types. *Nature biotechnology* 26, 916-924.
- White, P.M., Doetzlhofer, A., Lee, Y.S., Groves, A.K., Segil, N., 2006. Mammalian cochlear supporting cells can divide and trans-differentiate into hair cells. *Nature* 441, 984-987.
- Wilkie, A.O., Slaney, S.F., Oldridge, M., Poole, M.D., Ashworth, G.J., Hockley, A.D., Hayward, R.D., David, D.J., Pulleyn, L.J., Rutland, P., et al., 1995. Apert syndrome results from localized mutations of FGFR2 and is allelic with Crouzon syndrome. *Nature genetics* 9, 165-172.
- Wilson, B.S., Dorman, M.F., 2008. Cochlear implants: a remarkable past and a brilliant future. *Hearing research* 242, 3-21.

- Wilson, S.I., Rydstrom, A., Trimborn, T., Willert, K., Nusse, R., Jessell, T.M., Edlund, T., 2001. The status of Wnt signalling regulates neural and epidermal fates in the chick embryo. *Nature* 411, 325-330.
- Wise, A.K., Hume, C.R., Flynn, B.O., Jeelall, Y.S., Suhr, C.L., Sgro, B.E., O'Leary, S.J., Shepherd, R.K., Richardson, R.T., 2010. Effects of localized neurotrophin gene expression on spiral ganglion neuron resprouting in the deafened cochlea. *Molecular therapy : the journal of the American Society of Gene Therapy* 18, 1111-1122.
- Wise, A.K., Richardson, R., Hardman, J., Clark, G., O'Leary, S., 2005. Resprouting and survival of guinea pig cochlear neurons in response to the administration of the neurotrophins brain-derived neurotrophic factor and neurotrophin-3. *The Journal of comparative neurology* 487, 147-165.
- Woda, J.M., Pastagia, J., Mercola, M., Artinger, K.B., 2003. Dlx proteins position the neural plate border and determine adjacent cell fates. *Development* 130, 331-342.
- Wright, T.J., Hatch, E.P., Karabagli, H., Karabagli, P., Schoenwolf, G.C., Mansour, S.L., 2003. Expression of mouse fibroblast growth factor and fibroblast growth factor receptor genes during early inner ear development. *Developmental dynamics : an official publication of the American Association of Anatomists* 228, 267-272.
- Wright, T.J., Ladher, R., McWhirter, J., Murre, C., Schoenwolf, G.C., Mansour, S.L., 2004. Mouse FGF15 is the ortholog of human and chick FGF19, but is not uniquely required for otic induction. *Developmental biology* 269, 264-275.
- Wright, T.J., Mansour, S.L., 2003. Fgf3 and Fgf10 are required for mouse otic placode induction. *Development* 130, 3379-3390.
- Wu, H., Xu, J., Pang, Z.P., Ge, W., Kim, K.J., Bianchi, B., Chen, C., Sudhof, T.C., Sun, Y.E., 2007. Integrative genomic and functional analyses reveal neuronal subtype differentiation bias in human embryonic stem cell lines. *Proceedings of the National Academy of Sciences of the United States of America* 104, 13821-13826.
- Xia, X., Zhang, Y., Zieth, C.R., Zhang, S.C., 2007. Transgenes delivered by lentiviral vector are suppressed in human embryonic stem cells in a promoter-dependent manner. *Stem cells and development* 16, 167-176.
- Xu, R.H., Chen, X., Li, D.S., Li, R., Addicks, G.C., Glennon, C., Zwaka, T.P., Thomson, J.A., 2002. BMP4 initiates human embryonic stem cell differentiation to trophoblast. *Nature biotechnology* 20, 1261-1264.
- Xu, R.H., Peck, R.M., Li, D.S., Feng, X., Ludwig, T., Thomson, J.A., 2005. Basic FGF and suppression of BMP signaling sustain undifferentiated proliferation of human ES cells. *Nature methods* 2, 185-190.

- Xu, X., Weinstein, M., Li, C., Naski, M., Cohen, R.I., Ornitz, D.M., Leder, P., Deng, C., 1998. Fibroblast growth factor receptor 2 (FGFR2)-mediated reciprocal regulation loop between FGF8 and FGF10 is essential for limb induction. *Development* 125, 753-765.
- Yamaguchi, T.P., Harpal, K., Henkemeyer, M., Rossant, J., 1994. *fgfr-1* is required for embryonic growth and mesodermal patterning during mouse gastrulation. *Genes & development* 8, 3032-3044.
- Yan, G., Fukabori, Y., McBride, G., Nikolaropolous, S., McKeehan, W.L., 1993. Exon switching and activation of stromal and embryonic fibroblast growth factor (FGF)-FGF receptor genes in prostate epithelial cells accompany stromal independence and malignancy. *Molecular and cellular biology* 13, 4513-4522.
- Yang, L., Zhang, H., Hu, G., Wang, H., Abate-Shen, C., Shen, M.M., 1998. An early phase of embryonic *Dlx5* expression defines the rostral boundary of the neural plate. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 18, 8322-8330.
- Yang, S.M., Chen, W., Guo, W.W., Jia, S., Sun, J.H., Liu, H.Z., Young, W.Y., He, D.Z., 2012. Regeneration of stereocilia of hair cells by forced *Atoh1* expression in the adult mammalian cochlea. *PloS one* 7, e46355.
- Yao, S., Chen, S., Clark, J., Hao, E., Beattie, G.M., Hayek, A., Ding, S., 2006. Long-term self-renewal and directed differentiation of human embryonic stem cells in chemically defined conditions. *Proceedings of the National Academy of Sciences of the United States of America* 103, 6907-6912.
- Ying, Q.L., Nichols, J., Chambers, I., Smith, A., 2003. BMP induction of *Id* proteins suppresses differentiation and sustains embryonic stem cell self-renewal in collaboration with STAT3. *Cell* 115, 281-292.
- Ying, Q.L., Stavridis, M., Griffiths, D., Li, M., Smith, A., 2003. Conversion of embryonic stem cells into neuroectodermal precursors in adherent monoculture. *Nature biotechnology* 21, 183-186.
- Ylikoski, J., Pirvola, U., Moshnyakov, M., Palgi, J., Arumae, U., Saarma, M., 1993. Expression patterns of neurotrophin and their receptor mRNAs in the rat inner ear. *Hearing research* 65, 69-78.
- Yoshida, T., Hakuba, N., Morizane, I., Fujita, K., Cao, F., Zhu, P., Uchida, N., Kameda, K., Sakanaka, M., Gyo, K., Hata, R., 2007. Hematopoietic stem cells prevent hair cell death after transient cochlear ischemia through paracrine effects. *Neuroscience* 145, 923-930.
- Yu, J.Y., DeRuiter, S.L., Turner, D.L., 2002. RNA interference by expression of short-interfering RNAs and hairpin RNAs in mammalian cells. *Proceedings of the National Academy of Sciences of the United States of America* 99, 6047-6052.

Yu, P., Pan, G., Yu, J., Thomson, J.A., 2011. FGF2 sustains NANOG and switches the outcome of BMP4-induced human embryonic stem cell differentiation. *Cell stem cell* 8, 326-334.

Zafarana, G., Avery, S.R., Avery, K., Moore, H.D., Andrews, P.W., 2009. Specific knockdown of OCT4 in human embryonic stem cells by inducible short hairpin RNA interference. *Stem cells* 27, 776-782.

Zelarayan, L.C., Vendrell, V., Alvarez, Y., Dominguez-Frutos, E., Theil, T., Alonso, M.T., Maconochie, M., Schimmang, T., 2007. Differential requirements for FGF3, FGF8 and FGF10 during inner ear development. *Developmental biology* 308, 379-391.

Zhang, X., Ibrahimi, O.A., Olsen, S.K., Umemori, H., Mohammadi, M., Ornitz, D.M., 2006. Receptor specificity of the fibroblast growth factor family. The complete mammalian FGF family. *The Journal of biological chemistry* 281, 15694-15700.

Zheng, J.L., Gao, W.Q., 2000. Overexpression of Math1 induces robust production of extra hair cells in postnatal rat inner ears. *Nature neuroscience* 3, 580-586.

Zwaka, T.P., Thomson, J.A., 2003. Homologous recombination in human embryonic stem cells. *Nature biotechnology* 21, 319-321.